

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 May 2006 (26.05.2006)

PCT

(10) International Publication Number
WO 2006/055760 A1

(51) International Patent Classification:

C07D 249/12 (2006.01) C07D 403/04 (2006.01)
C07D 401/04 (2006.01) A61K 31/4196 (2006.01)
C07D 405/04 (2006.01) A61P 35/00 (2006.01)

(21) International Application Number:

PCT/US2005/041779

(22) International Filing Date:

17 November 2005 (17.11.2005)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/628,979 18 November 2004 (18.11.2004) US
60/709,358 18 August 2005 (18.08.2005) US
60/725,044 6 October 2005 (06.10.2005) US

(71) Applicant (for all designated States except US): **SYNTA PHARMACEUTICALS CORP.** [US/US]; 45 Hartwell Avenue, Lexington, Massachusetts 02421 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **YING, Weiwen** [CN/US]; 55 Littleton Road, Unit 29b, Ayer, Massachusetts 01432 (US). **JAMES, David** [AU/US]; 229 Western Avenue, No. 2, Cambridge, Massachusetts 02139 (US). **ZHANG, Shijie** [CN/US]; 27 Lojko Drive, Nashua, New Hampshire 03062 (US). **PRZEWLOKA, Teresa** [PL/US]; 23 Merrimack Meadows Lane, Tewksbury, Massachusetts 01876 (US). **CHAE, Junghyun** [KR/US]; 3 Brattle Drive, No. 10, Arlington, Massachusetts 02474 (US). **CHIM-MANAMADA, Dinesh, U.** [IN/US]; 26-B Liberty Street, Waltham, Massachusetts 02452 (US). **LEE, Chi-Wan** [KR/US]; 240 Magill Drive, Grafton, Massachusetts 01519 (US). **KOSTIK, Elena** [RU/US]; 10 Park Terrace B, Arlington, Massachusetts 02474 (US). **NG, Howard, P.** [US/US]; 17 Worcester Street, No. 1, Belmont, Massachusetts 02478 (US). **FOLEY, Kevin** [US/US]; 30 Cambridge

Park Drive, No. 4109, Cambridge, Massachusetts 02140 (US). **DU, Zhenjian** [US/US]; 18 Overlock Drive, Northborough, Massachusetts 01532 (US). **BARSOUM, James** [US/US]; 6 Moreland Avenue, Lexington, Massachusetts 02421 (US).

(74) Agents: **DAVID, Steven, G.** et al.; HAMILTON, BROOK, SMITH & REYNOLDS, P.C., 530 VIRGINIA ROAD, P.O. Box 9133, Concord, Massachusetts 01742-9133 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GI, GM, GR, GU, HN, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: TRIAZOLE COMPOUNDS THAT MODULATE HSP90 ACTIVITY

(57) Abstract: The present invention relates to substituted triazole compounds and compositions comprising substituted triazole compounds. The invention further relates to methods of inhibiting the activity of Hsp90 in a subject in need thereof and methods for preventing or treating hyperproliferative disorders, such as cancer, in a subject in need thereof comprising administering to the subject a substituted triazole compound of the invention, or a composition comprising such a compound.



WO 2006/055760 A1

TRIAZOLE COMPOUNDS THAT MODULATE HSP90 ACTIVITY

RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application No. 60/628,979, filed November 18, 2004; U.S. Provisional Application No. 60/709,358, filed August 18, 2005, and U.S. Provisional Application No. 60/725,044, filed October 6, 2005. The entire teachings of the above applications are incorporated herein by reference.

BACKGROUND OF THE INVENTION

Although tremendous advances have been made in elucidating the genomic abnormalities that cause malignant cancer cells, currently available chemotherapy remains unsatisfactory, and the prognosis for the majority of patients diagnosed with cancer remains dismal. Most chemotherapeutic agents act on a specific molecular target thought to be involved in the development of the malignant phenotype. However, a complex network of signaling pathways regulate cell proliferation, and the majority of malignant cancers are facilitated by multiple genetic abnormalities in these pathway. Therefore, it is unlikely that a therapeutic agent that acts on one molecular target will be fully effective in curing a patient who has cancer.

Heat shock proteins (HSPs) are a class of chaperone proteins that are up-regulated in response to elevated temperature and other environmental stresses, such as ultraviolet light, nutrient deprivation, and oxygen deprivation. HSPs act as chaperones to other cellular proteins (called client proteins) and facilitate their proper folding and repair, and aid in the refolding of misfolded client proteins. There are several known families of HSPs, each having its own set of client proteins. The Hsp90 family is one of the most abundant HSP families, accounting for about 1-2% of proteins in a cell that is not under stress and increasing to about 4-6% in a cell under stress. Inhibition of Hsp90 results in degradation of its client proteins via the ubiquitin proteasome pathway. Unlike other chaperone proteins, the client proteins of Hsp90 are mostly protein kinases or transcription factors involved in signal

transduction, and a number of its client proteins have been shown to be involved in the progression of cancer. Examples of Hsp90 client proteins that have been implicated in the progression of cancer are described below.

Her-2 is a transmembrane tyrosine kinase cell surface growth factor receptor
5 that is expressed in normal epithelial cells. Her2 has an extracellular domain that interacts with extracellular growth factors and an internal tyrosine kinase portion that transmits the external growth signal to the nucleus of the cell. Her2 is overexpressed in a significant proportion of malignancies, such as breast cancer, ovarian cancer, prostate cancer, and gastric cancers, and is typically associated with
10 a poor prognosis.

c-Kit is a membrane receptor protein tyrosine kinase which binds Stem Cell Factor (SCF) to its extraellular domain. c-Kit is involved in the development of melanocytes, mast, germ and hematopoietic cells, and there is evidence that it plays a role in several types of cancer including leukemias, mast cell tumors, small cell
15 lung cancer, testicular cancer, cancers of the gastointesinal tract and cancers of the central nervous system.

c-Met is a receptor tyrosine kinase that is encoded by the Met protooncogene and transduces the biological effects of hepatocyte growth factor (HGF), which is also referred to as scatter factor (SF). Jiang *et al.*, *Crit. Rev. Oncol. Hemtol.* 29: 209-
20 248 (1999), the entire teachings of which are incorporated herein by reference. c-Met and HGF are expressed in numerous tissues, although their expression is normally confined predominantly to cells of epithelial and mesenchymal origin, respectively. c-Met and HGF are required for normal mammalian development and have been shown to be important in cell migration, cell proliferation and survival,
25 morphogenic differentiation, and organization of 3-dimensional tubular structures (e.g., renal tubular cells, gland formation, etc.). The c-Met receptor has been shown to be expressed in a number of human cancers. c-Met and its ligand, HGF, have also been shown to be co-expressed at elevated levels in a variety of human cancers (particularly sarcomas). However, because the receptor and ligand are usually
30 expressed by different cell types, c-Met signaling is most commonly regulated by tumor-stroma (tumor-host) interactions. Furthermore, c-Met gene amplification, mutation, and rearrangement have been observed in a subset of human cancers.

Families with germline mutations that activate c-Met kinase are prone to multiple kidney tumors as well as tumors in other tissues. Numerous studies have correlated the expression of c-Met and/or HGF/SF with the state of disease progression of different types of cancer (including lung, colon, breast, prostate, liver, pancreas, 5 brain, kidney, ovaries, stomach, skin, and bone cancers). Furthermore, the overexpression of c-Met or HGF have been shown to correlate with poor prognosis and disease outcome in a number of major human cancers including lung, liver, gastric, and breast.

Akt kinase is a serine/threonine kinase which is a downstream effector 10 molecule of phosphoinositide 3-kinase and is involved in protecting the cell from apoptosis. Akt kinase is thought to be involved in the progression of cancer because it stimulates cell proliferation and suppresses apoptosis.

Cdk4/cyclin D complexes are involved in phosphorylation of retinoblastoma protein which is an essential step in progression of a cell through the G1 phase of the 15 cell cycle. Disruption of Hsp90 activity has been shown to decrease the half life of newly synthesized Cdk4.

Raf-1 is a MAP 3-kinase (MAP3K) which when activated can phosphorylate and activate the serine/threonine specific protein kinases ERK1 and ERK2. Activated ERKs play an important role in the control of gene expression involved in 20 the cell division cycle, apoptosis, cell differentiation and cell migration.

The transforming protein of Rous sarcoma virus, v-src, is a prototype of an oncogene family that induces cellular transformation (i.e., tumorigenesis) by non-regulated kinase activity. Hsp90 has been shown to complex with v-src and inhibit its degradation.

25 The BCR-ABL fusion protein associated with chronic myelogenous leukemia and in a subset of patients with acute lymphoblastic leukemia. The fusion protein is a consequence of exchange of genetic material from the long arms of chromosomes 9 and 22 and results in unregulated tyrosine kinase activity. BCR-ABL exists as a complex with Hsp90 and is rapidly degraded when the action of 30 Hsp90 is inhibited.

Hsp90 is required to maintain steroid hormone receptors in a conformation capable of binding hormone with high affinity. Inhibition of the action of Hsp90

therefore is expected to be useful in treating hormone-associated malignancies such as breast cancer.

p53 is a tumor suppressor protein that causes cell cycle arrest and apoptosis. Mutation of the p53 gene is found in about half of all human cancers making it one of the most common genetic alterations found in cancerous cells. In addition, p53 mutation is associated with a poor prognosis. Wild-type p53 has been shown to interact with Hsp90, but mutated p53 forms a more stable association than wild-type p53 as a result of its misfolded conformations. A stronger interaction with Hsp90 protects the mutated protein from normal proteolytic degradation and prolongs its half-life. In a cell that is heterozygous for mutated and wild-type p53, inhibition of the stabilizing effect of Hsp90 causes mutant p53 to be degraded and restores the normal transcriptional activity of wild-type p53.

Hif-1 α is a hypoxia-inducible transcription factor that is up-regulated under low oxygen conditions. Under normal oxygen conditions Hif-1 α associates with Von Hippel-Lindau (VHL) tumor suppressor protein and is degraded. Low oxygen conditions inhibits this association and allows Hif-1 α to accumulate and complex with Hif-1 β to form an active transcription complex that associates with hypoxia-response elements to activate the transcription of vascular endothelial growth factor (VEGF). Increased Hif-1 α is associated with increased metastasis and a poor prognosis.

Hsp90 has been shown by mutational analysis to be necessary for the survival of normal eukaryotic cells. However, Hsp90 is over expressed in many tumor types indicating that it may play a significant role in the survival of cancer cells and that cancer cells may be more sensitive to inhibition of Hsp90 than normal cells. For example, cancer cells typically have a large number of mutated and overexpressed oncoproteins that are dependent on Hsp90 for folding. In addition, because the environment of a tumor is typically hostile due to hypoxia, nutrient deprivation, acidosis, etc., tumor cells may be especially dependent on Hsp90 for survival. Moreover, inhibition of Hsp90 causes simultaneous inhibition of a number of oncoproteins, as well as hormone receptors and transcription factors making it an attractive target for an anti-cancer agent. In fact, benzoquinone ansamycins, a


family of natural products that inhibit Hsp90, has shown evidence of therapeutic activity in clinical trials.

Although promising, benzoquinone ansamycins, and their derivatives, suffer from a number of limitations. For example, they have low oral bioavailability, and their limited solubility makes them difficult to formula. In addition, they are metabolized by polymorphic cytochrome P450 CYP3A4 and are a substrate for P-glycoprotein export pump involved in the development of multidrug resistance. Therefore, a need exist for new therapeutics that improve the prognosis of cancer patients and that reduces or overcomes the limitations of currently used anti-cancer agents.

SUMMARY OF THE INVENTION

The present invention provides novel compounds which inhibit the activity of Hsp90 and are useful in the treatment of proliferative disorders, such as cancer. The present invention also provides new uses for previously disclosed compounds.

The present invention provides compounds having the formula (I):



(I)

and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof. In formula (I), ring A is an aryl or a heteroaryl, wherein the aryl or the heteroaryl are optionally further substituted with one or more substituents in addition to R₃;

R₁ is -OH, -SH, -NR₇H, -OR₂₆, -SR₂₆, -NHR₂₆, -O(CH₂)_mOH, -O(CH₂)_mSH, -O(CH₂)_mNR₇H, -S(CH₂)_mOH, -S(CH₂)_mSH, -S(CH₂)_mNR₇H, -OC(O)NR₁₀R₁₁, -SC(O)NR₁₀R₁₁, -NR₇C(O)NR₁₀R₁₁, -OC(O)R₇, -SC(O)R₇, -NR₇C(O)R₇, -OC(O)OR₇, -SC(O)OR₇, -NR₇C(O)OR₇, -OCH₂C(O)R₇, -SCH₂C(O)R₇, -NR₇CH₂C(O)R₇, -OCH₂C(O)OR₇, -SCH₂C(O)OR₇,

- NR₇CH₂C(O)OR₇, -OCH₂C(O)NR₁₀R₁₁, -SCH₂C(O)NR₁₀R₁₁,
 -NR₇CH₂C(O)NR₁₀R₁₁, -OS(O)_pR₇, -SS(O)_pR₇, -S(O)_pOR₇, -NR₇S(O)_pR₇,
 -OS(O)_pNR₁₀R₁₁, -SS(O)_pNR₁₀R₁₁, -NR₇S(O)_pNR₁₀R₁₁, -OS(O)_pOR₇, -SS(O)_pOR₇,
 -NR₇S(O)_pOR₇, -OC(S)R₇, -SC(S)R₇, -NR₇C(S)R₇, -OC(S)OR₇, -SC(S)OR₇,
 5 -NR₇C(S)OR₇, -OC(S)NR₁₀R₁₁, -SC(S)NR₁₀R₁₁, -NR₇C(S)NR₁₀R₁₁, -OC(NR₈)R₇,
 -SC(NR₈)R₇, -NR₇C(NR₈)R₇, -OC(NR₈)OR₇, -SC(NR₈)OR₇, -NR₇C(NR₈)OR₇,
 -OC(NR₈)NR₁₀R₁₁, -SC(NR₈)NR₁₀R₁₁, -NR₇C(NR₈)NR₁₀R₁₁, -OP(O)(OR₇)₂, or
 -SP(O)(OR₇)₂;
 R₃ is -OH, -SH, -NR₇H, -OR₂₆, -SR₂₆, -NHR₂₆, -O(CH₂)_mOH,
 10 -O(CH₂)_mSH, -O(CH₂)_mNR₇H, -S(CH₂)_mOH, -S(CH₂)_mSH, -S(CH₂)_mNR₇H,
 -OC(O)NR₁₀R₁₁, -SC(O)NR₁₀R₁₁, -NR₇C(O)NR₁₀R₁₁, -OC(O)R₇, -SC(O)R₇,
 -NR₇C(O)R₇, -OC(O)OR₇, -SC(O)OR₇, -NR₇C(O)OR₇, -OCH₂C(O)R₇,
 -SCH₂C(O)R₇, -NR₇CH₂C(O)R₇, -OCH₂C(O)OR₇, -SCH₂C(O)OR₇,
 -NR₇CH₂C(O)OR₇, -OCH₂C(O)NR₁₀R₁₁, -SCH₂C(O)NR₁₀R₁₁,
 15 -NR₇CH₂C(O)NR₁₀R₁₁, -OS(O)_pR₇, -SS(O)_pR₇, -S(O)_pOR₇, -NR₇S(O)_pR₇,
 -OS(O)_pNR₁₀R₁₁, -SS(O)_pNR₁₀R₁₁, -NR₇S(O)_pNR₁₀R₁₁, -OS(O)_pOR₇, -SS(O)_pOR₇,
 -NR₇S(O)_pOR₇, -OC(S)R₇, -SC(S)R₇, -NR₇C(S)R₇, -OC(S)OR₇, -SC(S)OR₇,
 -NR₇C(S)OR₇, -OC(S)NR₁₀R₁₁, -SC(S)NR₁₀R₁₁, -NR₇C(S)NR₁₀R₁₁, -OC(NR₈)R₇,
 -SC(NR₈)R₇, -NR₇C(NR₈)R₇, -OC(NR₈)OR₇, -SC(NR₈)OR₇, -NR₇C(NR₈)OR₇,
 20 -OC(NR₈)NR₁₀R₁₁, -SC(NR₈)NR₁₀R₁₁, -NR₇C(NR₈)NR₁₀R₁₁, -OP(O)(OR₇)₂, or
 -SP(O)(OR₇)₂;

R₅ is an optionally substituted heteroaryl or an optionally substituted 8 to 14 membered aryl;

- R₇ and R₈, for each occurrence, are, independently, -H, an optionally
 25 substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl,
 an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an
 optionally substituted heterocyclyl, an optionally substituted aryl, an optionally
 substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted
 heteraralkyl;

- 30 R₁₀ and R₁₁, for each occurrence, are independently -H, an optionally
 substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl,
 an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an

optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R₁₀ and R₁₁, taken together with the nitrogen to which they are attached, form an optionally substituted heterocyclyl or an optionally substituted

5 heteroaryl;


R₂₆ is a lower alkyl;

p, for each occurrence, is, independently, 1 or 2; and

m, for each occurrence, is independently, 1, 2, 3, or 4.

In one embodiment, ring A of the the compounds of formula (I) is not a
10 substituted [1,2,3]triazole, and/or compounds represented by formula (I) do not include 3-(2,4-dihydroxy-phenyl)-4-(7-naphthalen-1-yl)-5-mercapto-triazole.

The present invention also provides compounds having the formula (II):



(II)

and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof. In formula (II), ring A, R₁, and R₃ are defined as for formula (I); and

R₂ is a substituted phenyl, wherein the phenyl group is substituted with:


- i) one substituent selected from nitro, cyano, a haloalkoxy, an
20 optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl,
25 hydroxylalkyl, alkoxyalkyl, guanadino, -NR₁₀R₁₁, -O-R₂₀, -C(O)R₇, -C(O)OR₂₀, -OC(O)R₇, -C(O)NR₁₀R₁₁, -NR₈C(O)R₇, -SR₇, -S(O)_pR₇, -OS(O)_pR₇, -S(O)_pOR₇, -NR₈S(O)_pR₇, or -S(O)_pNR₁₀R₁₁,
or

- ii) two to five substituents selected from the group consisting of an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, hydroxyalkyl, alkoxyalkyl, -F, -Br, -I, cyano, nitro, guanadino, a haloalkyl, a heteroalkyl, $-NR_{10}R_{11}$, $-OR_7$, $-C(O)R_7$, $-C(O)OR_7$, $-OC(O)R_7$, $-C(O)NR_{10}R_{11}$, $-NR_8C(O)R_7$, $-SR_7$, $-S(O)_pR_7$, $-OS(O)_pR_7$, $-S(O)_pOR_7$, $-NR_8S(O)_pR_7$, or $-S(O)_pNR_{10}R_{11}$; and

R_{20} , for each occurrence, is independently an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl.

In one embodiment, compounds represented by formula (II) do not include 3-(2,4-dihydroxy-phenyl)-4-(7-naphthalen-1-yl)-5-mercapto-triazole, 3-(2,4-dihydroxyphenyl)-4-(2,5-dimethoxyphenyl)-5-mercapto-triazole, 3-(1-phenyl-5-amino-pyrazol-4-yl)-4-(2,4-dichlorophenyl)-5-mercapto-triazole, or 3-(2-hydroxyphenyl)-4-(2,4-dimethylphenyl)-5-mercapto-triazole.

The present invention also provides compounds having the formula (III):




(III)

and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof. In formula (III), ring A, R_1 , and R_3 are defined as for formula (I); and


R_{18} is an optionally substituted cycloalkyl, and optionally substituted cycloalkenyl, or a substituted alkyl, wherein the alkyl group is substituted with one or more substituents independently selected from the group consisting of an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, halo, cyano, nitro, guanadino, a haloalkyl, $-NR_{10}R_{11}$, $-OR_7$, $-C(O)R_7$, $-C(O)OR_7$, $-OC(O)R_7$, $-C(O)NR_{10}R_{11}$, $-NR_8C(O)R_7$, $-SR_7$, $-S(O)_pR_7$, $-OS(O)_pR_7$, $-S(O)_pOR_7$, $-NR_8S(O)_pR_7$, or $-S(O)_pNR_{10}R_{11}$, $-S(O)_pOR_7$, $-OP(O)(OR_7)_2$, or $-SP(O)(OR_7)_2$;

In one embodiment, compounds represented by formula (III) do not include compounds in which R_{18} is not cyclohexyl.

The invention also provides compounds represented by formula (IV) or formula (V):



(IV)



(V)

and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs

thereof. In formulas (IV) and (V), R_1 and R_3 are defined as for formula (I); and


X_{14} is O, S, or NR_7 ;

R_{21} is an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

R_{22} , for each occurrence, is independently -H or is selected from the group consisting of an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl, a haloalkyl, -C(O) R_7 , -C(O)OR $_7$, -OC(O) R_7 , -C(O)NR $_{10}$ R $_{11}$, -NR $_8$ C(O) R_7 , -S(O) $_p$ R $_7$, -S(O) $_p$ OR $_7$, or -S(O) $_p$ NR $_{10}$ R $_{11}$; and

R_{23} and R_{24} , for each occurrence, are independently -H or are selected from the group consisting of an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl, halo, cyano, nitro, guanadino, a haloalkyl, a heteroalkyl, -NR $_{10}$ R $_{11}$, -OR $_7$, -C(O) R_7 , -C(O)OR $_7$, -OC(O) R_7 , -C(O)NR $_{10}$ R $_{11}$, -NR $_8$ C(O) R_7 , -SR $_7$, -S(O) $_p$ R $_7$, -OS(O) $_p$ R $_7$, -S(O) $_p$ OR $_7$, -NR $_8$ S(O) $_p$ R $_7$, or -S(O) $_p$ NR $_{10}$ R $_{11}$.

The present invention also provides compounds represented by formula (VI):



and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof, wherein:

X_{41} is O, S, or NR $_{42}$;
 X_{42} is CR $_{44}$ or N;

Y_{40} is N or CR_{43} ;

Y_{41} is N or CR_{45} ;

Y_{42} , for each occurrence, is independently N, C or CR_{46} ;

Z is OH, SH, or NHR_7 ;

- 5 R_{41} is -H, -OH, -SH, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, halo, cyano, nitro,
- 10 guanadino, a haloalkyl, a heteroalkyl, an alkoxy or cycloalkoxy, a haloalkoxy, $-NR_{10}R_{11}$, $-OR_7$, $-C(O)R_7$, $-C(O)OR_7$, $-C(S)R_7$, $-C(O)SR_7$, $-C(S)SR_7$, $-C(S)OR_7$, $-C(S)NR_{10}R_{11}$, $-C(NR_8)OR_7$, $-C(NR_8)R_7$, $-C(NR_8)NR_{10}R_{11}$, $-C(NR_8)SR_7$, $-OC(O)R_7$, $-OC(O)OR_7$, $-OC(S)OR_7$, $-OC(NR_8)OR_7$, $-SC(O)R_7$, $-SC(O)OR_7$, $-SC(NR_8)OR_7$, $-OC(S)R_7$, $-SC(S)R_7$, $-SC(S)OR_7$, $-OC(O)NR_{10}R_{11}$, $-OC(S)NR_{10}R_{11}$,
- 15 $-OC(NR_8)NR_{10}R_{11}$, $-SC(O)NR_{10}R_{11}$, $-SC(NR_8)NR_{10}R_{11}$, $-SC(S)NR_{10}R_{11}$, $-OC(NR_8)R_7$, $-SC(NR_8)R_7$, $-C(O)NR_{10}R_{11}$, $-NR_8C(O)R_7$, $-NR_7C(S)R_7$, $-NR_7C(S)OR_7$, $-NR_7C(NR_8)R_7$, $-NR_7C(O)OR_7$, $-NR_7C(NR_8)OR_7$, $-NR_7C(O)NR_{10}R_{11}$, $-NR_7C(S)NR_{10}R_{11}$, $-NR_7C(NR_8)NR_{10}R_{11}$, $-SR_7$, $-S(O)_pR_7$, $-OS(O)_pR_7$, $-OS(O)_pOR_7$, $-OS(O)_pNR_{10}R_{11}$, $-S(O)_pOR_7$, $-NR_8S(O)_pR_7$,
- 20 $-NR_7S(O)_pNR_{10}R_{11}$, $-NR_7S(O)_pOR_7$, $-S(O)_pNR_{10}R_{11}$, $-SS(O)_pR_7$, $-SS(O)_pOR_7$, $-SS(O)_pNR_{10}R_{11}$, $-OP(O)(OR_7)_2$, or $-SP(O)(OR_7)_2$;

- R_{42} is -H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, hydroxyalkyl, alkoxyalkyl, a haloalkyl, a heteroalkyl, $-C(O)R_7$, $-(CH_2)_mC(O)OR_7$, $-C(O)OR_7$, $-OC(O)R_7$, $-C(O)NR_{10}R_{11}$, $-S(O)_pR_7$, $-S(O)_pOR_7$, or $-S(O)_pNR_{10}R_{11}$;
- 25

- R_{43} and R_{44} are, independently, -H, -OH, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted
- 30

heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, hydroxyalkyl, alkoxyalkyl, halo, cyano, nitro, guanadino, a haloalkyl, a heteroalkyl, -C(O)R₇, -C(O)OR₇, -OC(O)R₇, -C(O)NR₁₀R₁₁, -NR₈C(O)R₇, -SR₇, -S(O)_pR₇, -OS(O)_pR₇, -S(O)_pOR₇, -NR₈S(O)_pR₇, -S(O)_pNR₁₀R₁₁, or R₄₃ and R₄₄ taken


- 5 together with the carbon atoms to which they are attached form an optionally substituted cycloalkenyl, an optionally substituted aryl, an optionally substituted heterocyclyl, or an optionally substituted heteroaryl;

R₄₅ is -H, -OH, -SH, -NR₇H, -OR₂₆, -SR₂₆, -NHR₂₆, -O(CH₂)_mOH, -O(CH₂)_mSH, -O(CH₂)_mNR₇H, -S(CH₂)_mOH, -S(CH₂)_mSH, -S(CH₂)_mNR₇H, -OC(O)NR₁₀R₁₁, -SC(O)NR₁₀R₁₁, -NR₇C(O)NR₁₀R₁₁, -OC(O)R₇, -SC(O)R₇, -NR₇C(O)R₇, -OC(O)OR₇, -SC(O)OR₇, -NR₇C(O)OR₇, -OCH₂C(O)R₇, -SCH₂C(O)R₇, -NR₇CH₂C(O)R₇, -OCH₂C(O)OR₇, -SCH₂C(O)OR₇, -NR₇CH₂C(O)OR₇, -OCH₂C(O)NR₁₀R₁₁, -SCH₂C(O)NR₁₀R₁₁, -NR₇CH₂C(O)NR₁₀R₁₁, -OS(O)_pR₇, -SS(O)_pR₇, -NR₇S(O)_pR₇, -OS(O)_pNR₁₀R₁₁, -SS(O)_pNR₁₀R₁₁, -NR₇S(O)_pNR₁₀R₁₁, -OS(O)_pOR₇, -SS(O)_pOR₇, -NR₇S(O)_pOR₇, -OC(S)R₇, -SC(S)R₇, -NR₇C(S)R₇, -OC(S)OR₇, -SC(S)OR₇, -NR₇C(S)OR₇, -OC(S)NR₁₀R₁₁, -SC(S)NR₁₀R₁₁, -NR₇C(S)NR₁₀R₁₁, -OC(NR₈)R₇, -SC(NR₈)R₇, -NR₇C(NR₈)R₇, -OC(NR₈)OR₇, -SC(NR₈)OR₇, -NR₇C(NR₈)OR₇, -OC(NR₈)NR₁₀R₁₁, -SC(NR₈)NR₁₀R₁₁, or -NR₇C(NR₈)NR₁₀R₁₁;

- 20 R₄₆, for each occurrence, is independently selected from the group consisting of H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, halo, cyano, nitro, guanadino, a haloalkyl, a heteroalkyl, -NR₁₀R₁₁, -OR₇, -C(O)R₇, -C(O)OR₇, -OC(O)R₇, -C(O)NR₁₀R₁₁, -NR₈C(O)R₇, -SR₇, -S(O)_pR₇, -OS(O)_pR₇, -S(O)_pOR₇, -NR₈S(O)_pR₇, or -S(O)_pNR₁₀R₁₁;

R₇, R₈, R₁₀, R₁₁, R₂₆, p, and m are defined as above.

- 30 The present invention also provides compounds represented by formula (VII):




(VII)

and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs, wherein:

5 Z₁ is -OH or -SH; and

X₄₂, R₄₁, R₄₂, R₄₃, and R₄₅ are defined as above.

The present invention also provides compounds having the formula (VIII):



(VIII)

and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof, wherein:

X₄₅ is CR₅₄ or N;

Z₁ is -OH or -SH;


15 R₅₂ is selected from the group consisting of -H, methyl, ethyl, n-propyl, isopropyl, n-butyl, n-pentyl, n-hexyl, -(CH₂)₂OCH₃, -CH₂C(O)OH, and -C(O)N(CH₃)₂;

R_{53} and R_{54} are each, independently, -H, methyl, ethyl, or isopropyl; or R_{53} and R_{54} taken together with the carbon atoms to which they are attached form a phenyl, cyclohexenyl, or cyclooctenyl ring;

R_{55} is selected from the group consisting of -H, -OH, -OCH₃, and -OCH₂CH₃; and

R_{56} is selected from the group consisting of -H, methyl, ethyl, isopropyl, and cyclopropyl.

The present invention also provides compounds having the formula (IX):



and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof, wherein,

X_{44} , for each occurrence, is independently, O, S, NR₄₂ or C(R₄₆)₂;

Y_{43} is NR₄₂, C(R₄₆)₂, C(R₄₆)₂-C(R₄₆)₂, C(O), C(S), C(R₄₆)₂C(O), or C(R₄₆)₂C(S);

Y_{41} , Y_{42} , Z, R_{41} , R_{42} , and R_{46} are defined as above.

In one embodiment, in formula (IX), R_{41} is selected from the group consisting of -H, lower alkyl, lower alkoxy, lower cycloalkyl, and lower cycloalkoxy.

In another embodiment, in formula (IX), R_{41} is selected from the group consisting of -H, methyl, ethyl, propyl, isopropyl, cyclopropyl, methoxy, ethoxy, propoxy, and cyclopropoxy.

In another embodiment, in formula (IX), R_{42} is selected from the group consisting of -H, methyl, ethyl, n-propyl, isopropyl, cyclopropyl, n-butyl, *sec*-butyl,

tert-butyl, n-pentyl, n-hexyl, -C(O)OH, -(CH₂)_mC(O)OH, -CH₂OCH₃, -CH₂CH₂OCH₃, and -C(O)N(CH₃)₂.

In another embodiment, in formula (IX), Y₄₁ is CR₄₅. Preferably, R₄₅ is H, a lower alkoxy, or -OH.

5 In another embodiment, in formula (IX), Y₄₂ is CH.

In another embodiment, in formula (IX), Y₄₃ is CH₂.

In another embodiment, in formula (IX), Y₄₃ is NR₄₂, wherein R₄₂ is H or a lower alkyl.

10 In another embodiment, in formula (IX), one of X₄₄ is NR₄₂ and the other is CH₂ or C(R₆)₂. Preferably, one of X₄₄ is NR₄₂ and the other is CH₂.

In another embodiment, in formula (VI), Z is -OH.

In another embodiment, Z is -SH.

In another embodiment, the compound is selected from the group consisting of:

3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1,3-benzodioxol-5-yl)-5-mercapto-[1,2,4] triazole;

3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(indan-5-yl)-5-mercapto-[1,2,4] triazole;

4-Ethyl-6-[5-mercapto-4-(1-methyl-2,3-dihydro-1H-indol-5-yl)-4H-[1,2,4]triazol-3-yl]-benzene-1,3-diol;

5-(3-(5-ethyl-2,4-dihydroxyphenyl)-5-mercapto-4H-1,2,4-triazol-4-yl)indolin-2-one;

5-(3-(5-ethyl-2,4-dihydroxyphenyl)-5-mercapto-4H-1,2,4-triazol-4-yl)-1H-benzo[d]imidazol-2(3H)-one;

5-(3-(5-ethyl-2,4-dihydroxyphenyl)-5-mercapto-4H-1,2,4-triazol-4-yl)-1-methylindolin-2-one;

4-isopropyl-6-(5-mercapto-4-(4-propyl-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl)-4H-1,2,4-triazol-3-yl)benzene-1,3-diol;

6-(3-(5-ethyl-2,4-dihydroxyphenyl)-5-mercapto-4H-1,2,4-triazol-4-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one;


6-(3-(5-ethyl-2,4-dihydroxyphenyl)-5-mercapto-4H-1,2,4-triazol-4-yl)-3-methylbenzo[d]thiazol-2(3H)-one;

6-(3-(5-ethyl-2,4-dihydroxyphenyl)-5-mercapto-4H-1,2,4-triazol-4-

yl)benzo[d]thiazol-2(3H)-one; and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof.

Compounds of formula (IX) inhibit the activity of Hsp90 and are particularly useful for treating or preventing proliferative disorders, such as cancer. In addition, compounds of formula (IX) are particularly useful in treating cancer when given in combination with other anti-cancer agent.

5 The present invention also provides compounds having the formula (X):



and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof, wherein:

10 X_{41} , Y_{41} , Y_{42} , Z , R_7 , R_8 , R_{10} , R_{11} , R_{41} , R_{46} and p are defined as above.

The compounds shown in Table 1 or compounds of any formula herein, or tautomers, pharmaceutically acceptable salts, solvates, clathrates, hydrates, polymorphs or prodrugs thereof, inhibit the activity of Hsp90 and, thereby facilitates
 15 the degradation of Hsp90 client proteins. Hsp90 is necessary for the survival of normal eukaryotic cells. However, Hsp90 is over expressed in many tumor types indicating that it may play a significant role in the survival of cancer cells and that cancer cells may be more sensitive to inhibition of Hsp90 than normal cells. Thus, the compounds shown in Table 1 or compounds of any formula herein, or tautomers,
 20 pharmaceutically acceptable salts, solvates, clathrates, hydrates, polymorphs or prodrugs thereof, are useful treating proliferative disorders such as cancer.

Although chemotherapeutic agents initially cause tumor regression, most agents that are currently used to treat cancer target only one pathway to tumor

progression. Therefore, in many instances, after treatment with one or more chemotherapeutic agents, a tumor develops multidrug resistance and no longer responds positively to treatment. One of the advantages of inhibiting Hsp90 activity is that several of its client proteins, which are mostly protein kinases or transcription factors involved in signal transduction, have been shown to be involved in the progression of cancer. Thus, inhibition of Hsp90 provides a method of short circuiting several pathways for tumor progression simultaneously. Therefore, treatment of tumors with an Hsp90 inhibitor of the invention either alone, or in combination with other chemotherapeutic agents, is more likely to result in regression or elimination of the tumor, and less likely to result in the development of more aggressive multidrug resistant tumors than other currently available therapies.

BRIEF DESCRIPTION OF THE DRAWINGS

The foregoing and other objects, features and advantages of the invention will be apparent from the following more particular description of preferred embodiments of the invention, as illustrated in the accompanying drawings in which like reference characters refer to the same parts throughout the different views. The drawings are not necessarily to scale, emphasis instead being placed upon illustrating the principles of the invention.

Figure 1 is a graph showing the ATPase activity of Hsp90 when untreated, when treated with 40 mM of Geldanamycin, a known Hsp90 inhibitor as a positive control, and when treated with 40 μ M or 4 μ M of Compound 108 of the invention;

Figure 2 is gel showing the amount of Her2, an Hsp90 client protein, in cells that are untreated, in cells that have been treated with 0.5 μ M, 2 μ M, or 5 μ M of 17AAG, a known Hsp90 inhibitor, and in cells that have been treated with 0.5 μ M, 2 μ M, or 5 μ M of Compound 108 or Compound 49;

Figure 3 is a graph showing an FACSCalibur flow cytometer analysis of the c-kit positive population of HEL92.1.7 cells treated with Hsp90 inhibitors of the invention or 17AAG (as a positive control). The results indicate that the Hsp90 inhibitors of the invention induce c-kit degradation at a lower concentration than 17AAG, an Hsp90 inhibitor that is currently in phase II clinical trials.

Figure 4 is a graph showing an FACSCalibur flow cytometer analysis of the c-kit positive population of Kasumi-1 cells treated with Hsp90 inhibitors of the invention or 17AAG (as a positive control). The results indicate that the Hsp90 inhibitors of the invention induce c-kit degradation at a lower concentration than 17AAG, an Hsp90 inhibitor that is currently in phase II clinical trials.

Figure 5 is a Western blot analysis of the c-kit from Kasumi-1 cells treated with Hsp90 inhibitors of the invention or 17AAG (as a positive control).

Figure 6 is a Western blot analysis of the c-met from NCI-H1193 cells treated with Hsp90 inhibitors of the invention or 17AAG (as a positive control).

Figure 7 displays the results of a *nude* mouse xenograft study to determine the effect of Compound 49 on the *in vivo* growth rate of the human tumor cell line MDA-MB-435S. Tumor bearing animals (8 mice/group) were intraperitoneal (IP) injected 5 times per week for 3 weeks (hatched bar) and the average tumor volumes for each group (+/- SEM) were determined every 3-5 days. Treatment with a dose of 300 mg/kg body weight of Compound 49 decreased the growth rate of MDA-MB-435S cells in *nude* mice to a greater extent than did a dose of 100 mg/kg body weight of the Hsp90 inhibitor 17-allylamino-17-demethoxygeldanamycin (17-AAG);

Figure 8 demonstrates that treatment with Compound 49 did not cause toxicity in a *nude* mouse xenograft model using the human tumor cell line MDA-MB-435S (tumor growth data from the same study is presented in Figure 3). Tumor bearing animals (8 mice/group) were intraperitoneal (IP) injected 5 times per week for 3 weeks (hatched bar) and the average percent changes in body weights for each group relative to the start of dosing were determined every 1-3 days (error bars not shown for clarity; mean coefficient of variation = 18%). Treatment with a dose of 300 mg/kg body weight of Compound 49 was not toxic, as indicated by its lack of an effect on the body weights in animals treated with Compound 49 versus vehicle treated animals;

Figure 9 displays the results of a *nude* mouse xenograft study to determine the effect of Compound 188 on the *in vivo* growth rate of RERF-LC-AI^{IVP} human lung tumor cells. Tumor bearing animals (8 mice/group) were i.p. injected 5 times per week for a total of 15 doses (hatched bar) and the average tumor volumes for

each group (error bars represent SEM) were determined every 3-4 days. Treatment with a dose of 200 mg/kg body weight of Compound 188 inhibited tumor growth, as did a dose of 75 mg/kg body weight of 17-AAG (both compounds were dosed at approximately their maximum tolerated doses in *nude* mice); and

- 5 Figure 10 demonstrates that treatment with Compound 188 does not cause overt toxicity in a *nude* mouse xenograft model using RERF-LC-AI^{IVP} human lung tumor cells (data derived from the same study presented in Figure 5). Tumor bearing animals (8 mice/group) were i.p. injected 5 times per week for a total of 15 doses (hatched bar) and the cumulative average percent changes in body weights for
10 each group relative to the start of dosing were determined every 2-3 days. Treatment with a dose of 200 mg/kg body weight of Compound 188 was not overtly toxic, as indicated by the minimal effects on the animal body weights in the test article-treated versus vehicle-treated groups.

DETAILED DESCRIPTION OF THE INVENTION

- 15 A description of preferred embodiments of the invention follows.

 The present invention provides compounds and uses of said compounds. The present invention encompasses the use of the compounds of the invention to inhibit Hsp90 activity and for the treatment of a proliferative disorder, such as cancer. In particular, the present invention encompasses the use of compounds of
20 the invention to slow or stop the growth of cancerous cells or to reduce or eliminate cancerous cells in a mammal.

 In certain embodiments, the compounds of the invention can be used in combination with other chemotherapeutic agents and may help to prevent or reduce the development of multidrug resistant cancerous cells in a mammal. In this
25 embodiment, the compounds of the invention may allow a reduced efficacious amount of a second chemotherapeutic agent given to a mammal, because compounds of the invention should inhibit the development of multidrug resistant cancerous cells.

30 A. Terminology

 Unless otherwise specified, the below terms used herein are defined as follows:

As used herein, the term "alkyl" means a saturated straight chain or branched non-cyclic hydrocarbon having from 1 to 10 carbon atoms. Representative saturated straight chain alkyls include methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl and n-decyl; while saturated branched alkyls include

5 isopropyl, *sec*-butyl, isobutyl, *tert*-butyl, isopentyl, 2-methylbutyl, 3-methylbutyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 2-methylhexyl, 3-methylhexyl, 4-methylhexyl, 5-methylhexyl, 2,3-dimethylbutyl, 2,3-dimethylpentyl, 2,4-dimethylpentyl, 2,3-dimethylhexyl, 2,4-dimethylhexyl, 2,5-dimethylhexyl, 2,2-dimethylpentyl, 2,2-dimethylhexyl, 3,3-dimethylpentyl, 3,3-dimethylhexyl, 4,4-

10 dimethylhexyl, 2-ethylpentyl, 3-ethylpentyl, 2-ethylhexyl, 3-ethylhexyl, 4-ethylhexyl, 2-methyl-2-ethylpentyl, 2-methyl-3-ethylpentyl, 2-methyl-4-ethylpentyl, 2-methyl-2-ethylhexyl, 2-methyl-3-ethylhexyl, 2-methyl-4-ethylhexyl, 2,2-diethylpentyl, 3,3-diethylhexyl, 2,2-diethylhexyl, 3,3-diethylhexyl and the like. The term "(C₁-C₆)alkyl" means a saturated straight chain or branched non-cyclic

15 hydrocarbon having from 1 to 6 carbon atoms. Representative (C₁-C₆)alkyl groups are those shown above having from 1 to 6 carbon atoms. Alkyl groups included in compounds of this invention may be optionally substituted with one or more substituents.

As used herein, the term "alkenyl" means a saturated straight chain or

20 branched non-cyclic hydrocarbon having from 2 to 10 carbon atoms and having at least one carbon-carbon double bond. Representative straight chain and branched (C₂-C₁₀)alkenyls include vinyl, allyl, 1-butenyl, 2-butenyl, isobutylenyl, 1-pentenyl, 2-pentenyl, 3-methyl-1-butenyl, 2-methyl-2-butenyl, 2,3-dimethyl-2-butenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 1-heptenyl, 2-heptenyl, 3-heptenyl, 1-octenyl, 2-

25 octenyl, 3-octenyl, 1-nonenyl, 2-nonenyl, 3-nonenyl, 1-decenyl, 2-decenyl, 3-decenyl and the like. Alkenyl groups may be optionally substituted with one or more substituents.

As used herein, the term "alkynyl" means a saturated straight chain or branched non-cyclic hydrocarbon having from 2 to 10 carbon atoms and having at

30 least one carbon-carbon triple bond. Representative straight chain and branched alkynyls include acetylenyl, propynyl, 1-butylnyl, 2-butylnyl, 1-pentylnyl, 2-pentylnyl, 3-methyl-1-butylnyl, 4-pentylnyl, 1-hexynyl, 2-hexynyl, 5-hexynyl, 1-heptynyl, 2-

heptynyl, 6-heptynyl, 1-octynyl, 2-octynyl, 7-octynyl, 1-nonyl, 2-nonyl, 8-nonyl, 1-decynyl, 2-decynyl, 9-decynyl, and the like. Alkynyl groups may be optionally substituted with one or more substituents.

As used herein, the term "cycloalkyl" means a saturated, mono- or polycyclic alkyl radical having from 3 to 20 carbon atoms. Representative cycloalkyls include cyclopropyl, 1-methylcyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, -cyclodecyl, octahydro-pentalenyl, and the like. Cycloalkyl groups may be optionally substituted with one or more substituents.

As used herein, the term "cycloalkenyl" means a mono- or poly- cyclic non-aromatic alkyl radical having at least one carbon-carbon double bond in the cyclic system and from 3 to 20 carbon atoms. Representative cycloalkenyls include cyclopentenyl, cyclopentadienyl, cyclohexenyl, cyclohexadienyl, cycloheptenyl, cycloheptadienyl, cycloheptatrienyl, cyclooctenyl, cyclooctadienyl, cyclooctatrienyl, cyclooctatetraenyl, cyclononenyl, cyclononadienyl, cyclodecenyl, cyclodecadienyl, 1,2,3,4,5,8-hexahydronaphthalenyl and the like. Cycloalkenyl groups may be optionally substituted with one or more substituents.

As used herein, the term "haloalkyl" means an alkyl group in which one or more (including all) the hydrogen radicals are replaced by a halo group, wherein each halo group is independently selected from -F, -Cl, -Br, and -I. The term "halomethyl" means a methyl in which one to three hydrogen radical(s) have been replaced by a halo group. Representative haloalkyl groups include trifluoromethyl, bromomethyl, 1,2-dichloroethyl, 4-iodobutyl, 2-fluoropentyl, and the like.

As used herein, an "alkoxy" is an alkyl group which is attached to another moiety via an oxygen linker.

As used herein, an "haloalkoxy" is an haloalkyl group which is attached to another moiety via an oxygen linker.

As used herein, the term an "aromatic ring" or "aryl" means a hydrocarbon monocyclic or polycyclic radical in which at least one ring is aromatic. Examples of suitable aryl groups include, but are not limited to, phenyl, tolyl, anthracenyl, fluorenyl, indenyl, azulenyl, and naphthyl, as well as benzo-fused carbocyclic moieties such as 5,6,7,8-tetrahydronaphthyl. Aryl groups may be optionally substituted with one or more substituents. In one embodiment, the aryl group is a

monocyclic ring, wherein the ring comprises 6 carbon atoms, referred to herein as “(C₆)aryl.”

As used herein, the term “aralkyl” means an aryl group that is attached to another group by a (C₁-C₆)alkylene group. Representative aralkyl groups include
5 benzyl, 2-phenyl-ethyl, naphth-3-yl-methyl and the like. Aralkyl groups may be optionally substituted with one or more substituents.

As used herein, the term “alkylene” refers to an alkyl group that has two points of attachment. The term “(C₁-C₆)alkylene” refers to an alkylene group that has from one to six carbon atoms. Straight chain (C₁-C₆)alkylene groups are
10 preferred. Non-limiting examples of alkylene groups include methylene (-CH₂-), ethylene (-CH₂CH₂-), n-propylene (-CH₂CH₂CH₂-), isopropylene (-CH₂CH(CH₃)-), and the like. Alkylene groups may be optionally substituted with one or more substituents.

As used herein, the term “heterocyclyl” means a monocyclic (typically
15 having 3- to 10-members) or a polycyclic (typically having 7- to 20-members) heterocyclic ring system which is either a saturated ring or a unsaturated non-aromatic ring. A 3- to 10-membered heterocycle can contain up to 5 heteroatoms; and a 7- to 20-membered heterocycle can contain up to 7 heteroatoms. Typically, a heterocycle has at least one carbon atom ring member. Each heteroatom is
20 independently selected from nitrogen, which can be oxidized (e.g., N(O)) or quaternized; oxygen; and sulfur, including sulfoxide and sulfone. The heterocycle may be attached via any heteroatom or carbon atom. Representative heterocycles include morpholinyl, thiomorpholinyl, pyrrolidinonyl, pyrrolidinyl, piperidinyl, piperazinyl, hydantoinyl, valerolactamyl, oxiranyl, oxetanyl, tetrahydrofuranyl,
25 tetrahydropyranyl, tetrahydropyrindinyl, tetrahydropyrimidinyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, and the like. A heteroatom may be substituted with a protecting group known to those of ordinary skill in the art, for example, the hydrogen on a nitrogen may be substituted with a tert-butoxycarbonyl group. Furthermore, the heterocyclyl may be optionally substituted with one or
30 more substituents. Only stable isomers of such substituted heterocyclic groups are contemplated in this definition.

As used herein, the term "heteroaromatic", "heteroaryl" or like terms means a monocyclic or polycyclic heteroaromatic ring comprising carbon atom ring members and one or more heteroatom ring members. Each heteroatom is independently selected from nitrogen, which can be oxidized (e.g., N(O)) or quaternized; oxygen; and sulfur, including sulfoxide and sulfone. Representative heteroaryl groups include pyridyl, 1-oxo-pyridyl, furanyl, benzo[1,3]dioxolyl, benzo[1,4]dioxinyl, thienyl, pyrrolyl, oxazolyl, imidazolyl, thiazolyl, a isoxazolyl, quinolinyl, pyrazolyl, isothiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, a triazinyl, triazolyl, thiadiazolyl, isoquinolinyl, indazolyl, benzoxazolyl, benzofuryl, indoliziny, imidazopyridyl, tetrazolyl, benzimidazolyl, benzothiazolyl, benzothiadiazo, benzoxadiazolyl, indolyl, tetrahydroindolyl, azaindolyl, imidazopyridyl, quinazolinyl, purinyl, pyrrolo[2,3]pyrimidinyl, pyrazolo[3,4]pyrimidinyl, imidazo[1,2-a]pyridyl, and benzothienyl. In one embodiment, the heteroaromatic ring is selected from 5-8 membered monocyclic heteroaryl rings. The point of attachment of a heteroaromatic or heteroaryl ring to another group may be at either a carbon atom or a heteroatom of the heteroaromatic or heteroaryl rings. Heteroaryl groups may be optionally substituted with one or more substituents.

As used herein, the term "(C₅)heteroaryl" means an aromatic heterocyclic ring of 5 members, wherein at least one carbon atom of the ring is replaced with a heteroatom such as, for example, oxygen, sulfur or nitrogen. Representative (C₅)heteroaryls include furanyl, thienyl, pyrrolyl, oxazolyl, imidazolyl, thiazolyl, isoxazolyl, pyrazolyl, isothiazolyl, pyrazinyl, triazolyl, thiadiazolyl, and the like.

As used herein, the term "(C₆)heteroaryl" means an aromatic heterocyclic ring of 6 members, wherein at least one carbon atom of the ring is replaced with a heteroatom such as, for example, oxygen, nitrogen or sulfur. Representative (C₆)heteroaryls include pyridyl, pyridazinyl, pyrazinyl, triazinyl, tetrazinyl and the like.

As used herein, the term "heteroaralkyl" means a heteroaryl group that is attached to another group by a (C₁-C₆)alkylene. Representative heteroaralkyls include 2-(pyridin-4-yl)-propyl, 2-(thien-3-yl)-ethyl, imidazol-4-yl-methyl and the

like. Heteroaralkyl groups may be optionally substituted with one or more substituents.

As used herein, the term "halogen" or "halo" means -F, -Cl, -Br or -I.

Suitable substituents for an alkyl, alkylene, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocyclyl, aryl, aralkyl, heteroaryl, and heteroaralkyl groups include any substituent which will form a stable compound of the invention. Examples of substituents for an alkyl, alkylene, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocyclyl, aryl, aralkyl, heteroaryl, and heteroarylalkyl include an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, a haloalkyl, -C(O)NR₂₈R₂₉, -C(S)NR₂₈R₂₉, -C(NR₃₂)NR₂₈R₂₉, -NR₃₀C(O)R₃₁, -NR₃₀C(S)R₃₁, -NR₃₀C(NR₃₂)R₃₁, halo, -OR₃₀, cyano, nitro, haloalkoxy, -C(O)R₃₀, -C(S)R₃₀, -C(NR₃₂)R₃₀, -NR₂₈R₂₉, -C(O)OR₃₀, -C(S)OR₃₀, -C(NR₃₂)OR₃₀, -OC(O)R₃₀, -OC(S)R₃₀, -OC(NR₃₂)R₃₀, -NR₃₀C(O)NR₂₈R₂₉, -NR₃₀C(S)NR₂₈R₂₉, -NR₃₀C(NR₃₂)NR₂₈R₂₉, -OC(O)NR₂₈R₂₉, -OC(S)NR₂₈R₂₉, -OC(NR₃₂)NR₂₈R₂₉, -NR₃₀C(O)OR₃₁, -NR₃₀C(S)OR₃₁, -NR₃₀C(NR₃₂)OR₃₁, -S(O)_hR₃₀, -OS(O)_pR₃₀, -NR₃₀S(O)_pR₃₀, -S(O)_pNR₂₈R₂₉, -OS(O)_pNR₂₈R₂₉, or -NR₃₀S(O)_pNR₂₈R₂₉, wherein R₂₈ and R₂₉, for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R₂₈ and R₂₉ taken together with the nitrogen to which they are attached is optionally substituted heterocyclyl or optionally substituted heteroaryl;

R₃₀ and R₃₁ for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally

substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; and

R_{32} , for each occurrence is, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, $-C(O)R_{30}$, $-C(O)NR_{28}R_{29}$, $-S(O)_pR_{30}$, or $-S(O)_pNR_{28}R_{29}$;

p , for each occurrence, is independently, 1 or 2; and
10 h is 0, 1 or 2.

In addition, alkyl, cycloalkyl, alkylene, a heterocyclyl, and any saturated portion of a alkenyl, cycloalkenyl, alkynyl, aralkyl, and heteroaralkyl groups, may also be substituted with $=O$, $=S$, $=N-R_{32}$.

When a heterocyclyl, heteroaryl, or heteroaralkyl group contains a nitrogen atom, it may be substituted or unsubstituted. When a nitrogen atom in the aromatic ring of a heteroaryl group has a substituent the nitrogen may be a quaternary nitrogen.

As used herein, the terms "subject", "patient" and "mammal" are used interchangeably. The terms "subject" and "patient" refer to an animal (e.g., a bird such as a chicken, quail or turkey, or a mammal), preferably a mammal including a non-primate (e.g., a cow, pig, horse, sheep, rabbit, guinea pig, rat, cat, dog, and mouse) and a primate (e.g., a monkey, chimpanzee and a human), and more preferably a human. In one embodiment, the subject is a non-human animal such as a farm animal (e.g., a horse, cow, pig or sheep), or a pet (e.g., a dog, cat, guinea pig or rabbit). In a preferred embodiment, the subject is a human.

As used herein, the term "lower" refers to a group having up to four atoms. For example, a "lower alkyl" refers to an alkyl radical having from 1 to 4 carbon atoms, "lower alkoxy" refers to $-O-(C_1-C_4)alkyl$ and a "lower alkenyl" or "lower alkynyl" refers to an alkenyl or alkynyl radical having from 2 to 4 carbon atoms, respectively.

Unless indicated otherwise, the compounds of the invention containing reactive functional groups (such as (without limitation) carboxy, hydroxy, thiol, and

amino moieties) also include protected derivatives thereof. "Protected derivatives" are those compounds in which a reactive site or sites are blocked with one or more protecting groups. Examples of suitable protecting groups for hydroxyl groups include benzyl, methoxymethyl, allyl, trimethylsilyl, tert-butyldimethylsilyl, acetate, and the like. Examples of suitable amine protecting groups include benzyloxycarbonyl, tert-butoxycarbonyl, tert-butyl, benzyl and fluorenylmethyloxycarbonyl (Fmoc). Examples of suitable thiol protecting groups include benzyl, tert-butyl, acetyl, methoxymethyl and the like. Other suitable protecting groups are well known to those of ordinary skill in the art and include those found in T. W. Greene, *Protecting Groups in Organic Synthesis*, John Wiley & Sons, Inc. 1981.

As used herein, the term "compound(s) of this invention" and similar terms refers to a compound of formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, hydrate, polymorph or prodrug thereof, and also include protected derivatives thereof.

The compounds of the invention may contain one or more chiral centers and/or double bonds and, therefore, exist as stereoisomers, such as double-bond isomers (*i.e.*, geometric isomers), enantiomers, or diastereomers. According to this invention, the chemical structures depicted herein, including the compounds of this invention, encompass all of the corresponding compounds' enantiomers, diastereomers and geometric isomers, that is, both the stereochemically pure form (*e.g.*, geometrically pure, enantiomerically pure, or diastereomerically pure) and isomeric mixtures (*e.g.*, enantiomeric, diastereomeric and geometric isomeric mixtures). In some cases, one enantiomer, diastereomer or geometric isomer will possess superior activity or an improved toxicity or kinetic profile compared to other isomers. In those cases, such enantiomers, diastereomers and geometric isomers of compounds of this invention are preferred.

As used herein, the term "polymorph" means solid crystalline forms of a compound of the present invention or complex thereof. Different polymorphs of the same compound can exhibit different physical, chemical and/or spectroscopic properties. Different physical properties include, but are not limited to stability (*e.g.*, to heat or light), compressibility and density (important in formulation and product manufacturing), and dissolution rates (which can affect bioavailability).

Differences in stability can result from changes in chemical reactivity (*e.g.*, differential oxidation, such that a dosage form discolors more rapidly when comprised of one polymorph than when comprised of another polymorph) or mechanical characteristics (*e.g.*, tablets crumble on storage as a kinetically favored polymorph converts to thermodynamically more stable polymorph) or both (*e.g.*, tablets of one polymorph are more susceptible to breakdown at high humidity). Different physical properties of polymorphs can affect their processing. For example, one polymorph might be more likely to form solvates or might be more difficult to filter or wash free of impurities than another due to, for example, the shape or size distribution of particles of it.

As used herein, the term "hydrate" means a compound of the present invention or a salt thereof, that further includes a stoichiometric or non-stoichiometric amount of water bound by non-covalent intermolecular forces.

As used herein, the term "clathrate" means a compound of the present invention or a salt thereof in the form of a crystal lattice that contains spaces (*e.g.*, channels) that have a guest molecule (*e.g.*, a solvent or water) trapped within.

As used herein and unless otherwise indicated, the term "prodrug" means a derivative of a compound that can hydrolyze, oxidize, or otherwise react under biological conditions (*in vitro* or *in vivo*) to provide a compound of this invention.

Prodrugs may become active upon such reaction under biological conditions, or they may have activity in their unreacted forms. Examples of prodrugs contemplated in this invention include, but are not limited to, analogs or derivatives of compounds of formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X), or Table 1 that comprise biohydrolyzable moieties such as biohydrolyzable amides, biohydrolyzable esters, biohydrolyzable carbamates, biohydrolyzable carbonates, biohydrolyzable ureides, and biohydrolyzable phosphate analogues. Other examples of prodrugs include derivatives of compounds of formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X), or Table 1 that comprise -NO, -NO₂, -ONO, or -ONO₂ moieties. Prodrugs can typically be prepared using well-known methods, such as those described by 1 BURGER'S MEDICINAL CHEMISTRY AND DRUG DISCOVERY (1995) 172-178, 949-982 (Manfred E. Wolff ed., 5th ed).

As used herein and unless otherwise indicated, the terms "biohydrolyzable amide", "biohydrolyzable ester", "biohydrolyzable carbamate", "biohydrolyzable carbonate", "biohydrolyzable ureide" and "biohydrolyzable phosphate analogue" mean an amide, ester, carbamate, carbonate, ureide, or phosphate analogue, respectively, that either: 1) does not destroy the biological activity of the compound and confers upon that compound advantageous properties *in vivo*, such as improved water solubility, improved circulating half-life in the blood (e.g., because of reduced metabolism of the prodrug), improved uptake, improved duration of action, or improved onset of action; or 2) is itself biologically inactive but is converted *in vivo* to a biologically active compound. Examples of biohydrolyzable amides include, but are not limited to, lower alkyl amides, α -amino acid amides, alkoxyacyl amides, and alkylaminoalkylcarbonyl amides. Examples of biohydrolyzable esters include, but are not limited to, lower alkyl esters, alkoxyacyloxy esters, alkyl acylamino alkyl esters, and choline esters. Examples of biohydrolyzable carbamates include, but are not limited to, lower alkylamines, substituted ethylenediamines, aminoacids, hydroxyalkylamines, heterocyclic and heteroaromatic amines, and polyether amines.

As used herein, "Hsp90" includes each member of the family of heat shock proteins having a mass of about 90-kiloDaltons. For example, in humans the highly conserved Hsp90 family includes cytosolic Hsp90 α and Hsp90 β isoforms, as well as GRP94, which is found in the endoplasmic reticulum, and HSP75/TRAP1, which is found in the mitochondrial matrix.

The term "c-kit" or "c-kit kinase" refers to a membrane receptor protein tyrosine kinase which is preferably activated upon binding Stem Cell Factor (SCF) to its extracellular domain (Yarden et al., 1987; Qiu et al., 1988). The full length amino acid sequence of a c-kit kinase preferably is as set forth in Yarden, *et al.*, 1987, *EMBO J.*, 11:3341-3351; and Qiu, *et al.*, 1988, *EMBO J.*, 7:1003-1011, which are incorporated by reference herein in their entirety, including any drawings. Mutant versions of c-kit kinase are encompassed by the term "c-kit kinase" and include those that fall into two classes: (1) having a single amino acid substitution at codon 816 of the human c-kit kinase, or its equivalent position in other species (Ma *et al.*, 1999, *J. Invest Dermatol.*, 112:165-170), and (2) those which have mutations involving the putative juxtamembrane α -helix of the protein (Ma, *et al.*, 1999, *J.*

Biol. Chem., 274:13399-13402). Both of these publications are incorporated by reference herein in their entirety, including any drawings.

As used herein, a “proliferative disorder” or a “hyperproliferative disorder,” and other equivalent terms, means a disease or medical condition involving
5 pathological growth of cells. Proliferative disorders include cancer, smooth muscle cell proliferation, systemic sclerosis, cirrhosis of the liver, adult respiratory distress syndrome, idiopathic cardiomyopathy, lupus erythematosus, retinopathy, e.g., diabetic retinopathy or other retinopathies, cardiac hyperplasia, reproductive system associated disorders such as benign prostatic hyperplasia and ovarian cysts,
10 pulmonary fibrosis, endometriosis, fibromatosis, hamartomas, lymphangiomatosis, sarcoidosis, desmoid tumors,

Smooth muscle cell proliferation includes hyperproliferation of cells in the vasculature, for example, intimal smooth muscle cell hyperplasia, restenosis and vascular occlusion, particularly stenosis following biologically- or mechanically-
15 mediated vascular injury, e.g., vascular injury associated with angioplasty. Moreover, intimal smooth muscle cell hyperplasia can include hyperplasia in smooth muscle other than the vasculature, e.g., bile duct blockage, bronchial airways of the lung in patients with asthma, in the kidneys of patients with renal interstitial fibrosis, and the like.

20 Non-cancerous proliferative disorders also include hyperproliferation of cells in the skin such as psoriasis and its varied clinical forms, Reiter's syndrome, pityriasis rubra pilaris, and hyperproliferative variants of disorders of keratinization (e.g., actinic keratosis, senile keratosis), scleroderma, and the like.

In a preferred embodiment, the proliferative disorder is cancer. Cancers that
25 can be treated or prevented by the methods of the present invention include, but are not limited to human sarcomas and carcinomas, e.g., fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma,
30 colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary

adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, retinoblastoma; leukemias, e.g., acute lymphocytic leukemia and acute myelocytic leukemia (myeloblastic, promyelocytic, myelomonocytic, monocytic and erythroleukemia); chronic leukemia (chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia); and polycythemia vera, lymphoma (Hodgkin's disease and non-Hodgkin's disease), multiple myeloma, Waldenstrom's macroglobulinemia, and heavy chain disease.

Other examples of leukemias include acute and/or chronic leukemias, e.g., lymphocytic leukemia (e.g., as exemplified by the p388 (murine) cell line), large granular lymphocytic leukemia, and lymphoblastic leukemia; T-cell leukemias, e.g., T-cell leukemia (e.g., as exemplified by the CEM, Jurkat, and HSB-2 (acute), YAC-1 (murine) cell lines), T-lymphocytic leukemia, and T-lymphoblastic leukemia; B cell leukemia (e.g., as exemplified by the SB (acute) cell line), and B-lymphocytic leukemia; mixed cell leukemias, e.g., B and T cell leukemia and B and T lymphocytic leukemia; myeloid leukemias, e.g., granulocytic leukemia, myelocytic leukemia (e.g., as exemplified by the HL-60 (promyelocyte) cell line), and myelogenous leukemia (e.g., as exemplified by the K562 (chronic) cell line); neutrophilic leukemia; eosinophilic leukemia; monocytic leukemia (e.g., as exemplified by the THP-1 (acute) cell line); myelomonocytic leukemia; Naegeli-type myeloid leukemia; and nonlymphocytic leukemia. Other examples of leukemias are described in Chapter 60 of *The Chemotherapy Sourcebook*, Michael C. Perry Ed., Williams & Williams (1992) and Section 36 of *Holland Frie Cancer Medicine* 5th Ed., Bast et al. Eds., B.C. Decker Inc. (2000). The entire teachings of the preceding references are incorporated herein by reference.

In one embodiment, the disclosed method is believed to be particularly effective in treating subject with non-solid tumors such as multiple myeloma. In another embodiment, the disclosed method is believed to be particularly effective

against T-leukemia (e.g., as exemplified by Jurkat and CEM cell lines); B-leukemia (e.g., as exemplified by the SB cell line); promyelocytes (e.g., as exemplified by the HL-60 cell line); uterine sarcoma (e.g., as exemplified by the MES-SA cell line); monocytic leukemia (e.g., as exemplified by the THP-1 (acute) cell line); and
5 lymphoma (e.g., as exemplified by the U937 cell line).

Some of the disclosed methods can be particularly effective at treating subjects whose cancer has become "multi-drug resistant". A cancer which initially responded to an anti-cancer drug becomes resistant to the anti-cancer drug when the anti-cancer drug is no longer effective in treating the subject with the cancer. For
10 example, many tumors will initially respond to treatment with an anti-cancer drug by decreasing in size or even going into remission, only to develop resistance to the drug. Drug resistant tumors are characterized by a resumption of their growth and/or reappearance after having seemingly gone into remission, despite the administration of increased dosages of the anti-cancer drug. Cancers that have
15 developed resistance to two or more anti-cancer drugs are said to be "multi-drug resistant". For example, it is common for cancers to become resistant to three or more anti-cancer agents, often five or more anti-cancer agents and at times ten or more anti-cancer agents.

As used herein, the term "c-kit associated cancer" refers to a cancer which
20 has aberrant expression and/or activation of c-kit. c-Kit associated cancers include leukemias, mast cell tumors, small cell lung cancer, testicular cancer, some cancers of the gastrointestinal tract and some central nervous system. In addition, c-kit has been implicated in playing a role in carcinogenesis of the female genital tract (Inoue, *et al.*, 1994, *Cancer Res.*, 54(11):3049-3053), sarcomas of neuroectodermal origin
25 (Ricotti, *et al.*, 1998, *Blood*, 91:2397-2405), and Schwann cell neoplasia associated with neurofibromatosis (Ryan, *et al.*, 1994, *J. Neuro. Res.*, 37:415-432).

As used herein, the term "pharmaceutically acceptable salt," is a salt formed from, for example, an acid and a basic group of one of the compounds of formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X), or Table 1. Illustrative salts
30 include, but are not limited, to sulfate, citrate, acetate, oxalate, chloride, bromide, iodide, nitrate, bisulfate, phosphate, acid phosphate, isonicotinate, lactate, salicylate, acid citrate, tartrate, oleate, tannate, pantothenate, bitartrate, ascorbate, succinate,

maleate, besylate, gentisinate, fumarate, gluconate, glucaronate, saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate, and pamoate (*i.e.*, 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)) salts. The term "pharmaceutically acceptable salt" also refers to a salt prepared from
5 a compound of formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X), or Table 1 having an acidic functional group, such as a carboxylic acid functional group, and a pharmaceutically acceptable inorganic or organic base. Suitable bases include, but are not limited to, hydroxides of alkali metals such as sodium, potassium, and lithium; hydroxides of alkaline earth metal such as calcium and
10 magnesium; hydroxides of other metals, such as aluminum and zinc; ammonia, and organic amines, such as unsubstituted or hydroxy-substituted mono-, di-, or trialkylamines; dicyclohexylamine; tributyl amine; pyridine; N-methyl,N-ethylamine; diethylamine; triethylamine; mono-, bis-, or tris-(2-hydroxy-lower alkyl amines), such as mono-, bis-, or tris-(2-hydroxyethyl)amine, 2-hydroxy-tert-
15 butylamine, or tris-(hydroxymethyl)methylamine, N, N,-di-lower alkyl-N-(hydroxy lower alkyl)-amines, such as N,N-dimethyl-N-(2-hydroxyethyl)amine, or tri-(2-hydroxyethyl)amine; N-methyl-D-glucamine; and amino acids such as arginine, lysine, and the like. The term "pharmaceutically acceptable salt" also refers to a salt prepared from a compound of formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII),
20 (IX), (X), or Table 1 having a basic functional group, such as an amine functional group, and a pharmaceutically acceptable inorganic or organic acid.. Suitable acids include, but are not limited to, hydrogen sulfate, citric acid, acetic acid, oxalic acid, hydrochloric acid (HCl), hydrogen bromide (HBr), hydrogen iodide (HI), nitric acid, hydrogen bisulfide, phosphoric acid, lactic acid, salicylic acid, tartaric acid,
25 bitartratic acid, ascorbic acid, succinic acid, maleic acid, besylic acid, fumaric acid, gluconic acid, glucaronic acid, formic acid, benzoic acid, glutamic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, and *p*-toluenesulfonic acid.

As used herein, the term "pharmaceutically acceptable solvate," is a solvate
30 formed from the association of one or more pharmaceutically acceptable solvent molecules to one of the compounds of formula (I), (II), (III), (IV), (V), (VI), (VII),

(VIII), (IX), (X), or Table 1. The term solvate includes hydrates (*e.g.*, hemihydrate, monohydrate, dihydrate, trihydrate, tetrahydrate, and the like).

A pharmaceutically acceptable carrier may contain inert ingredients which do not unduly inhibit the biological activity of the compounds. The

5 pharmaceutically acceptable carriers should be biocompatible, *i.e.*, non-toxic, non-inflammatory, non-immunogenic and devoid of other undesired reactions upon the administration to a subject. Standard pharmaceutical formulation techniques can be employed, such as those described in Remington's Pharmaceutical Sciences, *ibid*. Suitable pharmaceutical carriers for parenteral administration include, for example,
10 sterile water, physiological saline, bacteriostatic saline (saline containing about 0.9% mg/ml benzyl alcohol), phosphate-buffered saline, Hank's solution, Ringer's-lactate and the like. Methods for encapsulating compositions (such as in a coating of hard gelatin or cyclodextran) are known in the art (Baker, *et al.*, "Controlled Release of Biological Active Agents", John Wiley and Sons, 1986).

15 As used herein, the term "effective amount" refers to an amount of a compound of this invention which is sufficient to reduce or ameliorate the severity, duration, progression, or onset of a proliferative disorder, prevent the advancement of a proliferative disorder, cause the regression of a proliferative, prevent the recurrence, development, onset or progression of a symptom associated with a
20 proliferative disorder, or enhance or improve the prophylactic or therapeutic effect(s) of another therapy. The precise amount of compound administered to a subject will depend on the mode of administration, the type and severity of the disease or condition and on the characteristics of the subject, such as general health, age, sex, body weight and tolerance to drugs. It will also depend on the degree,
25 severity and type of cell proliferation, and the mode of administration. The skilled artisan will be able to determine appropriate dosages depending on these and other factors. When co-administered with other agents, *e.g.*, when co-administered with an anti-cancer agent, an "effective amount" of the second agent will depend on the type of drug used. Suitable dosages are known for approved agents and can be
30 adjusted by the skilled artisan according to the condition of the subject, the type of condition(s) being treated and the amount of a compound of the invention being

used. In cases where no amount is expressly noted, an effective amount should be assumed.

Non-limiting examples of an effective amount of a compound of the invention are provided herein below. In a specific embodiment, the invention provides a method of preventing, treating, managing, or ameliorating a proliferative disorder or one or more symptoms thereof, said methods comprising administering to a subject in need thereof a dose of at least 150 µg/kg, preferably at least 250 µg/kg, at least 500 µg/kg, at least 1 mg/kg, at least 5 mg/kg, at least 10 mg/kg, at least 25 mg/kg, at least 50 mg/kg, at least 75 mg/kg, at least 100 mg/kg, at least 125 mg/kg, at least 150 mg/kg, or at least 200 mg/kg or more of one or more compounds of the invention once every day, preferably, once every 2 days, once every 3 days, once every 4 days, once every 5 days, once every 6 days, once every 7 days, once every 8 days, once every 10 days, once every two weeks, once every three weeks, or once a month.

The dosages of a chemotherapeutic agents other than compounds of the invention, which have been or are currently being used to prevent, treat, manage, or ameliorate a proliferative disorder, or one or more symptoms thereof, can be used in the combination therapies of the invention. Preferably, dosages lower than those which have been or are currently being used to prevent, treat, manage, or ameliorate a proliferative disorder, or one or more symptoms thereof, are used in the combination therapies of the invention. The recommended dosages of agents currently used for the prevention, treatment, management, or amelioration of a proliferative disorder, or one or more symptoms thereof, can be obtained from any reference in the art including, but not limited to, Hardman *et al.*, eds., 1996, Goodman & Gilman's The Pharmacological Basis Of Therapeutics 9th Ed, Mc-Graw-Hill, New York; Physician's Desk Reference (PDR) 57th Ed., 2003, Medical Economics Co., Inc., Montvale, NJ, which are incorporated herein by reference in its entirety.

As used herein, the terms "treat", "treatment" and "treating" refer to the reduction or amelioration of the progression, severity and/or duration of a proliferative disorder, or the amelioration of one or more symptoms (preferably, one or more discernible symptoms) of a proliferative disorder resulting from the

administration of one or more therapies (*e.g.*, one or more therapeutic agents such as a compound of the invention). In specific embodiments, the terms “treat”, “treatment” and “treating” refer to the amelioration of at least one measurable physical parameter of a proliferative disorder, such as growth of a tumor, not necessarily discernible by the patient. In other embodiments the terms “treat”, “treatment” and “treating” refer to the inhibition of the progression of a proliferative disorder, either physically by, *e.g.*, stabilization of a discernible symptom, physiologically by, *e.g.*, stabilization of a physical parameter, or both. In other embodiments the terms “treat”, “treatment” and “treating” refer to the reduction or stabilization of tumor size or cancerous cell count.

As used herein, the terms “prevent”, “prevention” and “preventing” refer to the reduction in the risk of acquiring or developing a given proliferative disorder, or the reduction or inhibition of the recurrence of a proliferative disorder. In one embodiment, a compound of the invention is administered as a preventative measure to a patient, preferably a human, having a genetic predisposition to any of the disorders described herein.

As used herein, the terms “therapeutic agent” and “therapeutic agents” refer to any agent(s) which can be used in the treatment, management, or amelioration of a proliferative disorder or one or more symptoms thereof. In certain embodiments, the term “therapeutic agent” refers to a compound of the invention. In certain other embodiments, the term “therapeutic agent” refers does not refer to a compound of the invention. Preferably, a therapeutic agent is an agent which is known to be useful for, or has been or is currently being used for the treatment, management, prevention, or amelioration a proliferative disorder or one or more symptoms thereof.

As used herein, the term “synergistic” refers to a combination of a compound of the invention and another therapy (*e.g.*, a prophylactic or therapeutic agent), which is more effective than the additive effects of the therapies. A synergistic effect of a combination of therapies (*e.g.*, a combination of prophylactic or therapeutic agents) permits the use of lower dosages of one or more of the therapies and/or less frequent administration of said therapies to a subject with a proliferative disorder. The ability to utilize lower dosages of a therapy (*e.g.*, a prophylactic or

therapeutic agent) and/or to administer said therapy less frequently reduces the toxicity associated with the administration of said therapy to a subject without reducing the efficacy of said therapy in the prevention, management or treatment of a proliferative disorder. In addition, a synergistic effect can result in improved
5 efficacy of agents in the prevention, management or treatment of a proliferative disorder. Finally, a synergistic effect of a combination of therapies (*e.g.*, a combination of prophylactic or therapeutic agents) may avoid or reduce adverse or unwanted side effects associated with the use of either therapy alone.

As used herein, the phrase "side effects" encompasses unwanted and adverse
10 effects of a therapy (*e.g.*, a prophylactic or therapeutic agent). Side effects are always unwanted, but unwanted effects are not necessarily adverse. An adverse effect from a therapy (*e.g.*, prophylactic or therapeutic agent) might be harmful or uncomfortable or risky. Side effects include, but are not limited to fever, chills, lethargy, gastrointestinal toxicities (including gastric and intestinal ulcerations and
15 erosions), nausea, vomiting, neurotoxicities, nephrotoxicities, renal toxicities (including such conditions as papillary necrosis and chronic interstitial nephritis), hepatic toxicities (including elevated serum liver enzyme levels), myelotoxicities (including leukopenia, myelosuppression, thrombocytopenia and anemia), dry mouth, metallic taste, prolongation of gestation, weakness, somnolence, pain
20 (including muscle pain, bone pain and headache), hair loss, asthenia, dizziness, extra-pyramidal symptoms, akathisia, cardiovascular disturbances and sexual dysfunction.

As used herein, the term "in combination" refers to the use of more than one therapies (*e.g.*, one or more prophylactic and/or therapeutic agents). The use of the
25 term "in combination" does not restrict the order in which therapies (*e.g.*, prophylactic and/or therapeutic agents) are administered to a subject with a proliferative disorder. A first therapy (*e.g.*, a prophylactic or therapeutic agent such as a compound of the invention) can be administered prior to (*e.g.*, 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24
30 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concomitantly with, or subsequent to (*e.g.*, 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12

hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of a second therapy (e.g., a prophylactic or therapeutic agent such as an anti-cancer agent) to a subject with a proliferative disorder, such as cancer.

5 As used herein, the terms “therapies” and “therapy” can refer to any protocol(s), method(s), and/or agent(s) that can be used in the prevention, treatment, management, or amelioration of a proliferative disorder or one or more symptoms thereof.

 A used herein, a “protocol” includes dosing schedules and dosing regimens.

10 The protocols herein are methods of use and include prophylactic and therapeutic protocols.

 As used herein, the terms “manage,” “managing,” and “management” refer to the beneficial effects that a subject derives from a therapy (e.g., a prophylactic or therapeutic agent), which does not result in a cure of the disease. In certain
15 embodiments, a subject is administered one or more therapies (e.g., one or more prophylactic or therapeutic agents) to “manage” a disease so as to prevent the progression or worsening of the disease.

 As used herein, a composition that “substantially” comprises a compound means that the composition contains more than about 80% by weight, more
20 preferably more than about 90% by weight, even more preferably more than about 95% by weight, and most preferably more than about 97% by weight of the compound.

 As used herein, a reaction that is “substantially complete” means that the reaction contains more than about 80% by weight of the desired product, more
25 preferably more than about 90% by weight of the desired product, even more preferably more than about 95% by weight of the desired product, and most preferably more than about 97% by weight of the desired product.

 As used herein, a racemic mixture means about 50% of one enantiomer and about 50% of is corresponding enantiomer relative to a chiral center in the molecule.

30 The invention encompasses all enantiomerically-pure, enantiomerically-enriched, diastereomerically pure, diastereomerically enriched, and racemic mixtures of the compounds of the invention.

Enantiomeric and diastereomeric mixtures can be resolved into their component enantiomers or diastereomers by well known methods, such as chiral-phase gas chromatography, chiral-phase high performance liquid chromatography, crystallizing the compound as a chiral salt complex, or crystallizing the compound in a chiral solvent. Enantiomers and diastereomers can also be obtained from diastereomerically- or enantiomerically-pure intermediates, reagents, and catalysts by well known asymmetric synthetic methods.

The compounds of the invention are defined herein by their chemical structures and/or chemical names. Where a compound is referred to by both a chemical structure and a chemical name, and the chemical structure and chemical name conflict, the chemical structure is determinative of the compound's identity.

When administered to a patient, *e.g.*, to a non-human animal for veterinary use or for improvement of livestock, or to a human for clinical use, the compounds of the invention are administered in isolated form or as the isolated form in a pharmaceutical composition. As used herein, "isolated" means that the compounds of the invention are separated from other components of either (a) a natural source, such as a plant or cell, preferably bacterial culture, or (b) a synthetic organic chemical reaction mixture. Preferably, the compounds of the invention are purified via conventional techniques. As used herein, "purified" means that when isolated, the isolate contains at least 95%, preferably at least 98%, of a compound of the invention by weight of the isolate either as a mixture of stereoisomers or as a diastereomeric or enantiomeric pure isolate.


As used herein, a composition that is "substantially free" of a compound means that the composition contains less than about 20% by weight, more preferably less than about 10% by weight, even more preferably less than about 5% by weight, and most preferably less than about 3% by weight of the compound.

Only those choices and combinations of substituents that result in a stable structure are contemplated. Such choices and combinations will be apparent to those of ordinary skill in the art and may be determined without undue experimentation.

The invention can be understood more fully by reference to the following detailed description and illustrative examples, which are intended to exemplify non-limiting embodiments of the invention.

B. The Compounds of the Invention

The present invention encompasses compounds having Formulas (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X), and those set forth in Table 1 and
5 tautomers, pharmaceutically acceptable salts, solvates, clathrates, hydrates, polymorphs and prodrugs thereof. In one aspect, the invention provides compounds of formula (I) as set forth below:




(I)

and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof, wherein ring A, R₁, R₃ and R₅ are defined as above.

Compounds of formula (I) inhibit the activity of Hsp90 and are particularly useful for treating or preventing proliferative disorders, such as cancer. In addition,
15 compounds of formula (I) are particularly useful in treating cancer when given in combination with other anti-cancer agent.

In one embodiment, in the compounds of formula (I), R₅ is an optionally substituted naphthyl.

In another embodiment, in the compounds of formula (I), R₅ is represented
20 by the following formula:



wherein:


R₉, for each occurrence, is independently a substituent selected from the group consisting of an optionally substituted alkyl, an optionally substituted alkenyl,

an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, hydroxyalkyl, alkoxyalkyl, halo, cyano, nitro, guanadino, a haloalkyl, a heteroalkyl, $-NR_{10}R_{11}$, $-OR_7$, $-C(O)R_7$, $-C(O)OR_7$, $-OC(O)R_7$, $-C(O)NR_{10}R_{11}$, $-NR_8C(O)R_7$, $-SR_7$, $-S(O)_pR_7$, $-OS(O)_pR_7$, $-S(O)_pOR_7$, $-NR_8S(O)_pR_7$, or $-S(O)_pNR_{10}R_{11}$, $-S(O)_pOR_7$, $-OP(O)(OR_7)_2$, or $-SP(O)(OR_7)_2$;

or two R_9 groups taken together with the carbon atoms to which they are attached form a fused ring; and

m is zero or an integer from 1 to 7, wherein R_7 , R_8 , R_{10} , R_{11} , and p are defined as above.

In another embodiment, in the compounds represented by formula (I), R_5 is represented by one of the following formulas:







wherein R_9 is defined as above;

q is zero or an integer from 1 to 7; and



u is zero or an integer from 1 to 8.

In another embodiment, in the compounds represented by formula (I), R_5 is selected from the group consisting of:






5



10

and



wherein:

X_6 , for each occurrence, is independently CH, CR_9 , N, N(O), $N^+(R_{17})$,
provided that at least three X_6 groups are independently selected from CH and CR_9 ;

5 X_7 , for each occurrence, is independently CH, CR_9 , N, N(O), $N^+(R_{17})$,
provided that at least three X_7 groups are independently selected from CH and CR_9 ;

X_8 , for each occurrence, is independently CH_2 , CHR_9 , CR_9R_9 , O, S, $S(O)_p$,
 NR_7 , or NR_{17} ;

X_9 , for each occurrence, is independently N or CH;


10 X_{10} , for each occurrence, is independently CH, CR_9 , N, N(O), $N^+(R_{17})$,
provided that at least one X_{10} is selected from CH and CR_9 ;

R_{17} , for each occurrence, is independently -H, an alkyl, an aralkyl, $-C(O)R_7$,
 $-C(O)OR_7$, or $-C(O)NR_{10}R_{11}$; wherein R_7 , R_9 , R_{10} , R_{11} and p are defined as above.

In another embodiment, in the compounds represented by formula (I), R_5 is
15 an optionally substituted indolyl, an optionally substituted benzoimidazolyl, an
optionally substituted indazolyl, an optionally substituted 3H-indazolyl, an
optionally substituted indolizynyl, an optionally substituted quinolynyl, an optionally
substituted isoquinolynyl, an optionally substituted benzoxazolyl, an optionally
substituted benzo[1,3]dioxolyl, an optionally substituted benzofuryl, an optionally
20 substituted benzothiazolyl, an optionally substituted benzo[d]isoxazolyl, an
optionally substituted benzo[d]isothiazolyl, an optionally substituted thiazolo[4,5-
c]pyridinyl, an optionally substituted thiazolo[5,4-c]pyridinyl, an optionally
substituted thiazolo[4,5-b]pyridinyl, an optionally substituted thiazolo[5,4-
b]pyridinyl, an optionally substituted oxazolo[4,5-c]pyridinyl, an optionally
25 substituted oxazolo[5,4-c]pyridinyl, an optionally substituted oxazolo[4,5-
b]pyridinyl, an optionally substituted oxazolo[5,4-b]pyridinyl, an optionally
substituted imidazopyridinyl, an optionally substituted benzothiadiazolyl,
benzoxadiazolyl, an optionally substituted benzotriazolyl, an optionally substituted

tetrahydroindolyl, an optionally substituted azaindolyl, an optionally substituted quinazolinyl, an optionally substituted purinyl, an optionally substituted imidazo[4,5-a]pyridinyl, an optionally substituted imidazo[1,2-a]pyridinyl, an optionally substituted 3*H*-imidazo[4,5-b]pyridinyl, an optionally substituted 1*H*-imidazo[4,5-b]pyridinyl, an optionally substituted 1*H*-imidazo[4,5-c]pyridinyl, an optionally substituted 3*H*-imidazo[4,5-c]pyridinyl, an optionally substituted pyridopyrdazinyl, and optionally substituted pyridopyrimidinyl, an optionally substituted pyrrolo[2,3]pyrimidyl, an optionally substituted pyrazolo[3,4]pyrimidyl, an optionally substituted cyclopentaimidazolyl, an optionally substituted cyclopentatriazolyl, an optionally substituted pyrrolopyrazolyl, an optionally substituted pyrroloimidazolyl, an optionally substituted pyrrolotriazolyl, or an optionally substituted benzo(b)thienyl.

In another embodiment, in the compounds represented by formula (I), R_5 is an optionally substituted indolyl. Preferably, R_5 is an indolyl represented by the following structural formula:




wherein:

R_{33} is a halo, lower alkyl, a lower alkoxy, a lower haloalkyl, a lower haloalkoxy, and lower alkyl sulfanyl;

R_{34} is H, a lower alkyl, or a lower alkylcarbonyl; and

Ring B and Ring C are optionally substituted with one or more substituents.

In another embodiment, in the compounds represented by formula (I), R_5 is selected from the group consisting of:




wherein:

X_{11} , for each occurrence, is independently CH, CR_9 , N, N(O), or $N^+(R_{17})$, provided that at least one X_{11} is N, N(O), or $N^+(R_{17})$ and at least two X_{11} groups are independently selected from CH and CR_9 ;

X_{12} , for each occurrence, is independently CH, CR_9 , N, N(O), $N^+(R_{17})$, provided that at least one X_{12} group is independently selected from CH and CR_9 ;

X_{13} , for each occurrence, is independently O, S, S(O)_p, NR_7 , or NR_{17} ; wherein R_7 , R_9 and R_{17} are defined as above.

In another embodiment, in compounds represented by formula (I), or any of the embodiments of formula (I) in which particular groups are disclosed, the compound is represented by the following structural formula:




wherein R_1 , R_3 , and R_5 are defined as above; and

R_6 , for each occurrence, is independently an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteroaralkyl, halo, cyano, nitro, guanadino, a haloalkyl, a heteroalkyl, alkoxy, haloalkoxy, $-NR_{10}R_{11}$, $-OR_7$, $-C(O)R_7$, $-C(O)OR_7$, $-C(S)R_7$, $-C(O)SR_7$, $-C(S)SR_7$, $-C(S)OR_7$, $-C(S)NR_{10}R_{11}$, $-C(NR_8)OR_7$, $-C(NR_8)R_7$, $-C(NR_8)NR_{10}R_{11}$, $-C(NR_8)SR_7$, $-OC(O)R_7$, $-OC(O)OR_7$, $-OC(S)OR_7$, $-OC(NR_8)OR_7$, $-SC(O)R_7$, $-SC(O)OR_7$, $-SC(NR_8)OR_7$,

- OC(S)R₇, -SC(S)R₇, -SC(S)OR₇, -OC(O)NR₁₀R₁₁, -OC(S)NR₁₀R₁₁,
 -OC(NR₈)NR₁₀R₁₁, -SC(O)NR₁₀R₁₁, -SC(NR₈)NR₁₀R₁₁, -SC(S)NR₁₀R₁₁,
 -OC(NR₈)R₇, -SC(NR₈)R₇, -C(O)NR₁₀R₁₁, -NR₈C(O)R₇, -NR₇C(S)R₇,
 -NR₇C(S)OR₇, -NR₇C(NR₈)R₇, -NR₇C(O)OR₇, -NR₇C(NR₈)OR₇,
 5 -NR₇C(O)NR₁₀R₁₁, -NR₇C(S)NR₁₀R₁₁, -NR₇C(NR₈)NR₁₀R₁₁, -SR₇, -S(O)_pR₇,
 -OS(O)_pR₇, -OS(O)_pOR₇, -OS(O)_pNR₁₀R₁₁, -S(O)_pOR₇, -NR₈S(O)_pR₇,
 -NR₇S(O)_pNR₁₀R₁₁, -NR₇S(O)_pOR₇, -S(O)_pNR₁₀R₁₁, -SS(O)_pR₇, -SS(O)_pOR₇,
 -SS(O)_pNR₁₀R₁₁, -OP(O)(OR₇)₂, or -SP(O)(OR₇)₂; and

n is zero or an integer from 1 to 4, wherein R₇, R₈, R₁₀, R₁₁, and p are defined
 10 as above.

In another embodiment, in compounds represented by formula (I), or any of
 the embodiments of formula (I) in which particular groups are disclosed, the
 compound is represented by the following structural formula:



15

wherein R₁, R₃, R₅, and R₆ are defined as above; and

- R₂₅ is an optionally substituted alkyl, an optionally substituted alkenyl, an
 optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally
 substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally
 20 substituted aryl, an optionally substituted heteroaryl, an optionally substituted
 aralkyl, an optionally substituted heteroaralkyl, halo, cyano, nitro, guanidino, a
 haloalkyl, a heteroalkyl, alkoxy, haloalkoxy, -NR₁₀R₁₁, -OR₇, -C(O)R₇, -C(O)OR₇,
 -C(S)R₇, -C(O)SR₇, -C(S)SR₇, -C(S)OR₇, -C(S)NR₁₀R₁₁, -C(NR₈)OR₇, -C(NR₈)R₇,
 -C(NR₈)NR₁₀R₁₁, -C(NR₈)SR₇, -OC(O)R₇, -OC(O)OR₇, -OC(S)OR₇, -OC(NR₈)OR₇,
 25 -SC(O)R₇, -SC(O)OR₇, -SC(NR₈)OR₇, -OC(S)R₇, -SC(S)R₇, -SC(S)OR₇,
 -OC(O)NR₁₀R₁₁, -OC(S)NR₁₀R₁₁, -OC(NR₈)NR₁₀R₁₁, -SC(O)NR₁₀R₁₁,
 -SC(NR₈)NR₁₀R₁₁, -SC(S)NR₁₀R₁₁, -OC(NR₈)R₇, -SC(NR₈)R₇, -C(O)NR₁₀R₁₁,

- NR₈C(O)R₇, -NR₇C(S)R₇, -NR₇C(S)OR₇, -NR₇C(NR₈)R₇, -NR₇C(O)OR₇,
 -NR₇C(NR₈)OR₇, -NR₇C(O)NR₁₀R₁₁, -NR₇C(S)NR₁₀R₁₁, -NR₇C(NR₈)NR₁₀R₁₁,
 -SR₇, -S(O)_pR₇, -OS(O)_pR₇, -OS(O)_pOR₇, -OS(O)_pNR₁₀R₁₁, -S(O)_pOR₇,
 -NR₈S(O)_pR₇, -NR₇S(O)_pNR₁₀R₁₁, -NR₇S(O)_pOR₇, -S(O)_pNR₁₀R₁₁, -SS(O)_pR₇,
 5 -SS(O)_pOR₇, -SS(O)_pNR₁₀R₁₁, -OP(O)(OR₇)₂, or -SP(O)(OR₇)₂;
 k is 1, 2, 3, or 4; and
 r is zero or an integer from 1 to 3, wherein R₇, R₈, R₁₀, R₁₁, and p are defined
 as above.


- In another embodiment of the compound represented by the above formula,
 10 R₁, R₃ and R₂₅ are each independently -OH, -SH, -NHR₇, -OC(O)NR₁₀R₁₁,
 -SC(O)NR₁₀R₁₁, -OC(O)R₇, -SC(O)R₇, -OC(O)OR₇, -SC(O)OR₇, -OS(O)_pR₇,
 -S(O)_pOR₇, -SS(O)_pR₇, -OS(O)_pOR₇, -SS(O)_pOR₇, -OC(S)R₇, -SC(S)R₇, -OC(S)OR₇,
 -SC(S)OR₇, -OC(S)NR₁₀R₁₁, -SC(S)NR₁₀R₁₁, -OC(NR₈)R₇, -SC(NR₈)R₇,
 -OC(NR₈)OR₇, -SC(NR₈)OR₇, -OP(O)(OR₇)₂ or -SP(O)(OR₇)₂.
 15 In another embodiment of the compound represented by the above formula,
 R₁ and R₃ are each, independently, -OH, -SH, or -NHR₇. In this case, R₆ can be an
 optionally substituted alkyl, an optionally substituted alkenyl, an optionally
 substituted alkynyl, cyano, halo, nitro, an optionally substituted cycloalkyl,
 haloalkyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an
 20 optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally
 substituted heteroaralkyl, -OR₇, -SR₇, -NR₁₀R₁₁, -OC(O)NR₁₀R₁₁,
 -SC(O)NR₁₀R₁₁, -NR₇C(O)NR₁₀R₁₁, -OC(O)R₇, -SC(O)R₇, -NR₇C(O)R₇,
 -OC(O)OR₇, -SC(O)OR₇, -NR₇C(O)OR₇, -OCH₂C(O)R₇, -SCH₂C(O)R₇,
 -NR₇CH₂C(O)R₇, -OCH₂C(O)OR₇, -SCH₂C(O)OR₇, -NR₇CH₂C(O)OR₇,
 25 -OCH₂C(O)NR₁₀R₁₁, -SCH₂C(O)NR₁₀R₁₁, -NR₇CH₂C(O)NR₁₀R₁₁, -OS(O)_pR₇,
 -SS(O)_pR₇, -NR₇S(O)_pR₇, -OS(O)_pNR₁₀R₁₁, -SS(O)_pNR₁₀R₁₁, -NR₇S(O)_pNR₁₀R₁₁,
 -OS(O)_pOR₇, -SS(O)_pOR₇, -NR₇S(O)_pOR₇, -OC(S)R₇, -SC(S)R₇, -NR₇C(S)R₇,
 -OC(S)OR₇, -SC(S)OR₇, -NR₇C(S)OR₇, -OC(S)NR₁₀R₁₁, -SC(S)NR₁₀R₁₁,
 -NR₇C(S)NR₁₀R₁₁, -OC(NR₈)R₇, -SC(NR₈)R₇, -NR₇C(NR₈)R₇, -OC(NR₈)OR₇,
 30 -SC(NR₈)OR₇, -NR₇C(NR₈)OR₇, -OC(NR₈)NR₁₀R₁₁, -SC(NR₈)NR₁₀R₁₁,
 -NR₇C(NR₈)NR₁₀R₁₁, -C(O)R₇, -C(O)OR₇, -C(O)NR₁₀R₁₁, -C(O)SR₇, -C(S)R₇,

-C(S)OR₇, -C(S)NR₁₀R₁₁, -C(S)SR₇, -C(NR₈)OR₇, -C(NR₈)R₇, -C(NR₈)NR₁₀R₁₁,
-C(NR₈)SR₇, -S(O)_pOR₇, -S(O)_pNR₁₀R₁₁, or -S(O)_pR₇.

In another embodiment of the above compound, R₁ is -SH or -OH; R₃ and R₂₅ are -OH; R₆ is a lower alkyl, C3-C6 cycloalkyl, lower alkoxy, a lower alkyl
5 sulfanyl, or -NR₁₀R₁₁; and R₉, for each occurrence, is independently selected from the group consisting of -OH, -SH, halo, a lower haloalkyl, cyano, a lower alkyl, a lower alkoxy, and a lower alkyl sulfanyl.

In another embodiment, in compounds represented by formula (I), or any of the embodiments of formula (I) in which particular groups are disclosed, R₁ and R₃
10 are each, independently, -OH, -SH, or -NHR₇.

In another embodiment, in compounds represented by formula (I), or any of the embodiments of formula (I) in which particular groups are disclosed, the compound is represented by the following structural formula:





15 wherein R₁, R₃, R₅, and R₂₅ are defined as above; and

R₆ is an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, cyano, halo, nitro, an optionally substituted cycloalkyl, haloalkyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted
20 aralkyl, an optionally substituted heteroaralkyl, -OR₇, -SR₇, -NR₁₀R₁₁, -OC(O)NR₁₀R₁₁, -SC(O)NR₁₀R₁₁, -NR₇C(O)NR₁₀R₁₁, -OC(O)R₇, -SC(O)R₇, -NR₇C(O)R₇, -OC(O)OR₇, -SC(O)OR₇, -NR₇C(O)OR₇, -OCH₂C(O)R₇, -SCH₂C(O)R₇, -NR₇CH₂C(O)R₇, -OCH₂C(O)OR₇, -SCH₂C(O)OR₇, -NR₇CH₂C(O)OR₇, -OCH₂C(O)NR₁₀R₁₁, -SCH₂C(O)NR₁₀R₁₁,
25 -NR₇CH₂C(O)NR₁₀R₁₁, -OS(O)_pR₇, -SS(O)_pR₇, -NR₇S(O)_pR₇, -OS(O)_pNR₁₀R₁₁, -SS(O)_pNR₁₀R₁₁, -NR₇S(O)_pNR₁₀R₁₁, -OS(O)_pOR₇, -SS(O)_pOR₇, -NR₇S(O)_pOR₇, -OC(S)R₇, -SC(S)R₇, -NR₇C(S)R₇, -OC(S)OR₇, -SC(S)OR₇, -NR₇C(S)OR₇,

- OC(S)NR₁₀R₁₁, -SC(S)NR₁₀R₁₁, -NR₇C(S)NR₁₀R₁₁, -OC(NR₈)R₇, -SC(NR₈)R₇,
 -NR₇C(NR₈)R₇, -OC(NR₈)OR₇, -SC(NR₈)OR₇, -NR₇C(NR₈)OR₇,
 -OC(NR₈)NR₁₀R₁₁, -SC(NR₈)NR₁₀R₁₁, -NR₇C(NR₈)NR₁₀R₁₁, -C(O)R₇,
 -C(O)OR₇, -C(O)NR₁₀R₁₁, -C(O)SR₇, -C(S)R₇, -C(S)OR₇, -C(S)NR₁₀R₁₁,
 5 -C(S)SR₇, -C(NR₈)OR₇, -C(NR₈)R₇, -C(NR₈)NR₁₀R₁₁, -C(NR₈)SR₇, -S(O)_pOR₇,
 -S(O)_pNR₁₀R₁₁, or -S(O)_pR₇, wherein R₇, R₈, R₁₀, R₁₁, and p are defined as above.
 In a preferred embodiment, R₁ is -SH or -OH; R₃ and R₂₅ are -OH; R₁₂ is a lower
 alkyl, lower alkoxy, a lower alkyl sulfanyl, or -NR₁₀R₁₁; and R₉, for each
 occurrence, is independently selected from the group consisting of -OH, -SH, halo,
 10 a lower haloalkyl, cyano, a lower alkyl, a lower alkoxy, and a lower alkyl sulfanyl.

In another embodiment, in compounds represented by formula (I), or any of
 the embodiments of formula (I) in which particular groups are disclosed, the
 compound is represented by one of the following structural formulas:



15


wherein R₁, R₃, R₅, R₆ and n are as defined above; and

X₃ and X₄ are each, independently, N, N(O), N⁺(R₁₇), CH or CR₆; and


X₅ is O, S, NR₁₇, CH=CH, CH=CR₆, CR₆=CH, CR₆=CR₆, CH=N, CR₆=N,
 CH=N(O), CR₆=N(O), N=CH, N=CR₆, N(O)=CH, N(O)=CR₆, N⁺(R₁₇)=CH,

- 20 N⁺(R₁₇)=CR₆, CH=N⁺(R₁₇), CR₆=N⁺(R₁₇), or N=N; wherein R₁₇ is defined as above.


In another embodiment, in compounds represented by formula (I), or any of
 the embodiments of formula (I) in which particular groups are disclosed, the
 compound is selected from the group consisting of:




,




;




,




5 ;




,



;




,




10

;



wherein R_1 , R_3 , R_5 , and R_{25} are defined as above.

In another aspect, the invention provides compounds of formula (II) as set forth below:



and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof, wherein ring A, R_1 and R_3 are defined as above; and

R_2 is a substituted phenyl, wherein the phenyl group is substituted with:

- i) one substituent selected from nitro, cyano, a haloalkoxy, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, hydroxylalkyl, alkoxyalkyl, guanadino, $-NR_{10}R_{11}$, $-O-R_{20}$, $-C(O)R_7$, $-C(O)OR_{20}$, $-OC(O)R_7$, $-C(O)NR_{10}R_{11}$, $-NR_8C(O)R_7$, $-SR_7$, $-S(O)_pR_7$, $-OS(O)_pR_7$, $-S(O)_pOR_7$, $-NR_8S(O)_pR_7$, or $-S(O)_pNR_{10}R_{11}$, or
- ii) two to five substituents selected from the group consisting of an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl,

an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, hydroxyalkyl, alkoxyalkyl, -F, -Br, -I, cyano, nitro, 5 guanadino, a haloalkyl, a heteroalkyl, -NR₁₀R₁₁, -OR₇, -C(O)R₇, -C(O)OR₇, -OC(O)R₇, -C(O)NR₁₀R₁₁, -NR₈C(O)R₇, -SR₇, -S(O)_pR₇, -OS(O)_pR₇, -S(O)_pOR₇, -NR₈S(O)_pR₇, or -S(O)_pNR₁₀R₁₁;


R₂₀, for each occurrence, is independently an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

p, for each occurrence, is, independently, 1 or 2.

15 Compounds of formula (II) inhibit the activity of Hsp90 and are particularly useful for treating or preventing proliferative disorders, such as cancer. In addition, compounds of formula (II) are particularly useful in treating cancer when given in combination with other anti-cancer agent.


In one embodiment, the compounds represented by formula (II) do not 20 include 3-(2,4-dihydroxy-phenyl)-4-(7-naphthalen-1-yl)-5-mercapto-triazole, 3-(2,4-dihydroxyphenyl)-4-(2,5-dimethoxyphenyl)-5-mercapto-triazole, 3-(1-phenyl-5-amino-pyrazol-4-yl)-4-(2,4-dichlorophenyl)-5-mercapto-triazole, and 3-(2-hydroxyphenyl)-4-(2,4-dimethylphenyl)-5-mercapto-triazole.

In another embodiment, in compounds represented by formula (II), or any of 25 the embodiments of formula (II) in which particular groups are disclosed, the compound is represented by the following structural formula:



wherein R_1 , R_2 , R_3 , R_6 , and n are defined as above.


In another embodiment, in compounds represented by formula (II), or any of the embodiments of formula (II) in which particular groups are disclosed, the compound is represented by the following structural formula:



wherein R_1 , R_2 , R_3 , R_6 , R_{25} and r are defined as above.



In another embodiment, in compounds represented by formula (II), or any of the embodiments of formula (II) in which particular groups are disclosed, R_1 and R_3 are each, independently, -OH, -SH, or -NHR₇.

In another embodiment, in compounds represented by formula (II), or any of the embodiments of formula (II) in which particular groups are disclosed, the compound is represented by the following structural formula:





wherein R_1 , R_2 , R_3 , R_6 and R_{25} are defined as above. In a preferred embodiment, R_1 is -SH or -OH; R_3 and R_{25} are -OH; R_{12} is a lower alkyl, lower alkoxy, a lower alkyl sulfanyl, or -NR₁₀R₁₁; and R_9 , for each occurrence, is independently selected from the group consisting of -OH, -SH, halo, a lower haloalkyl, cyano, a lower alkyl, a lower alkoxy, and a lower alkyl sulfanyl.

In another embodiment, in compounds represented by formula (II), or any of the embodiments of formula (II) in which particular groups are disclosed, the compound is represented by one of the following structural formulas:







wherein R_1 , R_2 , R_3 , R_6 , X_3 , X_4 , X_5 and n are defined as above.


- 5 In another embodiment, in compounds represented by formula (II), or any of the embodiments of formula (II) in which particular groups are disclosed, the compound is selected from the group consisting of:




10



15




;



5 ;

wherein R₁, R₂, R₃, and R₂₅ are defined as above.

In another aspect, the invention provides compounds of formula (III) as set forth below:



10

(III)

and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs.

In formula (III), ring A, R₁, and R₃ are defined as above; and

15 R₁₈ is an optionally substituted cycloalkyl, and optionally substituted cycloalkenyl, or a substituted alkyl, wherein the alkyl group is substituted with one or more substituents independently selected from the group consisting of an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, halo, cyano, nitro,
 20 guanadino, a haloalkyl, -NR₁₀R₁₁, -OR₇, -C(O)R₇, -C(O)OR₇, -OC(O)R₇,

-C(O)NR₁₀R₁₁, -NR₈C(O)R₇, -SR₇, -S(O)_pR₇, -OS(O)_pR₇, -S(O)_pOR₇,
-NR₈S(O)_pR₇, or -S(O)_pNR₁₀R₁₁, wherein R₇, R₈, R₁₀, R₁₁, and p are defined as
above.

Compounds of formula (III) inhibit the activity of Hsp90 and are particularly
5 useful for treating or preventing proliferative disorders, such as cancer. In addition,
compounds of formula (III) are particularly useful in treating cancer when given in
combination with other anti-cancer agent.


In one embodiment, in formula (III) R₁₈ is not cyclohexyl.

In another embodiment, in formula (III) R₁₈ is an optionally substituted
10 cycloalkyl or an optionally substituted cycloalkenyl.

In another embodiment, in formula (III) R₁₈ is a substituted alkyl.


In another embodiment, in compounds represented by formula (III), or any of
the embodiments of formula (III) in which particular groups are disclosed, the
compound is represented by the following structural formula:

15



wherein R₁, R₃, R₆, R₁₈, and n are defined as above.


In another embodiment, in compounds represented by formula (III), or any of
20 the embodiments of formula (III) in which particular groups are disclosed, the
compound is represented by the following structural formula:



wherein R_1 , R_3 , R_6 , R_{18} , R_{25} and r are defined as above.


In another embodiment, in compounds represented by formula (III), or any of the embodiments of formula (III) in which particular groups are disclosed, R_1 and R_3 are each, independently, -OH, -SH, or -NHR₇.

- 5 In another embodiment, in compounds represented by formula (III), or any of the embodiments of formula (III) in which particular groups are disclosed, the compound is represented by the following structural formula:




- 10 wherein R_1 , R_3 , R_6 , R_{18} , and R_{25} are defined as above. In a preferred embodiment, R_1 is -SH or -OH; R_3 and R_{25} are -OH; and R_{12} is a lower alkyl, lower alkoxy, a lower alkyl sulfanyl, or -NR₁₀R₁₁.

In another embodiment, in compounds represented by formula (III), or any of the embodiments of formula (III) in which particular groups are disclosed, the compound is represented by one of the following structural formulas:





wherein R_1 , R_3 , R_6 , R_{18} , X_3 , X_4 , X_5 , and n are defined as above.

- 20 In another embodiment, in compounds represented by formula (III), or any of the embodiments of formula (III) in which particular groups are disclosed, the compound is selected from the group consisting of:





;

,





5 ;

,



;


,




10 ;


;

,






and




wherein R_1 , R_3 , R_{18} , and R_{25} are defined as above.

In another aspect, the invention provides compounds of formula (IV) or (V)

as set forth below:



(IV)



(V)

and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof. In formulas (IV) and (V), R_1 and R_3 are as defined above; and

X_{14} is O, S, or NR_7 ;

R_{21} is an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

R_{22} , for each occurrence, is independently an -H or is selected from the group consisting of an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl, a haloalkyl, $-C(O)R_7$, $-C(O)OR_7$,

-OC(O)R₇, -C(O)NR₁₀R₁₁, -NR₈C(O)R₇, -S(O)_pR₇, -S(O)_pOR₇, or -S(O)_pNR₁₀R₁₁; and

R₂₃ and R₂₄, for each occurrence, are independently -H or are selected from the group consisting of an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl, halo, cyano, nitro, guanadino, a haloalkyl, a heteroalkyl, -NR₁₀R₁₁, -OR₇, -C(O)R₇, -C(O)OR₇, -OC(O)R₇, -C(O)NR₁₀R₁₁, -NR₈C(O)R₇, -SR₇, -S(O)_pR₇, -OS(O)_pR₇, -S(O)_pOR₇, -NR₈S(O)_pR₇, or -S(O)_pNR₁₀R₁₁;

wherein R₇, R₈, R₁₀, R₁₁ and p are defined as above.

In one embodiment, in formulas (IV) and (V), R₂₁ is an optionally substituted alkyl, an optionally substituted cycloalkyl, an optionally substituted aryl or an optionally substituted heteroaryl.


In another embodiment, in the formulas (IV) and (V), R₁ is -OH, -SH, or -NHR₇.

In another embodiment, in the formulas (IV) and (V), R₂₂ is -H, an alkyl, an aralkyl, -C(O)R₇, -C(O)OR₇, or -C(O)NR₁₀R₁₁.

In another embodiment, in the formulas (IV) and (V), X₁₄ is O.

Compounds of formula (IV) or (V) inhibit the activity of Hsp90 and are particularly useful for treating or preventing proliferative disorders, such as cancer. In addition, compounds of formula (IV) or (V) are particularly useful in treating cancer when given in combination with other anti-cancer agent.

In another aspect, the invention provides compounds represented by formula (VI):



(VI)

and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof, wherein:

- 5 X_{41} is O, S, or NR_{42} ;
 X_{42} is CR_{44} or N;
 Y_{40} is N or CR_{43} ;
 Y_{41} is N or CR_{45} ;
 Y_{42} , for each occurrence, is independently N, C or CR_{46} ;
- 10 Z is OH, SH, or NHR_7 ;
- R_{41} is -H, -OH, -SH, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, halo, cyano, nitro,
- 15 guanadino, a haloalkyl, a heteroalkyl, an alkoxy or cycloalkoxy, a haloalkoxy, - $NR_{10}R_{11}$, - OR_7 , - $C(O)R_7$, - $C(O)OR_7$, - $C(S)R_7$, - $C(O)SR_7$, - $C(S)SR_7$, - $C(S)OR_7$, - $C(S)NR_{10}R_{11}$, - $C(NR_8)OR_7$, - $C(NR_8)R_7$, - $C(NR_8)NR_{10}R_{11}$, - $C(NR_8)SR_7$, - $OC(O)R_7$, - $OC(O)OR_7$, - $OC(S)OR_7$, - $OC(NR_8)OR_7$, - $SC(O)R_7$, - $SC(O)OR_7$, - $SC(NR_8)OR_7$, - $OC(S)R_7$, - $SC(S)R_7$, - $SC(S)OR_7$, - $OC(O)NR_{10}R_{11}$, - $OC(S)NR_{10}R_{11}$, - $OC(NR_8)NR_{10}R_{11}$, - $SC(O)NR_{10}R_{11}$, - $SC(NR_8)NR_{10}R_{11}$, - $SC(S)NR_{10}R_{11}$, - $OC(NR_8)R_7$, - $SC(NR_8)R_7$, - $C(O)NR_{10}R_{11}$, - $NR_8C(O)R_7$, - $NR_7C(S)R_7$, - $NR_7C(S)OR_7$, - $NR_7C(NR_8)R_7$, - $NR_7C(O)OR_7$, - $NR_7C(NR_8)OR_7$, - $NR_7C(O)NR_{10}R_{11}$, - $NR_7C(S)NR_{10}R_{11}$, - $NR_7C(NR_8)NR_{10}R_{11}$, - SR_7 , - $S(O)_pR_7$, - $OS(O)_pR_7$, - $OS(O)_pOR_7$, - $OS(O)_pNR_{10}R_{11}$, - $S(O)_pOR_7$, - $NR_8S(O)_pR_7$,
- 25

-NR₇S(O)_pNR₁₀R₁₁, -NR₇S(O)_pOR₇, -S(O)_pNR₁₀R₁₁, -SS(O)_pR₇, -SS(O)_pOR₇,
-SS(O)_pNR₁₀R₁₁, -OP(O)(OR₇)₂, or -SP(O)(OR₇)₂;

- R₄₂ is -H, an optionally substituted alkyl, an optionally substituted alkenyl,
an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally
5 substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally
substituted aryl, an optionally substituted heteroaryl, an optionally substituted
aralkyl, an optionally substituted heteraralkyl, hydroxyalkyl, alkoxyalkyl, a
haloalkyl, a heteroalkyl, -C(O)R₇, -(CH₂)_mC(O)OR₇, -C(O)OR₇, -OC(O)R₇,
-C(O)NR₁₀R₁₁, -S(O)_pR₇, -S(O)_pOR₇, or -S(O)_pNR₁₀R₁₁;
- 10 R₄₃ and R₄₄ are, independently, -H, -OH, an optionally substituted alkyl, an
optionally substituted alkenyl, an optionally substituted alkynyl, an optionally
substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally
substituted heterocyclyl, an optionally substituted aryl, an optionally substituted
heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl,
15 hydroxyalkyl, alkoxyalkyl, halo, cyano, nitro, guanadino, a haloalkyl, a heteroalkyl,
-C(O)R₇, -C(O)OR₇, -OC(O)R₇, -C(O)NR₁₀R₁₁, -NR₈C(O)R₇, -SR₇, -S(O)_pR₇,
-OS(O)_pR₇, -S(O)_pOR₇, -NR₈S(O)_pR₇, -S(O)_pNR₁₀R₁₁, or R₄₃ and R₄₄ taken
together with the carbon atoms to which they are attached form an optionally
substituted cycloalkenyl, an optionally substituted aryl, an optionally substituted
20 heterocyclyl, or an optionally substituted heteroaryl;

- R₄₅ is -H, -OH, -SH, -NR₇H, -OR₂₆, -SR₂₆, -NHR₂₆, -O(CH₂)_mOH,
-O(CH₂)_mSH, -O(CH₂)_mNR₇H, -S(CH₂)_mOH, -S(CH₂)_mSH, -S(CH₂)_mNR₇H,
-OC(O)NR₁₀R₁₁, -SC(O)NR₁₀R₁₁, -NR₇C(O)NR₁₀R₁₁, -OC(O)R₇, -SC(O)R₇,
-NR₇C(O)R₇, -OC(O)OR₇, -SC(O)OR₇, -NR₇C(O)OR₇, -OCH₂C(O)R₇,
25 -SCH₂C(O)R₇, -NR₇CH₂C(O)R₇, -OCH₂C(O)OR₇, -SCH₂C(O)OR₇,
-NR₇CH₂C(O)OR₇, -OCH₂C(O)NR₁₀R₁₁, -SCH₂C(O)NR₁₀R₁₁,
-NR₇CH₂C(O)NR₁₀R₁₁, -OS(O)_pR₇, -SS(O)_pR₇, -NR₇S(O)_pR₇, -OS(O)_pNR₁₀R₁₁,
-SS(O)_pNR₁₀R₁₁, -NR₇S(O)_pNR₁₀R₁₁, -OS(O)_pOR₇, -SS(O)_pOR₇, -NR₇S(O)_pOR₇,
-OC(S)R₇, -SC(S)R₇, -NR₇C(S)R₇, -OC(S)OR₇, -SC(S)OR₇, -NR₇C(S)OR₇,
30 -OC(S)NR₁₀R₁₁, -SC(S)NR₁₀R₁₁, -NR₇C(S)NR₁₀R₁₁, -OC(NR₈)R₇, -SC(NR₈)R₇,
-NR₇C(NR₈)R₇, -OC(NR₈)OR₇, -SC(NR₈)OR₇, -NR₇C(NR₈)OR₇,
-OC(NR₈)NR₁₀R₁₁, -SC(NR₈)NR₁₀R₁₁, or -NR₇C(NR₈)NR₁₀R₁₁;

R_{46} , for each occurrence, is independently selected from the group consisting of H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, halo, cyano, nitro, guanadino, a haloalkyl, a heteroalkyl, -NR₁₀R₁₁, -OR₇, -C(O)R₇, -C(O)OR₇, -OC(O)R₇, -C(O)NR₁₀R₁₁, -NR₈C(O)R₇, -SR₇, -S(O)_pR₇, -OS(O)_pR₇, -S(O)_pOR₇, -NR₈S(O)_pR₇, or -S(O)_pNR₁₀R₁₁;

R_7 , R_8 , R_{10} , R_{11} , R_{26} , p , and m are defined as above.

10 In one embodiment, in formula (VI), X_{41} is NR₄₂ and X_{42} is CR₄₄.

In another embodiment, in formula (VI), X_{41} is NR₄₂ and X_{42} is N.

In another embodiment, in formula (VI), R_{41} is selected from the group consisting of -H, lower alkyl, lower alkoxy, lower cycloalkyl, and lower cycloalkoxy.

15 In another embodiment, in formula (VI), R_{41} is selected from the group consisting of -H, methyl, ethyl, propyl, isopropyl, cyclopropyl, methoxy, ethoxy, propoxy, and cyclopropoxy.

In another embodiment, in formula (VI), X_{41} is NR₄₂, and R_{42} is selected from the group consisting of -H, a lower alkyl, a lower cycloalkyl, -C(O)N(R₂₇)₂, and -C(O)OH, wherein R_{27} is -H or a lower alkyl.

20 In another embodiment, in formula (VI), X_{41} is NR₄₂, and R_{42} is selected from the group consisting of -H, methyl, ethyl, n-propyl, isopropyl, cyclopropyl, n-butyl, *sec*-butyl, *tert*-butyl, n-pentyl, n-hexyl, -C(O)OH, -(CH₂)_mC(O)OH, -CH₂OCH₃, -CH₂CH₂OCH₃, and -C(O)N(CH₃)₂.

25 In one embodiment, Y_{40} is CR₄₃. Preferably, Y_{40} is CR₄₃ and R_{43} is H or a lower alkyl.

In another embodiment, in formula (VI), R_{43} and R_{44} are, independently, selected from the group consisting of -H, methyl, ethyl, propyl, isopropyl, cyclopropyl, methoxy, ethoxy, propoxy, and cyclopropoxy.

30 In another embodiment, in formula (VI), X_{42} is CR₄₄; Y is CR₄₃; and R_{43} and R_{44} together with the carbon atoms to which they are attached form a cycloalkenyl, an aryl, heterocyclyl, or heteroaryl ring. In one aspect of this embodiment, R_{43} and

R₄₄ together with the carbon atoms to which they are attached form a C₅-C₈ cycloalkenyl or a C₅-C₈ aryl.

In another embodiment, in formula (VI), R₄₅ is selected from the group consisting of -H, -OH, -SH, -NH₂, a lower alkoxy, a lower alkyl amino, and a lower dialkyl amino.

In another embodiment, in formula (VI), R₄₅ is selected from the group consisting of -H, -OH, methoxy and ethoxy.

In another embodiment, in formula (VI), X₄₁ is O.

In another embodiment, the compound is selected from the group consisting of:

3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(2-methyl-7-methoxy-benzofuran-4-yl)-5-mercapto-[1,2,4]triazole,

3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(benzofuran-5-yl)-5-mercapto-[1,2,4]triazole,

3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(2-methyl-1,3-benzoxaz-5-yl)-5-mercapto-[1,2,4]triazole, and

tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof.

In another embodiment, in formula (VI), Z is -OH.

In another embodiment, the compound is selected from the group consisting of:

3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole,

3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole,

3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole,

3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-isopropyl-indol-4-yl)-5-hydroxy-[1,2,4]triazole, and

tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof.

In another embodiment, Z is -SH.

In another embodiment, the compound is selected from the group consisting of:


3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indazol-5-yl)-5-mercapto-[1,2,4]triazole,

5 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indazol-6-yl)-5-mercapto-[1,2,4]triazole, and

tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof.

Compounds of formula (VI) inhibit the activity of Hsp90 and are particularly useful for treating or preventing proliferative disorders, such as cancer. In addition, compounds of formula (VI) are particularly useful in treating cancer when given in combination with other anti-cancer agent.

In another aspect, the invention provides compounds represented by formula (VII):



and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof, wherein:

20 Z₁ is -OH or -SH;

X₄₂, R₄₁, R₄₂, R₄₃, and R₄₅ are defined as above.

In one embodiment, in formula (VII), Z₁ is -OH.

In another embodiment, in formula (VII), Z₁ is -SH.

In another embodiment, in formula (VII), R_{41} is selected from the group consisting of -H, lower alkyl, lower alkoxy, lower cycloalkyl, and lower cycloalkoxy.

5 In another embodiment, in formula (VII), R_{41} is selected from the group consisting of -H, methyl, ethyl, propyl, isopropyl, cyclopropyl, methoxy, ethoxy, propoxy, and cyclopropoxy.

In another embodiment, in formula (VII), R_{42} is selected from the group consisting of lower alkyl, lower cycloalkyl, $-C(O)N(R_{27})_2$, or $-C(O)OH$, wherein R_{27} is -H or a lower alkyl.

10 In another embodiment, in formula (VII), R_{42} is selected from the group consisting of -H, methyl, ethyl, n-propyl, isopropyl, cyclopropyl, n-butyl, *sec*-butyl, *tert*-butyl, n-pentyl, n-hexyl, $-C(O)OH$, $-(CH_2)_mC(O)OH$, $-CH_2OCH_3$, $-CH_2CH_2OCH_3$, and $-C(O)N(CH_3)_2$.

In another embodiment, R_{43} is H or a lower alkyl.

15 In another embodiment, in formula (VII), X_{42} is CR_{44} , and R_{43} and R_{44} are, independently, selected from the group consisting of -H, methyl, ethyl, propyl, isopropyl, cyclopropyl, methoxy, ethoxy, propoxy, and cyclopropoxy.

20 In another embodiment, in formula (VII), X_{42} is CR_{44} , and R_{43} and R_{44} , taken together with the carbon atoms to which they are attached, form a cycloalkenyl, aryl, heterocyclyl, or heteroaryl ring. Preferably, in this embodiment, R_{43} and R_{44} , taken together with the carbon atoms to which they are attached, form a C_5 - C_8 cycloalkenyl or a C_5 - C_8 aryl.

25 In another embodiment, in formula (VII), R_{45} is selected from the group consisting of -H, -OH, -SH, -NH₂, a lower alkoxy, a lower alkyl amino, and a lower dialkyl amino.

In another embodiment, in formula (VII), R_{45} is selected from the group consisting of -H, -OH, methoxy, and ethoxy.

In another embodiment, in formula (VII), X_{43} is CR_{44} .

30 of:

3-(2,4-dihydroxyphenyl)-4-(1-ethyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,

- 3-(2,4-dihydroxyphenyl)-4-(1-isopropyl-indol-4-yl)-5-mercapto-
[1,2,4]triazole,
3-(2,4-dihydroxyphenyl)-4-(indol-4-yl)-5-mercapto-[1,2,4]triazole,
3-(2,4-dihydroxyphenyl)-4-(1-methoxyethyl-indol-4-yl)-5-mercapto-
5 [1,2,4]triazole,
3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-indol-4-yl)-5-mercapto-
[1,2,4]triazole,
3-(2,4-dihydroxyphenyl)-4-(1-dimethylcarbamoyl-indol-4-yl)-5-mercapto-
[1,2,4]triazole,
10 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-propyl-indol-4-yl)-5-mercapto-
[1,2,4]triazole,
3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,2,3-trimethyl-indol-5-yl)-5-mercapto-
[1,2,4]triazole,
3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(2,3-dimethyl-indol-5-yl)-5-mercapto-
15 [1,2,4]triazole,
3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-acetyl-2,3-dimethyl-indol-5-yl)-5-
mercapto-[1,2,4]triazole,
3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-5-
mercapto-[1,2,4]triazole,
20 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-propyl-2,3-dimethyl-indol-5-yl)-5-
mercapto-[1,2,4]triazole,
3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(N-methyl-tetrahydrocarbozol-7-yl)-5-
mercapto-[1,2,4]triazole,
3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(N-methyl-cyclononan[a]indol-5-yl)-5-
25 mercapto-[1,2,4]triazole,
3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-n-butyl-indol-4-yl)-5-mercapto-
[1,2,4]triazole,
3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-n-pentyl-indol-4-yl)-5-mercapto-
[1,2,4]triazole,
30 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-n-hexyl-indol-4-yl)-5-mercapto-
[1,2,4]triazole,

- 3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1-(1-methylcyclopropyl)-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- 5 3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1,2,3-trimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-5-mercapto-[1,2,4]triazole disodium salt,
- 3-(2,4-dihydroxy-5-*tert*-butyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-
- 10 5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1-propyl-7-methoxy-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-methyl-3-ethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- 15 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-methyl-3-isopropyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- 20 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(N-ethyl-carbozol-7-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-7-hydroxy-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- 25 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-7-ethoxy-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,2-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(N-methyl-indol-5-yl)-5-mercapto-
- 30 [1,2,4]triazole,
- 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,

3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,

3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,

5 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1H-indol-5-yl)-5-mercapto-[1,2,4]triazole,

3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,2-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,

10 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-ethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,

3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-propyl-indol-5-yl)-5-mercapto-[1,2,4]triazole, and

tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof.

15 In another embodiment, in formula (VII), X₄₂ is N.

In another embodiment, the compound is selected from the group consisting of

3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-ethyl-benzimidazol-4-yl)-5-mercapto-[1,2,4]triazole,

20 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-ethyl-benzimidazol-4-yl)-5-mercapto-[1,2,4]triazole HCL salt,

3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(2-methyl-3-ethyl-benzimidazol-5-yl)-5-mercapto-[1,2,4]triazole,

25 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-ethyl-2-methyl-benzimidazol-5-yl)-5-mercapto-[1,2,4]triazole,


3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-2-trifluoromethyl-benzimidazol-5-yl)-5-mercapto-[1,2,4]triazole, and

tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof.

30 Compounds of formula (VII) inhibit the activity of Hsp90 and are particularly useful for treating or preventing proliferative disorders, such as cancer.

In addition, compounds of formula (VII) are particularly useful in treating cancer when given in combination with other anti-cancer agent.

In another aspect, the invention provides compounds represented by formula (VIII):



and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof, wherein:

10 X_{45} is CR_{54} or N;

Z_1 is $-OH$ or $-SH$;

R_{52} is selected from the group consisting of $-H$, methyl, ethyl, n-propyl, isopropyl, n-butyl, n-pentyl, n-hexyl, $-(CH_2)_2OCH_3$, $-CH_2C(O)OH$, and $-C(O)N(CH_3)_2$;

15 R_{53} and R_{54} are each, independently, $-H$, methyl, ethyl, or isopropyl; or R_{53} and R_{54} taken together with the carbon atoms to which they are attached form a phenyl, cyclohexenyl, or cyclooctenyl ring;

R_{55} is selected from the group consisting of $-H$, $-OH$, $-OCH_3$, and $-OCH_2CH_3$; and

20 R_{56} is selected from the group consisting of $-H$, methyl, ethyl, isopropyl, and cyclopropyl.

In one embodiment, in formula (VIII), Z_1 is $-OH$.

In another embodiment, in formula (VIII), Z_1 is $-SH$.

In another embodiment, in formula (VIII), R_{53} is H or a lower alkyl.


In another embodiment, in formula (VIII), X_{45} is CR_{54} . Preferably, R_{54} is H or a lower alkyl.

In another embodiment, X_{45} is N.

5 In another embodiment, the compound is 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(N-methyl-indol-5-yl)-5-mercapto-[1,2,4]triazole.

Compounds of formula (VIII) inhibit the activity of Hsp90 and are particularly useful for treating or preventing proliferative disorders, such as cancer. In addition, compounds of formula (VIII) are particularly useful in treating cancer when given in combination with other anti-cancer agent.

In another aspect, the invention provides compounds represented by formula (IX):



15

and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof, wherein,

X_{44} , for each occurrence, is independently, O, NR_{42} or $\text{C}(\text{R}_{46})_2$;

20 Y_{43} is NR_{42} or $C(R_{46})_2$;

Y_{41} , Y_{42} , Z , R_{41} , R_{42} , and R_{46} are defined as above.

In one embodiment, in formula (IX), R₄₁ is selected from the group consisting of -H, lower alkyl, lower alkoxy, lower cycloalkyl, and lower cycloalkoxy.

In another embodiment, in formula (IX), R_{41} is selected from the group consisting of -H, methyl, ethyl, propyl, isopropyl, cyclopropyl, methoxy, ethoxy, propoxy, and cyclopropoxy.

5 In another embodiment, in formula (IX), R_{42} is selected from the group consisting of -H, methyl, ethyl, n-propyl, isopropyl, cyclopropyl, n-butyl, *sec*-butyl, *tert*-butyl, n-pentyl, n-hexyl, $-C(O)OH$, $-(CH_2)_mC(O)OH$, $-CH_2OCH_3$, $-CH_2CH_2OCH_3$, and $-C(O)N(CH_3)_2$.

In another embodiment, in formula (IX), Y_{41} is CR_{45} . Preferably, R_{45} is H, a lower alkoxy, or -OH.

10 In another embodiment, in formula (IX), Y_{42} is CH.

In another embodiment, in formula (IX), Y_{43} is CH_2 .

In another embodiment, in formula (IX), Y_{43} is NR_{42} , wherein R_{42} is H or a lower alkyl.

15 In another embodiment, in formula (IX), one of X_{44} is NR_{42} and the other is CH_2 or $C(R_6)_2$. Preferably, one of X_{44} is NR_{42} and the other is CH_2 .


In another embodiment, in formula (VI), Z is -OH.

In another embodiment, Z is -SH.

20 Compounds of formula (IX) inhibit the activity of Hsp90 and are particularly useful for treating or preventing proliferative disorders, such as cancer. In addition, compounds of formula (IX) are particularly useful in treating cancer when given in combination with other anti-cancer agent.

In another aspect, the invention provides compounds represented by formula (X):

25



and tautomers, pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof, wherein:

5 X_{41} , Y_{41} , Y_{42} , Z , R_7 , R_8 , R_{10} , R_{11} , R_{41} , R_{46} , and p are defined as above.

In one embodiment, in formula (X), R_{41} is selected from the group consisting of -H, lower alkyl, lower alkoxy, lower cycloalkyl, and lower cycloalkoxy.

In another embodiment, in formula (X), R_{41} is selected from the group consisting of -H, methyl, ethyl, propyl, isopropyl, cyclopropyl, methoxy, ethoxy,
10 propoxy, and cyclopropoxy.

In another embodiment, in formula (X), X_{41} is NR_{42} . Preferably, R_{42} is selected from the group consisting of -H, methyl, ethyl, n-propyl, isopropyl, cyclopropyl, n-butyl, *sec*-butyl, *tert*-butyl, n-pentyl, n-hexyl, -C(O)OH, -(CH₂)_mC(O)OH, -CH₂OCH₃, -CH₂CH₂OCH₃, and -C(O)N(CH₃)₂. More preferably,
15 R_{42} is H or a lower alkyl.

In another embodiment, in formula (X), X_{41} is O.

In another embodiment, in formula (X), X_{41} is S.

In another embodiment, in formula (X), Y_{41} is CR₄₅. Preferably, R_{45} is H, a lower alkoxy, or -OH.

20 In another embodiment, in formula (X), Y_{42} is CH.

In another embodiment, in formula (X), R_{46} is H or a lower alkyl.

In one embodiment, the compound is 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(2-methyl-indazol-6-yl)-5-mercapto-[1,2,4]triazole.







Compounds of formula (X) inhibit the activity of Hsp90 and are particularly
25 useful for treating or preventing proliferative disorders, such as cancer. In addition,















compounds of formula (X) are particularly useful in treating cancer when given in combination with other anti-cancer agent.













i) **Exemplary Compounds of the Invention**











- 5 Exemplary compounds of the invention are depicted in Table 1 below, including tautomers, pharmaceutically acceptable salts, solvates, clathrates, hydrates, polymorphs or prodrugs thereof .













Table 1











No.	Structure	Tautomeric Structure	Name
1			3-(2-Hydroxyphenyl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
2			3-(2,4-Dihydroxyphenyl)-4-[4-(2-methoxyethoxy)-naphthalen-1-yl]-5-mercapto-[1,2,4] triazole
3			3-(2,4-Dihydroxyphenyl)-4-(2-methyl-4-bromophenyl)-5-mercapto-[1,2,4] triazole











4			3-(2,4-Dihydroxyphenyl)-4-(4-bromophenyl)-5-mercapto-[1,2,4] triazole
5			3-(3,4-Dihydroxyphenyl)-4-(6-methoxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
6			3-(3,4-Dihydroxyphenyl)-4-(6-ethoxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
7			3-(3,4-Dihydroxyphenyl)-4-(6-propoxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
8			3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(5-methoxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
9			3-(3,4-Dihydroxyphenyl)-4-(6-isopropoxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
10			3-(2,4-Dihydroxyphenyl)-4-(2,6-diethylphenyl)-5-mercapto-[1,2,4] triazole













11			3-(2,4-Dihydroxyphenyl)- 4-(2-methyl-6-ethylphenyl)-5-mercapto- [1,2,4] triazole
12			3-(2,4-Dihydroxyphenyl)- 4-(2,6-diisopropylphenyl)- 5-mercapto-[1,2,4] triazole
13			3-(2,4-Dihydroxyphenyl)- 4-(1-ethyl-indol-4-yl)-5- mercapto-[1,2,4] triazole
14			3-(2,4-Dihydroxyphenyl)- 4-(2,3-dihydro- benzo[1,4]dioxin-5-yl)-5- mercapto-[1,2,4] triazole
15			3-(2,4-Dihydroxyphenyl)- 4-(3-methylphenyl)-5- mercapto-[1,2,4] triazole
16			3-(2,4-Dihydroxyphenyl)- 4-(4-methylphenyl)-5- mercapto-[1,2,4] triazole

17			3-(2,4-Dihydroxyphenyl)- 4-(2-chlorophenyl)-5- mercapto-[1,2,4] triazole
18			3-(2,4-Dihydroxyphenyl)- 4-(3-chlorophenyl)-5- mercapto-[1,2,4] triazole
19			3-(2,4-Dihydroxyphenyl)- 4-(4-chlorophenyl)-5- mercapto-[1,2,4] triazole
20			3-(2,4-Dihydroxyphenyl)- 4-(2-methoxyphenyl)-5- mercapto-[1,2,4] triazole
21			3-(2,4-Dihydroxyphenyl)- 4-(3-methoxyphenyl)-5- mercapto-[1,2,4] triazole

22			3-(2,4-Dihydroxyphenyl)- 4-(4-methoxyphenyl)-5- mercapto-[1,2,4] triazole
23			3-(2,4-Dihydroxyphenyl)- 4-(3-fluorophenyl)-5- mercapto-[1,2,4] triazole
24			3-(2,4-Dihydroxyphenyl)- 4-(2-ethylphenyl)-5- mercapto-[1,2,4] triazole
25			3-(2-Hydroxy-4- fluorophenyl)-4- (naphthalen-1-yl)-5- mercapto-[1,2,4] triazole
26			3-(2-Hydroxy-4- aminophenyl)-4- (naphthalen-1-yl)-5- mercapto-[1,2,4] triazole
27			3-(2,4-Dihydroxyphenyl)- 4-(2-methyl-4-butyl- phenyl)-5-mercapto-[1,2,4] triazole

28			3-(2,4-Dihydroxyphenyl)- 4-(2,4-dimethyl-phenyl)-5- mercapto-[1,2,4] triazole
29			3-(2,4-Dihydroxyphenyl)- 4-(2,6-dimethyl-phenyl)-5- mercapto-[1,2,4] triazole
30			3-(2,4-Dihydroxyphenyl)- 4-(2,6-dimethyl-phenyl)-5- mercapto-[1,2,4] triazole
31			3-(2,4-Dihydroxyphenyl)- 4-(4-fluorophenyl)-5- mercapto-[1,2,4] triazole
32			3-(2,4-Dihydroxyphenyl)- 4-(2- methylsulfanylphenyl)-5- mercapto-[1,2,4] triazole

33			3-(2,4-Dihydroxyphenyl)- 4-(naphthalene-2-yl)-5- mercapto-[1,2,4] triazole
34			3-(2,4-Dihydroxyphenyl)- 4-(2,3-dimethylphenyl)-5- mercapto-[1,2,4] triazole
35			3-(2,4-Dihydroxyphenyl)- 4-(2-methyl-4- fluorophenyl)-5-mercapto- [1,2,4] triazole
36			3-(2,4-Dihydroxyphenyl)- 4-(acenaphthalen-5-yl)-5- mercapto-[1,2,4] triazole
37			3-(2-Hydroxy-4-methoxy- phenyl)-4-(naphthalen-1- yl)-5-mercapto-[1,2,4] triazole

38			3-(2,4-Dihydroxyphenyl)- 4-(2,3-dichlorophenyl)-5- mercapto-[1,2,4] triazole
39			3-(2,4-Dihydroxyphenyl)- 4-(5-methoxynaphthalen- 1-yl)-5-mercapto-[1,2,4] triazole
40			3-(2,4-Dihydroxyphenyl)- 4-(pyren-1-yl)-5-mercapto- [1,2,4] triazole
41			3-(2,4-Dihydroxyphenyl)- 4-(quinolin-5-yl)-5- mercapto-[1,2,4] triazole
42			3-(2,4-Dihydroxyphenyl)- 4-(1,2,3,4- tetrahydronaphthalen-5- yl)-5-mercapto-[1,2,4] triazole
43			3-(2,4-Dihydroxyphenyl)- 4-(anthracen-1-yl)-5- mercapto-[1,2,4] triazole

44			3-(2,4-Dihydroxyphenyl)- 4-(biphenyl-2-yl)-5- mercapto-[1,2,4] triazole
45			3-(2,4-Dihydroxy-6- methyl-phenyl)-4- (naphthalene-1-yl)-5- mercapto-[1,2,4] triazole
46			3-(2,4-Dihydroxyphenyl)- 4-(4-pentyloxyphenyl)-5- mercapto-[1,2,4] triazole
47			3-(2,4-Dihydroxyphenyl)- 4-(4-octyloxyphenyl)-5- mercapto-[1,2,4] triazole
48			3-(2,4-Dihydroxyphenyl)- 4-(4-chloronaphthalen-1- yl)-5-mercapto-[1,2,4] triazole
49			3-(2,4-Dihydroxy-5-ethyl- phenyl)-4-(naphthalen-1- yl)-5-mercapto-[1,2,4] triazole

50			3-(2,4-Dihydroxy-5-ethylphenyl)-4-(7-carboxymethoxynaphthalen-1-yl)-5-mercapto-[1,2,4] triazole
51			3-(2,4-Dihydroxyphenyl)-4-(2-methyl-quinolin-4-yl)-5-mercapto-[1,2,4] triazole
52			3-(3-Hydroxypyridin-4-yl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
53			3-(2-Hydroxy-4-acetylamino-phenyl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
54			3-(2,4-Dihydroxy-phenyl)-4-(1,2,3,4-tetrahydronaphthalen-1-yl)-5-mercapto-[1,2,4] triazole
55			3-(2,4-Dihydroxy-phenyl)-4-(2,3-dihydrobenzo[1,4]dioxin-5-yl)-5-mercapto-[1,2,4] triazole













56			3-(2,4-Dihydroxy-phenyl)- 4-(3,5-dimethoxyphenyl)- 5-mercapto-[1,2,4] triazole
57			3-(2,4-Dihydroxy-phenyl)- 4-(2,3-dimethyl-1 <i>H</i> -indol- 4-yl)-5-mercapto-[1,2,4] triazole
58			3-(2,4-Dihydroxy-3- propyl-phenyl)-4- (naphthalen-1-yl)-5- mercapto-[1,2,4] triazole
59			3-(1-ethyl-4-hydroxy-6- oxo-1,6-dihydro-pyridin-3- yl)-4-(naphthalen-1-yl)-5- mercapto-[1,2,4] triazole
60			3-(4-hydroxy-6-oxo- pyridin-3-yl)-4- (naphthalen-1-yl)-5- mercapto-[1,2,4] triazole
61			3-(2,4-Dihydroxy-phenyl)- 4-(3,5-di- <i>tert</i> -butylphenyl)- 5-mercapto-[1,2,4] triazole

62			3-(2,6-Dihydroxy-5-fluoro-pyridin-3-yl) 4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
63			3-(2,4-Dihydroxy-5-methyl-phenyl)-4-(naphthalene-1-yl)-5-mercapto-[1,2,4] triazole
64			3-[2,4-Dihydroxy-phenyl]-4-(3-benzoylphenyl)-5-mercapto-[1,2,4] triazole
65			3-(2,4-Dihydroxy-phenyl)-4-(4-carboxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
66			3-(2,4-Dihydroxy-phenyl)-4-[4-(N,N-dimethylcarbamoyl)-naphthalen-1-yl]-5-mercapto-[1,2,4] triazole
67			3-(2,4-Dihydroxy-phenyl)-4-(4-propoxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole

68			3-(2,4-Dihydroxy-phenyl)-4-(4-isopropoxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
69			3-(2,4-Dihydroxy-phenyl)-4-(5-isopropoxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
70			3-(2,4-Dihydroxy-phenyl)-4-(isoquinolin-5-yl)-5-mercapto-[1,2,4] triazole
71			3-(2,4-Dihydroxy-phenyl)-4-(5-propoxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
72			3-(2-Hydroxy-4-methanesulfonamino-phenyl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
73			3-(2,4-Dihydroxy-3,6-dimethyl-phenyl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole













74		3-(2,4-Dihydroxy-phenyl)-4-[7-(2-methoxyethoxy)-naphthalen-1-yl]-5-mercapto-[1,2,4] triazole
75		3-(2,4-Dihydroxy-5-hexyl-phenyl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
76		3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(4-methoxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
77		3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(6-methoxy-naphthalin-1-yl)-5-mercapto-[1,2,4] triazole
78		3-(2,4-Dihydroxy-3-chloro-5-ethyl-phenyl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
79		3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(2,3-dimethoxy-4-methoxy-phenyl)-5-mercapto-[1,2,4] triazole
80		3-(2,4-Dihydroxy-phenyl)-4-(7-isopropoxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole










81			3-(2,4-Dihydroxy-phenyl)-4-(7-ethoxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
82			3-(2,4-Dihydroxy-phenyl)-4-(7-propoxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
83			3-(2-Hydroxy-4-methoxymethoxy-phenyl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
84			3-[2-Hydroxy-4-(2-hydroxy-ethoxy)-phenyl]-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
85			3-(2,4-Dihydroxyphenyl)-4-(7-methoxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
86			3-(2,4-Dihydroxyphenyl)-4-(5-methoxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
87			3-(2,4-Dihydroxyphenyl)-4-(4-hydroxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole









88			3-(2,4-Dihydroxyphenyl)- 4-(1-isopropyl-indol-4-yl)- 5-mercapto-[1,2,4] triazole
89			3-(2,4-Dihydroxy-5-tert- butyl-phenyl)-4- (naphthalen-1-yl)-5- mercapto-[1,2,4] triazole
90			3-(2,4-Dihydroxy-5- propyl-phenyl)-4- (naphthalen-1-yl)-5- mercapto-[1,2,4] triazole
91			3-(2,4-Dihydroxy-3- methyl-5-ethyl-phenyl)-4- (naphthalen-1-yl)-5- mercapto-[1,2,4] triazole
92			3-(2,4-Dihydroxy-5- isobutyl-phenyl)-4- (naphthalen-1-yl)-5- mercapto-[1,2,4] triazole
93			3-(2,4-Dihydroxy-phenyl)- 4-(2,3-dimethoxy-phenyl)- 5-mercapto-[1,2,4] triazole

94			3-(2,4-Dihydroxy-phenyl)-4-(2-methoxy-3-chloro-phenyl)-5-mercapto-[1,2,4] triazole
95			3-(2,4-Dihydroxy-phenyl)-4-(indol-4-yl)-5-mercapto-[1,2,4] triazole
96			3-(2,4-Dihydroxy-phenyl)-4-[1-(2-methoxyethoxy)-indol-4-yl]-5-mercapto-[1,2,4] triazole
97			3-(2,4-Dihydroxy-phenyl)-4-(naphthalen-1-yl)-5-hydroxy-[1,2,4] triazole
98			3-(1-Oxo-3-hydroxy-pyridin-4-yl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
99			3-(2,5-Dihydroxy-4-carboxy)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole

100			3-(2,4-Dihydroxy-5-ethylphenyl)-4-(1-isopropylindol-4-yl)-5-mercapto-[1,2,4] triazole
101			3-(2,4-Dihydroxy-5-ethylphenyl)-4-[1-(dimethylcarbamoyl)indol-4-yl]-5-mercapto-[1,2,4] triazole
102			3-(2,4-Dihydroxy-5-ethylphenyl)-4-(1-ethylbenzoimidazol-4-yl)-5-mercapto-[1,2,4] triazole
103			3-(2,4-Dihydroxy-5-ethylphenyl)-4-(1,2,3-trimethylindol-5-yl)-5-mercapto-[1,2,4] triazole
104			3-(2,5-Dihydroxy-4-hydroxymethylphenyl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
105			3-(2-Hydroxy-4-amino-phenyl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole











106			3-(2-Hydroxy-4-acetylamino-phenyl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
107			3-(2,4-Dihydroxy-3-chloro-phenyl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
108			3-(2,4-Dihydroxy-phenyl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
109			3-(2,4-Dihydroxy-phenyl)-4-(2-methyl-phenyl)-5-mercapto-[1,2,4] triazole
110			3-(2,4-Dihydroxy-phenyl)-4-(2,5-dimethoxy-phenyl)-5-mercapto-[1,2,4] triazole
111			3-(2,4-Dihydroxy-phenyl)-4-phenyl-5-mercapto-[1,2,4] triazole













112			3-(2-Hydroxy-phenyl)-4-(2-methoxy-phenyl)-5-mercapto-[1,2,4] triazole
113			3-(2-Hydroxy-phenyl)-4-(4-methyl-phenyl)-5-mercapto-[1,2,4] triazole
114			3-(2-Hydroxy-phenyl)-4-(4-bromo-phenyl)-5-mercapto-[1,2,4] triazole
115			3-(2,4-Dihydroxy-phenyl)-4-(naphthalen-1-yl)-5-(methyl sulfanyl)- [1,2,4] triazole
116			3-(2,4-Dimethoxy-phenyl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole

117			3-[2,4-Di-(dimethylcarbamoyloxy)-phenyl]-4-(naphthalen-1-yl)-5-(dimethylcarbamoylsulfanyl)- [1,2,4] triazole
118			3-(2,4-Dihydroxy-phenyl)-4-(naphthalen-1-yl)-5-(dimethylcarbamoylsulfanyl)- [1,2,4] triazole
119			3-(2,4-Diethoxycarbonyloxy-phenyl)-4-(naphthalen-1-yl)-5-(ethoxycarbonylsulfanyl)- [1,2,4] triazole
120			3-(2,4-Di-isobutyryloxy-phenyl)-4-(naphthalen-1-yl)-5-(isobutyrylsulfanyl)- [1,2,4] triazole
121			3-[2,4-Di-(dimethylcarbamoyloxy)-phenyl]-4-(quinolin-5-yl)-5-(dimethylcarbamoylsulfanyl)- [1,2,4] triazole
122			3-(2,4-Diacetoxy-phenyl)-4-(naphthalen-1-yl)-5-(acetylsulfanyl)- [1,2,4] triazole
123			3-(2,4-Diacetoxy-phenyl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole

124			3-(2,4-Diethylcarbamoyloxy-phenyl)-4-(naphthalen-1-yl)-5-(ethylcarbamoylsulfanyl)-[1,2,4] triazole
125			3-(2,4-Dihydroxy-phenyl)-4-(naphthalen-1-yl)-5-(2-hydroxyethylsulfanyl)-[1,2,4] triazole
126			3-(2,4-Dihydroxy-phenyl)-4-ethyl-5-mercapto-[1,2,4] triazole
127			3-(2,4-Dihydroxy-phenyl)-4-propyl-5-mercapto-[1,2,4] triazole
128			3-(2,4-Dihydroxy-phenyl)-4-isopropyl-5-mercapto-[1,2,4] triazole
129			3-(2,4-Dihydroxy-phenyl)-4-butyl-5-mercapto-[1,2,4] triazole
130			3-(2,4-Dihydroxy-phenyl)-4-cyclopropyl-5-mercapto-[1,2,4] triazole

131			3-(2,4-Dihydroxy-phenyl)- 4-(naphthalen-1-yl)-5- (carboxyethylsulfanyl)- [1,2,4] triazole
132			3-(2,6-Dimethoxy-5- fluoro-pyridin-3-yl)-4- (naphthalen-1-yl)-5- mercapto-[1,2,4] triazole
133			3-(2-Methanesulfonyloxy- 4-methanesulfonylamino- phenyl)-4-(naphthalen-1- yl)-5-mercapto-[1,2,4] triazole
134			3-(2-Methoxy-phenyl)-4- (4-methoxy-phenyl)-5- mercapto-[1,2,4] triazole
135			3-(3-Hydroxy-naphthalen- 2-yl)-4-phenyl-5- mercapto-[1,2,4] triazole












136			3-(2-Methoxy-phenyl)-4-(4-methyl-phenyl)-5-mercapto-[1,2,4] triazole
137			3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(3-methoxy-phenyl)-5-hydroxy-[1,2,4] triazole
138			3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(naphthalen-1-yl)-5-hydroxy-[1,2,4] triazole
139			3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-indol-3-yl)-5-hydroxy-[1,2,4] triazole
140			3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-indol-4-yl)-5-amino-[1,2,4] triazole

141			3-(2,4-Dihydroxy-5-ethylphenyl)-4-(3-methoxyphenyl)-5-amino-[1,2,4] triazole
142			3-(2,4-Dihydroxy-5-ethylphenyl)-4-(naphthalen-1-yl)-5-amino-[1,2,4] triazole
143			3-(2-Hydroxy-5-ethoxyphenyl)-4-(naphthalen-1-yl)-5-hydroxy-[1,2,4] triazole
144			3-(2-Hydroxy-5-isopropylphenyl)-4-(naphthalen-1-yl)-5-hydroxy-[1,2,4] triazole
145			3-(2-Dihydroxy-phenyl)-4-(7-fluoro-naphthalen-1-yl)-5-hydroxy-[1,2,4] triazole
146			3-(2,4-Dihydroxy-phenyl)-4-(2,3-difluorophenyl)-5-hydroxy-[1,2,4] triazole

147			3-(2,4-Dihydroxy-phenyl)-4-[2-(1 <i>H</i> -tetrazol-5-yl)-phenyl]-5-hydroxy-[1,2,4] triazole
148			3-(2,4-Dihydroxy-phenyl)-4-(benzothiazol-4-yl)-5-hydroxy-[1,2,4] triazole
149			3-(2,4-Dihydroxy-phenyl)-4-(9 <i>H</i> -purin-6-yl)-5-hydroxy-[1,2,4] triazole
150			3-(2,4-Dihydroxy-phenyl)-4-{4-[2-(morpholin-1-yl)-ethoxy]-phenyl}-5-hydroxy-[1,2,4] triazole
151			3-(2,4-Dihydroxy-phenyl)-4-cyclopentyl-5-hydroxy-[1,2,4] triazole
152			3-(2,4-Dihydroxy-phenyl)-4-phenyl-5-(sulfamoylamino)- [1,2,4] triazole

153			3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(naphthalene-1-yl)-5-ureido-[1,2,4] triazole
154			3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(2,3-difluorophenyl)-5-ureido-[1,2,4] triazole
155			3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-indol-4-yl)-5-ureido-[1,2,4] triazole
156			3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(quinolin-5-yl)-5-ureido-[1,2,4] triazole
157			3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(naphthalene-1-yl)-5-carbamoyloxy-[1,2,4] triazole
158			3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(3-trifluoromethyl-phenyl)-5-carbamoyloxy-[1,2,4] triazole
159			3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(1-methyl-indol-4-yl)-5-carbamoyloxy-[1,2,4] triazole
160			3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(8-methoxy-quinolin-5-yl)-5-carbamoyloxy-[1,2,4] triazole

161			3-(2,4-Dihydroxy-5-isopropyl-phenyl)-4-(3-methyl-quinolin-5-yl)-5-carboxyamino-[1,2,4] triazole
162			3-(2,4-Dihydroxy-phenyl)-4-(1-methyl-2-chloro-indol-4-yl)-5-carbamoyloxy-[1,2,4] triazole
163			3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-[3,5-di(trifluoromethyl)-phenyl]-5-carbamoyloxy-[1,2,4] triazole
164			3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(3-trifluoromethyl-phenyl)-5-(sulfamoylamino)-[1,2,4] triazole
165			3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(naphthalene-1-yl)-5-(sulfamoylamino)-[1,2,4] triazole
166			3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(1-isopropyl-benzoimidazol-4-yl)-5-(sulfamoylamino)-[1,2,4] triazole
167			3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(3-isopropylphenyl)-5-(thiocarboxyamino)-[1,2,4] triazole

168			3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(3-isopropoxy-phenyl)-5-(sulfamoyloxy)- [1,2,4] triazole
169			3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(naphthalene-1-yl)-5-(sulfamoyloxy)- [1,2,4] triazole
170			3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(1-isopropyl-benzoimidazol-4-yl)-5-(sulfamoyloxy)- [1,2,4] triazole
171			3-(2-Hydroxy-4-ethoxycarbonyoxy-5-methoxy-phenyl)-4-(1-isopropyl-benzoimidazol-4-yl)-5-hydroxy-[1,2,4] triazole
172			3-(2-Hydroxy-4-ethoxycarbonyoxy-5-ethyl-phenyl)-4-(naphthalin-2-yl)-5-hydroxy-[1,2,4] triazole
173			3-[2-Hydroxy-4-(dimethyl-carbamoyoxy)-5-ethyl-phenyl]-4-(naphthalin-2-yl)-5-hydroxy-[1,2,4] triazole
174			3-[2-Hydroxy-4-(dimethyl-carbamoyoxy)-5-chloro-phenyl]-4-(quinolin-5-yl)-5-mercapto-[1,2,4] triazole

175		3-[2-Hydroxy-4-(dimethyl-carbamoyoxy)-5-ethyl-phenyl]-4-(2,3-difluoro-phenyl)-5-mercapto-[1,2,4] triazole
176		3-[2-Hydroxy-4-isobutyryloxy-5-ethyl-phenyl]-4-(1-methyl-benzo-imidazol-4-yl)-5-hydroxy-[1,2,4] triazole
177		3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
178		3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(5-hydroxy-naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
179		3-(2,4-Dihydroxy-phenyl)-4-(naphthalen-1-ylmethyl)-5-mercapto-[1,2,4] triazole
180		3-(2-Hydroxy-4-methoxyphenyl)-4-(naphthalen-1-yl)-5-mercapto-[1,2,4] triazole
181		3-(2,4-Dihydroxy-phenyl)-4-(biphenyl-3-yl)-5-mercapto-[1,2,4] triazole

182		3-(2,4-Dihydroxy-phenyl)- 4-(2-methyl-5- hydroxymethyl-phenyl)-5- mercapto-[1,2,4] triazole
183		3-(2,4-Dihydroxy-phenyl)- 4-(1-dimethylcarbamoyl- indol-4-yl)-5-mercapto- [1,2,4] triazole
184		3-(2,4,5-Trihydroxy- phenyl)-4-(naphthalene-1- yl)-5-mercapto-[1,2,4] triazole
185		3-(2,4-Dihydroxy-5-ethyl- phenyl)-4-(2,3-dimethyl- indol-5-yl)-5-mercapto- [1,2,4] triazole
186		3-(2,4-Dihydroxy-5-ethyl- phenyl)-4-(3-t-butyl-4- methoxy-phenyl)-5- mercapto-[1,2,4] triazole
187		3-(2,4-Dihydroxy-5-ethyl- phenyl)-4-(1-ethyl-1H- benzoimidazol-4-yl)-5- mercapto-[1,2,4] triazole, HCl salt











188			3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-5-mercapto-[1,2,4] triazole
189			3-(2,4-Dihydroxy-5-cyclopropyl-phenyl)-4-(naphthalene-1-yl)-5-mercapto-[1,2,4] triazole
190			3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-propyl-indol-4-yl)-5-mercapto-[1,2,4] triazole
191			3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-acetyl-2,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4] triazole
192			3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(2-methyl-3-ethyl-benzimidazol-5-yl)-5-mercapto-[1,2,4] triazole
193			3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-ethyl-2-methyl-benzimidazol-5-yl)-5-mercapto-[1,2,4] triazole











194			3-(2,4-dihydroxy-5-ethylphenyl)-4-(1-propyl-2,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4] triazole
195			3-(2,4-dihydroxy-5-ethylphenyl)-4-(N-methyl-tetrahydrocarbozol-7-yl)-5-mercapto-[1,2,4] triazole
196			3-(2,4-dihydroxy-5-ethylphenyl)-4-(N-methyl-cyclononan[a]indol-5-yl)-5-mercapto-[1,2,4] triazole
197			3-(2,4-dihydroxy-5-ethylphenyl)-4-(1-n-butyl-indol-4-yl)-5-mercapto-[1,2,4] triazole
198			3-(2,4-dihydroxy-5-ethylphenyl)-4-(1-n-pentyl-indol-4-yl)-5-mercapto-[1,2,4] triazole
199			3-(2,4-dihydroxy-5-ethylphenyl)-4-(1-n-hexyl-indol-4-yl)-5-mercapto-[1,2,4] triazole

200			3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1-(1-methylcyclopropyl)-indol-4-yl)-5-mercapto-[1,2,4] triazole
201			3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-5-mercapto-[1,2,4] triazole
202			3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1,2,3-trimethyl-indol-5-yl)-5-mercapto-[1,2,4] triazole
203			3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-5-mercapto-[1,2,4] triazole disodium salt
204			3-(2,4-dihydroxy-5-tert-butyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-5-mercapto-[1,2,4] triazole
205			3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1-propyl-7-methoxy-indol-4-yl)-5-mercapto-[1,2,4] triazole

206		3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-methyl-3-ethyl-indol-5-yl)-5-mercapto-[1,2,4] triazole
207		3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4] triazole
208		3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-5-mercapto-[1,2,4] triazole
209		3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-methyl-3-isopropyl-indol-5-yl)-5-mercapto-[1,2,4] triazole
210		3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(N-ethyl-carbozol-7-yl)-5-mercapto-[1,2,4] triazole
211		3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-7-hydroxy-indol-4-yl)-5-mercapto-[1,2,4] triazole

212		3-(2,4-dihydroxy-5-ethylphenyl)-4-(1-isopropyl-7-ethoxy-indol-4-yl)-5-mercapto-[1,2,4] triazole
213		3-(2,4-dihydroxy-5-ethylphenyl)-4-(1,2-dimethyl-indol-5-yl)-5-mercapto-[1,2,4] triazole
214		3-(2,4-dihydroxy-5-ethylphenyl)-4-(N-methyl-indol-5-yl)-5-mercapto-[1,2,4] triazole
215		3-(2,4-dihydroxy-5-ethylphenyl)-4-(2-methyl-7-methoxy-benzofuran-4-yl)-5-mercapto-[1,2,4] triazole
216		3-(2,4-dihydroxy-5-ethylphenyl)-4-(benzofuran-5-yl)-5-mercapto-[1,2,4] triazole
217		3-(2,4-dihydroxy-5-ethylphenyl)-4-(2-methyl-1,3-benzoxaz-5-yl)-5-mercapto-[1,2,4] triazole

218			3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4] triazole
219			3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4] triazole
220			3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-hydroxy-[1,2,4] triazole
221			3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(N-methyl-indol-5-yl)-5-mercapto-[1,2,4] triazole
222			3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1,2-dimethyl-indol-5-yl)-5-mercapto-[1,2,4] triazole

223			3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-hydroxy-[1,2,4] triazole
224			3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-mercapto-[1,2,4] triazole
225			3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1H-indol-5-yl)-5-mercapto-[1,2,4] triazole
226			3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4] triazole
227			3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-ethyl-indol-5-yl)-5-mercapto-[1,2,4] triazole

228			3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-propyl-indol-5-yl)-5-mercapto-[1,2,4] triazole
229			3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-2-trifluoromethyl-benzimidazol-5-yl)-5-mercapto-[1,2,4] triazole
230			3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indazol-5-yl)-5-mercapto-[1,2,4] triazole
231			3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indazol-6-yl)-5-mercapto-[1,2,4] triazole
232			3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-isopropyl-indol-4-yl)-5-hydroxy-[1,2,4] triazole



233			3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1,3-benzodioxol-5-yl)-5-mercapto-[1,2,4] triazole
234			3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(indan-5-yl)-5-mercapto-[1,2,4] triazole
235			3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(2-methyl-indazol-6-yl)-5-mercapto-[1,2,4] triazole
236			3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(3-oxo-benzo[1,4]oxazin-6-yl)-5-mercapto-[1,2,4] triazole
237			3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(2-oxo-1,3-dihydro-benzoimidazol-5-yl)-5-mercapto-[1,2,4] triazole
238			3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(2H-benzo[1,4]oxazin-6-yl)-5-mercapto-[1,2,4] triazole

239			4-Ethyl-6-[5-mercapto-4-(1-methyl-2,3-dihydro-1H-indol-5-yl)-4H-[1,2,4]triazol-3-yl]-benzene-1,3-diol
240			5-(3-(5-ethyl-2,4-dihydroxyphenyl)-5-mercapto-4H-1,2,4-triazol-4-yl)indolin-2-one
241			5-(3-(5-ethyl-2,4-dihydroxyphenyl)-5-mercapto-4H-1,2,4-triazol-4-yl)-1H-benzo[d]imidazol-2(3H)-one
242			5-(3-(5-ethyl-2,4-dihydroxyphenyl)-5-mercapto-4H-1,2,4-triazol-4-yl)-1-methylindolin-2-one
243			4-isopropyl-6-(5-mercapto-4-(4-propyl-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl)-4H-1,2,4-triazol-3-yl)benzene-1,3-diol

244		6-(3-(5-ethyl-2,4-dihydroxyphenyl)-5-mercapto-4H-1,2,4-triazol-4-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one
245		6-(3-(5-ethyl-2,4-dihydroxyphenyl)-5-mercapto-4H-1,2,4-triazol-4-yl)-3-methylbenzo[d]thiazol-2(3H)-one
246		6-(3-(5-ethyl-2,4-dihydroxyphenyl)-5-mercapto-4H-1,2,4-triazol-4-yl)benzo[d]thiazol-2(3H)-one





Preferred compounds of the invention are those compounds that can form a tautomeric structure as shown below and as exemplified by the tautomeric structures shown in Table 1:

5



Similarly, prodrugs, *i.e.* compounds which can be metabolized or hydrolyzed *in vivo* to a compound of the present invention are encompassed by the present

description. For example, the following embodiments of a compound of the present invention can be produced *in vivo* in the following reaction:



where R_{200} is R_2 , R_5 or R_{18} .

One skilled in the art will understand that other hydrolyzable protecting groups can be employed with the compounds of the present invention to obtain prodrugs encompassed by the present description.

15 Without wishing to be bound by any theory, it is believed that the compounds of the invention preferentially bind to Hsp90 in the tautomeric form shown above, and thereby inhibit the activity of Hsp90.

C. Methods for Making Compounds of the Invention

Compounds of the invention can be obtained via standard, well-known synthetic methodology, see *e.g.*, March, J. Advanced Organic Chemistry; Reactions Mechanisms, and Structure, 4th ed., 1992. In particular, compounds of the invention

5 can be obtained by heating a hydrazide (A) with an isocyanate ($X_{14} = O$), isothiocyanate, ($X_{14} = S$) or carbodiimide ($X_{14} = NR_7$) (B) in an alcohol to form intermediate (C). Intermediate (C) can be cyclized to form a triazole core (D) by heating it in an aqueous solution which includes about 2 molar equivalents of NaOH (see Scheme I below). Starting materials useful for preparing compounds of the

10 invention and intermediates therefore, are commercially available or can be prepared from commercially available materials using known synthetic methods and reagents. For example, a hydrazide can be prepared by reacting an ester (such as 2,4-dihydroxybenzoic acid methyl ester) or acid chloride with hydrazine. Isocyanates and isothiocyanates (X_{14} is O or S, respectively) can be formed in a number of ways

15 from compounds that have a primary amine group. For example, a primary amine can be reacted with phosgene or thiophosgene to form an isocyanate or an isothiocyanate, respectively. Alternatively, a cyanate or thiocyanate ion can be reacted with an alkyl halide to form an alkyl isocyanate or an alkyl isothiocyanate. In addition, a isothiocyanate can be prepared by reacting a diazonium salt with a

20 thiocyanate ion. Carbodiimides (X_{14} is NR_7) can be prepared by dehydration of ureas using a dehydration agent such as tosyl chloride in pyridine, $POCl_3$, PCl_5 , P_2O_5 -pyridine, and $Ph_3PBr_2-Et_3N$. Other methods of preparing isocyanates, thioisocyanates, and carbodiimides can be found in March, J. Advanced Organic Chemistry; Reactions Mechanisms, and Structure, 4th ed., 1992, the entire teachings

25 of which are incorporated by reference.


Compounds represented by formulas (IV) and (V) can be made in an analogous fashion as compounds depicted in Scheme I.

Reactive functional groups can be protected during one or more reaction step, then deprotected to restore the original functionality. Examples of suitable

30 protecting groups for hydroxyl groups include benzyl, methoxymethyl, allyl, trimethylsilyl, tert-butyldimethylsilyl, acetate, and the like. Examples of suitable amine protecting groups include benzyloxycarbonyl, tert-butoxycarbonyl, tert-butyl,

- benzyl and fluorenylmethyloxy-carbonyl (Fmoc). Examples of suitable thiol protecting groups include benzyl, tert-butyl, acetyl, methoxymethyl and the like. Other suitable protecting groups are well known to those of ordinary skill in the art and include those found in T. W. Greene, *Protecting Groups in Organic Synthesis*, 5 John Wiley & Sons, Inc. 1981.


Scheme I: Synthesis of triazole compounds of the invention



- 10 An alternative method of preparing the compounds of the invention is shown in Scheme II. In this method, an aryl, heteroaryl, cycloalkyl, or alkyl amine compound (i) is stirred at about room temperature with a thiocarbonyl (ii) which has two leaving groups, L₁ and L₂, such as imidazole-1-yl groups, to form compound (iii). Typically, the thiocarbonyl compound is present in a slight molar excess of about 1.05 eq. to about 1.3 eq. compared with compound (i). Compound (iii) is then combined with a hydrazide compound (iv) in a solvent and heated to about 50°C to about 100°C for about 0.5 to 5 hrs to form compound (v). Typically, compound (iii) and compound (iv) can be present in about equal molar ratio or a slight excess of compound (iii), such as about 1.01 to about 1.1 molar eq. of compound (iii) compare to compound (iv). Compound (v) can then be cyclized to form a triazole compound of the invention (vi) by suspending it in aqueous solution containing about 2 molar eq. of NaOH and heating the solution to about 75°C to about 110°C for about 0.5 hr

to about 2 hrs. Typically, the NaOH solution containing compound (v) is degassed before heating by bubbling an inert gas, such as nitrogen or argon, through it.

Scheme II: Alternative synthesis of triazole compounds of the invention



In one embodiment, ring A of the compounds of the invention is a 2,4-dihydroxyphenyl group. In this embodiment, it is sometimes desirable to prepare a prodrug by protecting the 4-hydroxy group with a moiety that can be hydrolyzed in vivo. Protection of the 4-hydroxy group is expected to improve the circulating half-life of compound compounds of the invention. In addition, it is desirable that a group added to the 4-hydroxy group increase the water solubility of the compounds of the invention. In one embodiment, 4-methyl-piperazine-1-carbamoyl group is used to protect the 4-hydroxy group (see Scheme III). In this embodiment, a compound of the invention, such as compound (E), is treated with about one molar equivalents of 4-methyl-piperazine-1-carbonyl chloride (F) in the presence of a base to form compound (G) in which the 4-hydroxy group is protected. Alternatively, the metcapto group can be protected first by reacting compound (E) with about one molar equivalent of acyl chloride in the presence of a base to form intermediate (H). Intermediate (H) can then be reacted with about one molar equivalent of 4-methyl-


10

15

20


piperazine-1-carbonyl chloride (**F**) in the presence of a base, then the acetyl group can be removed by treatment with a mild acid to form compound (**G**).

5 Scheme III: Preparation of prodrugs in which the 4-hydroxy group of compounds of the invention is protected with 4-methyl-piperazine-1-carbamoyl.



Another prodrug of compounds of the the invention can be formed by
 10 addition of a phosphate group to the 4-hydroxy group (Scheme IV). In this embodiment, a compound of the invention, such as compound (**E**), is treated with about one molar equivalent of diisopropyl phosphoramidous acid di-t-butyl ester in the presence of tetrazole to yield compound (**J**). The phosphorous group is then
 15 oxidized with m-CPBA to form a phosphoric acid di-t-butyl ester group of compound **K**. The t-butyl groups are then hydrolyzed with trifluoroacetic acid (TFA) to yield a phosphoric acid group or compound **L**.

Scheme IV: Preparation of prodrugs in which the 4-hydroxy group of compounds of the invention is protected with a phosphate group.



D. Uses of Compounds of the Invention

5 The present invention is directed to therapies which involve administering one or more compounds of the invention, or compositions comprising said compounds to a subject, preferably a human subject, to inhibit the activity of Hsp90 or to prevent, treat, manage, or ameliorate a proliferative disorder, such as cancer, or one or more symptoms thereof. In one embodiment, the present invention is

10 directed to treating cancers in which aberrant expression and/or activation of c-kit has been implicated as contributing to neoplastic pathology by administering one or more compounds of the invention.

In one aspect, the invention provides a method of inhibiting the activity of Hsp90 in a cell, comprising administering to the cell an effective amount of a

15 compound represented by formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X), or any embodiment thereof, or a compound shown in Table 1. In one embodiment, the compound is administered to a cell in a subject, preferably a mammal, and more preferably a human.

In another aspect, the invention provides a method of treating or preventing a

20 proliferation disorder in a mammal, comprising administering to the mammal an effective amount of a compound represented by formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X), or any embodiment thereof, or a compound shown in Table 1. In one embodiment, the compound is administered to a human to treat or

prevent a proliferative disorder. In another embodiment, the proliferation disorder is cancer. In another embodiment, the compound is administered with one or more additional therapeutic agents. In a preferred embodiment, the additional therapeutic agent is an anticancer agent.

5 In another aspect, the invention provides a method for treating cancer in a mammal, comprising administering to the mammal an effective amount of a compound represented by formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X), or any embodiment thereof, or a compound shown in Table 1. In one embodiment, the compound is administered to a human to treat or prevent cancer.

10 In another embodiment, the compound is administered with one or more additional therapeutic agents. In a preferred embodiment, the one or more additional therapeutic agents are anticancer agents.

In another aspect, the invention provides a method for treating a c-kit associated cancer in a mammal, comprising administering to the mammal an effective amount of a compound represented by formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X), or any embodiment thereof, or a compound shown in Table 1. In one embodiment, the compound is administered to a human to treat or prevent the c-kit associated cancer. In another embodiment, the compound is administered with one or more additional therapeutic agents. In a preferred embodiment, the one or more additional therapeutic agents are anticancer agents.

1. c-Kit Associated Cancers

SCF binding to the c-kit protects hematopoietic stem and progenitor cells from apoptosis (Lee, *et al.*, 1997, *J. Immunol.*, 159:3211-3219), thereby contributing to colony formation and hematopoiesis. Expression of c-kit is frequently observed in acute myelocytic leukemia (AML) and sometimes observed in acute lymphocytic leukemia (ALL) (for reviews, see Sperling, *et al.*, 1997, *Haemat.*, 82:617-621; Escribano, *et al.*, 1998, *Leuk. Lymph.*, 30:459-466). Although c-kit is expressed in the majority of AML cells, its expression does not appear to be prognostic of disease progression (Sperling, *et al.*, 1997, *Haemat.* 82:617-621). However, SCF protected AML cells from apoptosis induced by chemotherapeutic agents (Hassan, *et al.*, 1996, *Acta. Hem.*, 95:257-262). Therefore, degradation of c-kit caused by the

inhibition of Hsp90 by the compounds of the invention will enhance the efficacy of these agents and may induce apoptosis of AML cells.

The clonal growth of cells from patients with myelodysplastic syndrome (Sawada, *et al.*, 1996, *Blood*, 88:319-327) or chronic myelogenous leukemia (CML) (Sawai, *et al.*, 1996, *Exp. Hem.*, 2:116-122) was found to be significantly enhanced by SCF in combination with other cytokines. CML is characterized by expansion of Philadelphia chromosome positive cells of the marrow (Verfaillie, *et al.*, 1998, *Leuk.*, 12:136-138), which appears to primarily result from inhibition of apoptotic death (Jones, 1997, *Curr. Opin. Onc.*, 9:3-7). The product of the Philadelphia chromosome, p210.sup.BCR-ABL, has been reported to mediate inhibition of apoptosis (Bedi, *et al.*, 1995, *Blood*, 86:1148-1158). Since p210.sup.BCR-ABL and the c-kit RTK both inhibit apoptosis and p62.sup.dok has been suggested as a substrate (Carpino, *et al.*, 1997, *Cell*, 88:197-204), it is possible that clonal expansion mediated by these kinases occurs through a common signaling pathway. However, c-kit has also been reported to interact directly with p210.sup.BCR-ABL (Hallek, *et al.*, 1996, *Brit. J Haem.*, 94:5-16), which suggests that c-kit may have a more causative role in CML pathology. Therefore, degradation of c-kit caused by the inhibition of Hsp90 by the compounds of the invention will prove useful in the treatment of CML.

Normal colorectal mucosa does not express c-kit (Bellone, *et al.*, 1997, *J. Cell Physiol.*, 172:1-11). However, c-kit is frequently expressed in colorectal carcinoma (Bellone, *et al.*, 1997, *J. Cell Physiol.*, 172: 1-11), and autocrine loops of SCF and c-kit have been observed in several colon carcinoma cell lines (Toyota, *et al.*, 1993, *Turn. Biol.*, 14:295-302; Lahm, *et al.*, 1995, *Cell Growth & Differ.*, 6:1111-1118; Bellone, *et al.*, 1997, *J. Cell Physiol.*, 172:1-11). Furthermore, disruption of the autocrine loop by the use of neutralizing antibodies (Lahm, *et al.*, 1995, *Cell Growth & Differ.*, 6:1111-1118) and downregulation of c-kit and/or SCF significantly inhibits cell proliferation (Lahm, *et al.*, 1995, *Cell Growth & Differ.*, 6:1111-1118; Bellone, *et al.*, 1997, *J. Cell Physiol.*, 172:1-11).

SCF/c-kit autocrine loops have been observed in gastric carcinoma cell lines (Turner, *et al.*, 1992, *Blood*, 80:374-381; Hassan, *et al.*, 1998, *Digest. Dis. Science*, 43:8-14), and constitutive c-kit activation also appears to be important for

gastrointestinal stromal tumors (GISTs). GISTs are the most common mesenchymal tumor of the digestive system. More than 90% of GISTs express c-kit, which is consistent with the putative origin of these tumor cells from interstitial cells of Cajal (ICCs) (Hirota, *et al.*, 1998, *Science*, 279:577-580). The c-kit expressed in GISTs from several different patients was observed to have mutations in the intracellular juxtamembrane domain leading to constitutive activation (Hirota, *et al.*, 1998, *Science* 279:577-580). Therefore, degradation of c-kit caused by the inhibition of Hsp90 by the compounds of the invention will be an efficacious means for the treatment of these cancers.

Male germ cell tumors have been histologically categorized into seminomas, which retain germ cell characteristics, and nonseminomas which can display characteristics of embryonal differentiation. Both seminomas and nonseminomas are thought to initiate from a preinvasive stage designated carcinoma in situ (CIS) (Murty, *et al.*, 1998, *Sem. Oncol.*, 25:133-144). Both c-kit and SCF have been reported to be essential for normal gonadal development during embryogenesis (Loveland, *et al.*, 1997, *J. Endocrinol.*, 153:337-344). Loss of either the receptor or the ligand resulted in animals devoid of germ cells. In postnatal testes, c-kit has been found to be expressed in Leydig cells and spermatogonia, while SCF was expressed in Sertoli cells (Loveland, *et al.*, 1997, *J. Endocrinol.*, 153:337-344). Testicular tumors develop from Leydig cells with high frequency in transgenic mice expressing human papilloma virus 16 (HPV16) E6 and E7 oncogenes (Kondoh, *et al.*, 1991, *J. Virol.*, 65:3335-3339; Kondoh, *et al.*, 1994, *J. Urol.*, 152:2151-2154). These tumors express both c-kit and SCF, and an autocrine loop may contribute to the tumorigenesis (Kondoh, *et al.*, 1995, *Oncogene*, 10:341-347) associated with cellular loss of functional p53 and the retinoblastoma gene product by association with E6 and E7 (Dyson, *et al.*, 1989, *Science*, 243:934-937; Werness, *et al.*, 1990, *Science*, 248:76-79; Scheffner, *et al.*, 1990, *Cell*, 63:1129-1136). Defective signaling mutants of SCF (Kondoh, *et al.*, 1995, *Oncogene*, 10:341-347) or c-kit (Li, *et al.*, 1996, *Canc. Res.*, 56:4343-4346) inhibited formation of testicular tumors in mice expressing HPV16 E6 and E7. Since c-kit kinase activation is pivotal to tumorigenesis in these animals, the compounds of the invention which inhibit Hsp90

and thereby cause the degradation of c-kit will be useful for preventing or treating testicular tumors associated with human papilloma virus.

Expression of c-kit on germ cell tumors shows that the receptor is expressed by the majority of carcinomas in situ and seminomas, but c-kit is expressed in only a minority of nonseminomas (Strohmeyer, *et al.*, 1991, *Canc. Res.*, 51:1811-1816; Rajpert-de Meyts, *et al.*, 1994, *Int. J. Androl.*, 17:85-92; Izquierdo, *et al.*, 1995, *J. Pathol.*, 177:253-258; Strohmeyer, *et al.*, 1995, *J. Urol.*, 153:511-515; Bokenmeyer, *et al.*, 1996, *J. Cance. Res., Clin. Oncol.*, 122:301-306; Sandlow, *et al.*, 1996, *J. Androl.*, 17:403-408). Therefore, degradation of c-kit caused by the inhibition of Hsp90 by the compounds of the invention will be an efficacious means for the treatment of these cancers.

SCF and c-kit are expressed throughout the central nervous system of developing rodents, and the pattern of expression suggests a role in growth, migration and differentiation of neuroectodermal cells. Expression of SCF and c-kit have also been reported in the adult brain (Hamel, *et al.*, 1997, *J. Neuro-Onc.*, 35:327-333). Expression of c-kit has also been observed in normal human brain tissue (Tada, *et al.* 1994, *J. Neuro.*, 80:1063-1073). Glioblastoma and astrocytoma, which define the majority of intracranial tumors, arise from neoplastic transformation of astrocytes (Levin, *et al.*, 1997, *Principles & Practice of Oncology*, 2022-2082). Expression of c-kit has been observed in glioblastoma cell lines and tissues (Berdel, *et al.*, 1992, *Canc. Res.*, 52:3498-3502; Tada, *et al.*, 1994, *J. Neuro.*, 80:1063-1073; Stanulla, *et al.*, 1995, *Act. Neuropath.*, 89:158-165).

The association of c-kit with astrocytoma pathology is less clear. Reports of expression of c-kit in normal astrocytes have been made (Natali, *et al.*, 1992, *Int. J. Canc.*, 52:197-201), (Tada, *et al.* 1994, *J. Neuro.*, 80:1063-1073), while others report it is not expressed (Kristt, *et al.*, 1993, *Neuro.*, 33:106-115). In the former case, high levels of c-kit expression in high grade tumors were observed (Kristt, *et al.*, 1993, *Neuro.*, 33:106-115), whereas in the latter case researchers were unable to detect any expression in astrocytomas. In addition, contradictory reports of c-kit and SCF expression in neuroblastomas also exist. One study found that neuroblastoma cell lines often express SCF, but rarely express c-kit. In primary tumors, c-kit was detected in about 8% of neuroblastomas, while SCF was found in 18% of tumors

(Beck, et al., 1995, *Blood*, 86:3132-3138). In contrast, other studies (Cohen, et al., 1994, *Blood*, 84:3465-3472) have reported that all 14 neuroblastoma cell lines examined contained c-kit/SCF autocrine loops, and expression of both the receptor and ligand were observed in 45% of tumor samples examined. In two cell lines, 5 anti-c-kit antibodies inhibited cell proliferation, suggesting that the SCF/c-kit autocrine loop contributed to growth (Cohen, et al., 1994, *Blood*, 84:3465-3472). Therefore, degradation of c-kit caused by the inhibition of Hsp90 by the compounds of the invention will be an efficacious means for treating some cancers of the central nervous system.

10

2. Combination Therapies and Treatment of Refractory Cancers

The invention also provides methods of preventing, treating, managing, or ameliorating a proliferative disorder, such as cancer, or one or more symptoms thereof, said methods comprising administering to a subject in need thereof one or 15 more compounds of the invention and one or more other therapies (e.g., one or more prophylactic or therapeutic agents that are currently being used, have been used, are known to be useful or in development for use in the prevention, treatment or amelioration of a proliferative disorder, such as cancer, or one or more symptoms associated with said proliferative disorder).

20 The prophylactic or therapeutic agents of the combination therapies of the invention can be administered sequentially or concurrently. In a specific embodiment, the combination therapies of the invention comprise one or more compounds and at least one other therapy (e.g., another prophylactic or therapeutic agent) which has the same mechanism of action as said compounds. In another 25 specific embodiment, the combination therapies of the invention comprise one or more compounds of the invention and at least one other therapy (e.g., another prophylactic or therapeutic agent) which has a different mechanism of action than said compounds. In certain embodiments, the combination therapies of the present invention improve the prophylactic or therapeutic effect of one or more compounds 30 of the invention by functioning together with the compounds to have an additive or synergistic effect. In certain embodiments, the combination therapies of the present invention reduce the side effects associated with the therapies (e.g., prophylactic or

therapeutic agents). In certain embodiments, the combination therapies of the present invention reduce the effective dosage of one or more of the therapies.

The prophylactic or therapeutic agents of the combination therapies can be administered to a subject, preferably a human subject, in the same pharmaceutical composition. In alternative embodiments, the prophylactic or therapeutic agents of the combination therapies can be administered concurrently to a subject in separate pharmaceutical compositions. The prophylactic or therapeutic agents may be administered to a subject by the same or different routes of administration.

In a specific embodiment, a pharmaceutical composition comprising one or more compounds of the invention is administered to a subject, preferably a human, to prevent, treat, manage, or ameliorate a proliferative disorder, such as cancer, or one or more symptom thereof. In accordance with the invention, pharmaceutical compositions of the invention may also comprise one or more other agents (*e.g.*, prophylactic or therapeutic agents which are currently being used, have been used, or are known to be useful in the prevention, treatment or amelioration of a proliferative disorder or a symptom thereof).

The invention provides methods for preventing, managing, treating or ameliorating a proliferative disorder, such as cancer, or one or more symptoms thereof in a subject refractory (either completely or partially) to existing agent therapies for such a proliferative disorder, said methods comprising administering to said subject a dose of an effective amount of one or more compounds of the invention and a dose of an effective amount of one or more therapies (*e.g.*, one or more prophylactic or therapeutic agents useful for the prevention, treatment, management, or amelioration of a proliferative disorder or a symptom thereof). The invention also provides methods for preventing, treating, managing, or ameliorating a proliferative disorder or a symptom thereof by administering one or more compounds of the invention in combination with any other therapy(ies) to patients who have proven refractory to other therapies but are no longer on these therapies.

The compounds of the invention and/or other therapies can be administered to a subject by any route known to one of skill in the art. Examples of routes of administration include, but are not limited to, parenteral, *e.g.*, intravenous,

intradermal, subcutaneous, oral (e.g., inhalation), intranasal, transdermal (topical), transmucosal, and rectal administration.

3. Agents Useful In Combination With the Compounds of the Invention

5 Without wishing to be bound by theory, it is believed that the compounds of the invention can be particularly effective at treating subjects whose cancer has become multi-drug resistant. Although chemotherapeutic agents initially cause tumor regression, most agents that are currently used to treat cancer target only one pathway to tumor progression. Therefore, in many instances, after treatment with
10 one or more chemotherapeutic agents, a tumor develops multidrug resistance and no longer response positively to treatment. One of the advantages of inhibiting Hsp90 activity is that several of its client proteins, which are mostly protein kinases or transcription factors involved in signal transduction, have been shown to be involved in the progression of cancer. Thus, inhibition of Hsp90 provides a method of short
15 circuiting several pathways for tumor progression simultaneously. Therefore, it is believed that treatment of cancer with an Hsp90 inhibitor of the invention either alone, or in combination with other chemotherapeutic agents, is more likely to result in regression or elimination of the tumor, and less likely to result in the development of more aggressive multidrug resistant tumors than other currently available
20 therapies.

Anticancer agents that can be co-administered with the compounds of the invention include TaxolTM, also referred to as "paclitaxel", is a well-known anti-cancer drug which acts by enhancing and stabilizing microtubule formation, and analogs of TaxolTM, such as TaxotereTM. Compounds that have the basic taxane
25 skeleton as a common structure feature, have also been shown to have the ability to arrest cells in the G2-M phases due to stabilization or inhibition of microtubules.

Other anti-cancer agents that can be employed in combination with the compounds of the invention include Avastin, Adriamycin, Dactinomycin, Bleomycin, Vinblastine, Cisplatin, acivicin; aclarubicin; acodazole hydrochloride;
30 acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; aminoglutethimide; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene

- hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; bropirimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefingol; chlorambucil; cirolemycin; cladribine; crisnatol mesylate; cyclophosphamide;
- 5 cytarabine; dacarbazine; daunorubicin hydrochloride; decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziquone; doxorubicin; doxorubicin hydrochloride; droloxifene; droloxifene citrate; dromostanolone propionate; duazomycin; edatrexate; eflornithine hydrochloride; elsamitrucin; enloplatin; enpromate; epipropidine; epirubicin hydrochloride; erbulozole; esorubicin
- 10 hydrochloride; estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; fluocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; ilmofoesine; interleukin II (including recombinant
- 15 interleukin II, or rIL2), interferon alfa-2a; interferon alfa-2b; interferon alfa-n1 ; interferon alfa-n3; interferon beta-I a; interferon gamma-I b; iproplatin; irinotecan hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride; lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprocil; maytansine; mechlorethamine hydrochloride; megestrol acetate;
- 20 melengestrol acetate; melphalan; menogaril; mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedopa; mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane; mitoxantrone hydrochloride; mycophenolic acid; nocodazole; nogalamycin; ormaplatin; oxisuran; pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide;
- 25 pipobroman; piposulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium; porfiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride; pyrazofurin; riboprine; rogletimide; safingol; safingol hydrochloride; semustine; simtrazene; sparfosate sodium; sparsomycin; spirogermanium hydrochloride; spiromustine; spiroplatin; streptonigrin;
- 30 streptozocin; sulofenur; talisomycin; tecogalan sodium; tegafur; teloxantrone hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprine; thioguanine; thiotepa; tiazofurin; tirapazamine; toremifene citrate; trestolone acetate;

tricitabine phosphate; trimetrexate; trimetrexate glucuronate; triptorelin; tubulozole hydrochloride; uracil mustard; uredepa; vapreotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine sulfate; vinepidine sulfate; vinglycinate sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; zorubicin hydrochloride.

Other anti-cancer drugs that can be employed in combination with the compounds of the invention include: 20-epi-1,25 dihydroxyvitamin D₃; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; anti-dorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen; antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin III derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstauroporine; beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; broprimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives; canarypox IL-2; capecitabine; carboxamide-amino-triazole; carboxyamidotriazole; CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B; cetorelix; chlorlins; chloroquinoxaline sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin analogue; conagenin; crambescidin 816; crisnatol; cryptophycin 8; cryptophycin A derivatives; curacin A; cyclopentantraquinones; cycloplatam; cypemycin; cytarabine ocfosfate; cytolytic factor; cytostatin; dacliximab; decitabine; dehydrodidemnin B; deslorelin; dexamethasone; dexifosfamide; dexrazoxane; dexverapamil; diaziquone; didemnin B; didox; diethylnorspermine; dihydro-5-azacytidine; 9- dioxamycin; diphenyl spiromustine; docosanol; dolasetron; doxifluridine; droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine;

- edelfosine; edrecolomab; eflornithine; elemene; emitefur; epirubicin; epristeride; estramustine analogue; estrogen agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane; fadrozole; fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; flezelastine; fluasterone; fludarabine; fluorodaunorubicin
- 5 hydrochloride; forfenimex; formestane; fostriecin; fotemustine; gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; heregulin; hexamethylene bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene; idramantone; ilmofofosine; ilomastat; imidazoacridones; imiquimod; immunostimulant peptides;
- 10 insulin-like growth factor-1 receptor inhibitor; interferon agonists; interferons; interleukins; iobenguane; iododoxorubicin; ipomeanol, 4-; iroplact; irsogladine; isobengazole; isohomohalicondrin B; itasetron; jasplakinolide; kahalalide F; lamellarin-N triacetate; lanreotide; leinamycin; lenograstim; lentinan sulfate; leptolstatin; letrozole; leukemia inhibiting factor; leukocyte alpha interferon;
- 15 leuprolide+estrogen+progesterone; leuprorelin; levamisole; liarozole; linear polyamine analogue; lipophilic disaccharide peptide; lipophilic platinum compounds; lissoclinamide 7; lobaplatin; lombricine; lometrexol; lonidamine; losoxantrone; lovastatin; loxoribine; lurtotecan; lutetium texaphyrin; lysofylline; lytic peptides; maitansine; mannostatin A; marimastat; masoprocol; maspin;
- 20 matrilysin inhibitors; matrix metalloproteinase inhibitors; menogaril; merbarone; meterelin; methioninase; metoclopramide; MIF inhibitor; mifepristone; miltefosine; mirimostim; mismatched double stranded RNA; mitoguazone; mitolactol; mitomycin analogues; mitonafide; mitotoxin fibroblast growth factor-saporin; mitoxantrone; mofarotene; molgramostim; monoclonal antibody, human chorionic gonadotrophin; monophosphoryl lipid A+myobacterium cell wall sk; mopidamol;
- 25 multiple drug resistance gene inhibitor; multiple tumor suppressor 1-based therapy; mustard anticancer agent; mycaperoxide B; mycobacterial cell wall extract; myriaporone; N-acetyldinaline; N-substituted benzamides; nafarelin; nagrestip; naloxone+pentazocine; napavin; naphterpin; nartograstim; nedaplatin; nemorubicin;
- 30 neridronic acid; neutral endopeptidase; nilutamide; nisamycin; nitric oxide modulators; nitroxide antioxidant; nitrullyn; O6-benzylguanine; octreotide; okicenone; oligonucleotides; onapristone; ondansetron; ondansetron; oracin; oral

- cytokine inducer; ormaplatin; osaterone; oxaliplatin; oxaunomycin; palauamine; palmitoylrhizoxin; pamidronic acid; panaxytriol; panomifene; parabactin; pazelliptine; pegaspargase; peldesine; pentosan polysulfate sodium; pentostatin; pentozole; perflubron; perfosfamide; perillyl alcohol; phenazinomycin;
- 5 phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride; pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum complex; platinum compounds; platinum-triamine complex; porfimer sodium; porfiromycin; prednisone; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based immune modulator; protein kinase C
- 10 inhibitor; protein kinase C inhibitors, microalgal; protein tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists; raltitrexed; ramosetron; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes;
- 15 RII retinamide; rogletimide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safingol; saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; signal transduction modulators; single chain antigen-binding protein; sizofiran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol;
- 20 somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin; spongistatin 1; squalamine; stem cell inhibitor; stem-cell division inhibitors; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista; suramin; swainsonine; synthetic glycosaminoglycans; tallimustine; tamoxifen methiodide; tauromustine; tazarotene;
- 25 tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors; temoporfin; temozolomide; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor agonist; thymotrinan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine; titanocene bichloride; topsentin; toremifene; totipotent stem cell
- 30 factor; translation inhibitors; tretinoin; triacetyluridine; triciribine; trimetrexate; triptorelin; tropisetron; turosteride; tyrosine kinase inhibitors; tyrphostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth inhibitory factor; urokinase

receptor antagonists; vapreotide; variolin B; vector system, erythrocyte gene therapy; velaresol; veramine; verdins; verteporfin; vinorelbine; vinxaltine; vitaxin; vorozole; zanoterone; zeniplatin; zilascorb; and zinostatin stimalamer. Preferred anti-cancer drugs are 5-fluorouracil and leucovorin.

- 5 Other chemotherapeutic agents that can be employed in combination with the compounds of the invention include but are not limited to alkylating agents, antimetabolites, natural products, or hormones. Examples of alkylating agents useful for the treatment or prevention of T-cell malignancies in the methods and compositions of the invention include but are not limited to, nitrogen mustards (*e.g.*,
10 mechchloroethamine, cyclophosphamide, chlorambucil, *etc.*), alkyl sulfonates (*e.g.*, busulfan), nitrosoureas (*e.g.*, carmustine, lomusitne, *etc.*), or triazenes (decabazine, *etc.*). Examples of antimetabolites useful for the treatment or prevention of T-cell malignancies in the methods and compositions of the invention include but are not limited to folic acid analog (*e.g.*, methotrexate), or pyrimidine analogs (*e.g.*,
15 Cytarabine), purine analogs (*e.g.*, mercaptopurine, thioguanine, pentostatin). Examples of natural products useful for the treatment or prevention of T-cell malignancies in the methods and compositions of the invention include but are not limited to vinca alkaloids (*e.g.*, vinblastin, vincristine), epipodophyllotoxins (*e.g.*, etoposide), antibiotics (*e.g.*, daunorubicin, doxorubicin, bleomycin), enzymes (*e.g.*,
20 L-asparaginase), or biological response modifiers (*e.g.*, interferon alpha).

- Examples of alkylating agents that can be employed in combination with the compounds of the invention include but are not limited to, nitrogen mustards (*e.g.*, mechchloroethamine, cyclophosphamide, chlorambucil, melphalan, *etc.*), ethylenimine and methylmelamines (*e.g.*, hexamethylmelamine, thiotepa), alkyl sulfonates (*e.g.*,
25 busulfan), nitrosoureas (*e.g.*, carmustine, lomusitne, semustine, streptozocin, *etc.*), or triazenes (decabazine, *etc.*). Examples of antimetabolites useful for the treatment or prevention of cancer in the methods and compositions of the invention include but are not limited to folic acid analog (*e.g.*, methotrexate), or pyrimidine analogs (*e.g.*, fluorouracil, floxouridine, Cytarabine), purine analogs (*e.g.*, mercaptopurine,
30 thioguanine, pentostatin). Examples of natural products useful for the treatment or prevention of cancer in the methods and compositions of the invention include but are not limited to vinca alkaloids (*e.g.*, vinblastin, vincristine), epipodophyllotoxins

(*e.g.*, etoposide, teniposide), antibiotics (*e.g.*, actinomycin D, daunorubicin, doxorubicin, bleomycin, plicamycin, mitomycin), enzymes (*e.g.*, L-asparaginase), or biological response modifiers (*e.g.*, interferon alpha). Examples of hormones and antagonists useful for the treatment or prevention of cancer in the methods and

5 compositions of the invention include but are not limited to adrenocorticosteroids (*e.g.*, prednisone), progestins (*e.g.*, hydroxyprogesterone caproate, megestrol acetate, medroxyprogesterone acetate), estrogens (*e.g.*, diethylstilbestrol, ethinyl estradiol), antiestrogen (*e.g.*, tamoxifen), androgens (*e.g.*, testosterone propionate, fluoxymesterone), antiandrogen (*e.g.*, flutamide), gonadotropin releasing hormone

10 analog (*e.g.*, leuprolide). Other agents that can be used in the methods and compositions of the invention for the treatment or prevention of cancer include platinum coordination complexes (*e.g.*, cisplatin, carboplatin), anthracenedione (*e.g.*, mitoxantrone), substituted urea (*e.g.*, hydroxyurea), methyl hydrazine derivative (*e.g.*, procarbazine), adrenocortical suppressant (*e.g.*, mitotane, aminoglutethimide).

15 Examples of anti-cancer agents which act by arresting cells in the G2-M phases due to stabilization or inhibition of microtubules and which can be used in combination with the compounds of the invention include without limitation the following marketed drugs and drugs in development: Erbulozole (also known as R-55104), Dolastatin 10 (also known as DLS-10 and NSC-376128), Mivobulin

20 isethionate (also known as CI-980), Vincristine, NSC-639829, Discodermolide (also known as NVP-XX-A-296), ABT-751 (Abbott, also known as E-7010), Altorhyrtins (such as Altorhyrtin A and Altorhyrtin C), Spongistatins (such as Spongistatin 1, Spongistatin 2, Spongistatin 3, Spongistatin 4, Spongistatin 5, Spongistatin 6, Spongistatin 7, Spongistatin 8, and Spongistatin 9), Cemadotin hydrochloride (also

25 known as LU-103793 and NSC-D-669356), Epothilones (such as Epothilone A, Epothilone B, Epothilone C (also known as desoxyepothilone A or dEpoA), Epothilone D (also referred to as KOS-862, dEpoB, and desoxyepothilone B), Epothilone E, Epothilone F, Epothilone B N-oxide, Epothilone A N-oxide, 16-aza-epothilone B, 21-aminoepothilone B (also known as BMS-310705), 21-

30 hydroxyepothilone D (also known as Desoxyepothilone F and dEpoF), 26-fluoroepothilone), Auristatin PE (also known as NSC-654663), Soblidotin (also known as TZT-1027), LS-4559-P (Pharmacia, also known as LS-4577), LS-4578

- (Pharmacia, also known as LS-477-P), LS-4477 (Pharmacia), LS-4559 (Pharmacia), RPR-112378 (Aventis), Vincristine sulfate, DZ-3358 (Daiichi), FR-182877 (Fujisawa, also known as WS-9885B), GS-164 (Takeda), GS-198 (Takeda), KAR-2 (Hungarian Academy of Sciences), BSF-223651 (BASF, also known as ILX-651 and LU-223651), SAH-49960 (Lilly/Novartis), SDZ-268970 (Lilly/Novartis), AM-97 (Armad/Kyowa Hakko), AM-132 (Armad), AM-138 (Armad/Kyowa Hakko), IDN-5005 (Indena), Cryptophycin 52 (also known as LY-355703), AC-7739 (Ajinomoto, also known as AVE-8063A and CS-39.HCl), AC-7700 (Ajinomoto, also known as AVE-8062, AVE-8062A, CS-39-L-Ser.HCl, and RPR-258062A), Vitilevuamide, Tubulysin A, Canadensol, Centaureidin (also known as NSC-106969), T-138067 (Tularik, also known as T-67, TL-138067 and TI-138067), COBRA-1 (Parker Hughes Institute, also known as DDE-261 and WHI-261), H10 (Kansas State University), H16 (Kansas State University), Oncocidin A1 (also known as BTO-956 and DIME), DDE-313 (Parker Hughes Institute), Fijianolide B, Laulimalide, SPA-2 (Parker Hughes Institute), SPA-1 (Parker Hughes Institute, also known as SPIKET-P), 3-IAABU (Cytoskeleton/Mt. Sinai School of Medicine, also known as MF-569), Narcosine (also known as NSC-5366), Nascapine, D-24851 (Asta Medica), A-105972 (Abbott), Hemiasterlin, 3-BAABU (Cytoskeleton/Mt. Sinai School of Medicine, also known as MF-191), TMPN (Arizona State University), Vanadocene acetylacetonate, T-138026 (Tularik), Monsatrol, Inanocine (also known as NSC-698666), 3-IAABE (Cytoskeleton/Mt. Sinai School of Medicine), A-204197 (Abbott), T-607 (Tularik, also known as T-900607), RPR-115781 (Aventis), Eleutherobins (such as Desmethyleleutherobin, Desaeyleleutherobin, Isoeleutherobin A, and Z-Eleutherobin), Caribaeoside, Caribaeolin, Halichondrin B, D-64131 (Asta Medica), D-68144 (Asta Medica), Diazonamide A, A-293620 (Abbott), NPI-2350 (Nereus), Taccalonolide A, TUB-245 (Aventis), A-259754 (Abbott), Diozostatin, (-)-Phenylahistin (also known as NSCL-96F037), D-68838 (Asta Medica), D-68836 (Asta Medica), Myoseverin B, D-43411 (Zentaris, also known as D-81862), A-289099 (Abbott), A-318315 (Abbott), HTI-286 (also known as SPA-110, trifluoroacetate salt) (Wyeth), D-82317 (Zentaris), D-82318 (Zentaris), SC-12983 (NCI), Resverastatin phosphate sodium, BPR-0Y-007 (National Health Research Institutes), and SSR-250411 (Sanofi).

E. Compositions and Methods for Administering Therapies

The present invention provides compositions for the treatment, prophylaxis, and amelioration of proliferative disorders, such as cancer. In a specific
5 embodiment, a composition comprises one or more compounds of the invention, or a pharmaceutically acceptable salt, solvate, clathrate, hydrate or prodrug thereof. In another embodiment, a composition of the invention comprises one or more prophylactic or therapeutic agents other than a compound of the invention, or a pharmaceutically acceptable salt, solvate, clathrate, hydrate, prodrug thereof. In
10 another embodiment, a composition of the invention comprises one or more compounds of the invention, or a pharmaceutically acceptable salt, solvate, clathrate, hydrate or prodrug thereof, and one or more other prophylactic or therapeutic agents. In another embodiment, the composition comprises a compound of the invention, or a pharmaceutically acceptable salt, solvate, clathrate, hydrate, or prodrug thereof,
15 and a pharmaceutically acceptable carrier, diluent or excipient.

In a preferred embodiment, a composition of the invention is a pharmaceutical composition or a single unit dosage form. Pharmaceutical compositions and dosage forms of the invention comprise one or more active ingredients in relative amounts and formulated in such a way that a given
20 pharmaceutical composition or dosage form can be used to treat or prevent proliferative disorders, such as cancer. Preferred pharmaceutical compositions and dosage forms comprise a compound of formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X), or Table 1, or a pharmaceutically acceptable prodrug, salt, solvate, clathrate, hydrate, or prodrug thereof, optionally in combination with one or more
25 additional active agents.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include, but are not limited to, parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), intranasal, transdermal (topical),
30 transmucosal, and rectal administration. In a specific embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous, subcutaneous, intramuscular, oral, intranasal

or topical administration to human beings. In a preferred embodiment, a pharmaceutical composition is formulated in accordance with routine procedures for subcutaneous administration to human beings.

Single unit dosage forms of the invention are suitable for oral, mucosal (*e.g.*,
5 nasal, sublingual, vaginal, buccal, or rectal), parenteral (*e.g.*, subcutaneous, intravenous, bolus injection, intramuscular, or intraarterial), or transdermal administration to a patient. Examples of dosage forms include, but are not limited to: tablets; caplets; capsules, such as soft elastic gelatin capsules; cachets; troches; lozenges; dispersions; suppositories; ointments; cataplasms (poultices); pastes;
10 powders; dressings; creams; plasters; solutions; patches; aerosols (*e.g.*, nasal sprays or inhalers); gels; liquid dosage forms suitable for oral or mucosal administration to a patient, including suspensions (*e.g.*, aqueous or non-aqueous liquid suspensions, oil-in-water emulsions, or a water-in-oil liquid emulsions), solutions, and elixirs; liquid dosage forms suitable for parenteral administration to a patient; and sterile
15 solids (*e.g.*, crystalline or amorphous solids) that can be reconstituted to provide liquid dosage forms suitable for parenteral administration to a patient.

The composition, shape, and type of dosage forms of the invention will typically vary depending on their use. For example, a dosage form suitable for mucosal administration may contain a smaller amount of active ingredient(s) than an
20 oral dosage form used to treat the same indication. This aspect of the invention will be readily apparent to those skilled in the art. *See, e.g.*, Remington's Pharmaceutical Sciences (1990) 18th ed., Mack Publishing, Easton PA.

Typical pharmaceutical compositions and dosage forms comprise one or more excipients. Suitable excipients are well known to those skilled in the art of
25 pharmacy, and non-limiting examples of suitable excipients are provided herein. Whether a particular excipient is suitable for incorporation into a pharmaceutical composition or dosage form depends on a variety of factors well known in the art including, but not limited to, the way in which the dosage form will be administered to a patient. For example, oral dosage forms such as tablets may contain excipients
30 not suited for use in parenteral dosage forms.

The suitability of a particular excipient may also depend on the specific active ingredients in the dosage form. For example, the decomposition of some

active ingredients can be accelerated by some excipients such as lactose, or when exposed to water. Active ingredients that comprise primary or secondary amines (*e.g.*, N-desmethylvenlafaxine and N,N-didesmethylvenlafaxine) are particularly susceptible to such accelerated decomposition. Consequently, this invention

5 encompasses pharmaceutical compositions and dosage forms that contain little, if any, lactose. As used herein, the term "lactose-free" means that the amount of lactose present, if any, is insufficient to substantially increase the degradation rate of an active ingredient. Lactose-free compositions of the invention can comprise excipients that are well known in the art and are listed, for example, in the U.S.

10 Pharmacopia (USP) SP (XXI)/NF (XVI). In general, lactose-free compositions comprise active ingredients, a binder/filler, and a lubricant in pharmaceutically compatible and pharmaceutically acceptable amounts. Preferred lactose-free dosage forms comprise active ingredients, microcrystalline cellulose, pre-gelatinized starch, and magnesium stearate.

15 This invention further encompasses anhydrous pharmaceutical compositions and dosage forms comprising active ingredients, since water can facilitate the degradation of some compounds. For example, the addition of water (*e.g.*, 5%) is widely accepted in the pharmaceutical arts as a means of simulating long-term storage in order to determine characteristics such as shelf-life or the stability of

20 formulations over time. *See, e.g.*, Jens T. Carstensen (1995) Drug Stability: Principles & Practice, 2d. Ed., Marcel Dekker, NY, NY, 379-80. In effect, water and heat accelerate the decomposition of some compounds. Thus, the effect of water on a formulation can be of great significance since moisture and/or humidity are commonly encountered during manufacture, handling, packaging, storage,

25 shipment, and use of formulations.

Anhydrous pharmaceutical compositions and dosage forms of the invention can be prepared using anhydrous or low moisture containing ingredients and low moisture or low humidity conditions. Pharmaceutical compositions and dosage forms that comprise lactose and at least one active ingredient that comprises a

30 primary or secondary amine are preferably anhydrous if substantial contact with moisture and/or humidity during manufacturing, packaging, and/or storage is expected.

An anhydrous pharmaceutical composition should be prepared and stored such that its anhydrous nature is maintained. Accordingly, anhydrous compositions are preferably packaged using materials known to prevent exposure to water such that they can be included in suitable formulary kits. Examples of suitable packaging include, but are not limited to, hermetically sealed foils, plastics, unit dose containers (e.g., vials), blister packs, and strip packs.

The invention further encompasses pharmaceutical compositions and dosage forms that comprise one or more compounds that reduce the rate by which an active ingredient will decompose. Such compounds, which are referred to herein as "stabilizer" include, but are not limited to, antioxidants such as ascorbic acid, pH buffers, or salt buffers.

1) Oral Dosage Forms

Pharmaceutical compositions of the invention that are suitable for oral administration can be presented as discrete dosage forms, such as, but are not limited to, tablets (e.g., chewable tablets), caplets, capsules, and liquids (e.g., flavored syrups). Such dosage forms contain predetermined amounts of active ingredients, and may be prepared by methods of pharmacy well known to those skilled in the art. *See generally*, Remington's Pharmaceutical Sciences (1990) 18th ed., Mack Publishing, Easton PA.

Typical oral dosage forms of the invention are prepared by combining the active ingredient(s) in an admixture with at least one excipient according to conventional pharmaceutical compounding techniques. Excipients can take a wide variety of forms depending on the form of preparation desired for administration. For example, excipients suitable for use in oral liquid or aerosol dosage forms include, but are not limited to, water, glycols, oils, alcohols, flavoring agents, preservatives, and coloring agents. Examples of excipients suitable for use in solid oral dosage forms (e.g., powders, tablets, capsules, and caplets) include, but are not limited to, starches, sugars, micro-crystalline cellulose, diluents, granulating agents, lubricants, binders, and disintegrating agents.

Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid excipients are employed. If desired, tablets can be coated by standard aqueous or nonaqueous

techniques. Such dosage forms can be prepared by any of the methods of pharmacy. In general, pharmaceutical compositions and dosage forms are prepared by uniformly and intimately admixing the active ingredients with liquid carriers, finely divided solid carriers, or both, and then shaping the product into the desired presentation if necessary.

For example, a tablet can be prepared by compression or molding. Compressed tablets can be prepared by compressing in a suitable machine the active ingredients in a free-flowing form such as powder or granules, optionally mixed with an excipient. Molded tablets can be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

Examples of excipients that can be used in oral dosage forms of the invention include, but are not limited to, binders, fillers, disintegrants, and lubricants. Binders suitable for use in pharmaceutical compositions and dosage forms include, but are not limited to, corn starch, potato starch, or other starches, gelatin, natural and synthetic gums such as acacia, sodium alginate, alginic acid, other alginates, powdered tragacanth, guar gum, cellulose and its derivatives (*e.g.*, ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose), polyvinyl pyrrolidone, methyl cellulose, pre-gelatinized starch, hydroxypropyl methyl cellulose, (*e.g.*, Nos. 2208, 2906, 2910), microcrystalline cellulose, and mixtures thereof.

Suitable forms of microcrystalline cellulose include, but are not limited to, the materials sold as AVICEL-PH-101, AVICEL-PH-103 AVICEL RC-581, AVICEL-PH-105 (available from FMC Corporation, American Viscose Division, Avicel Sales, Marcus Hook, PA), and mixtures thereof. One specific binder is a mixture of microcrystalline cellulose and sodium carboxymethyl cellulose sold as AVICEL RC-581. Suitable anhydrous or low moisture excipients or additives include AVICEL-PH-103J and Starch 1500 LM.

Examples of fillers suitable for use in the pharmaceutical compositions and dosage forms disclosed herein include, but are not limited to, talc, calcium carbonate (*e.g.*, granules or powder), microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, pre-gelatinized starch, and mixtures thereof. The binder or filler in pharmaceutical compositions of the invention is

typically present in from about 50 to about 99 weight percent of the pharmaceutical composition or dosage form.

Disintegrants are used in the compositions of the invention to provide tablets that disintegrate when exposed to an aqueous environment. Tablets that contain too much disintegrant may disintegrate in storage, while those that contain too little may not disintegrate at a desired rate or under the desired conditions. Thus, a sufficient amount of disintegrant that is neither too much nor too little to detrimentally alter the release of the active ingredients should be used to form solid oral dosage forms of the invention. The amount of disintegrant used varies based upon the type of formulation, and is readily discernible to those of ordinary skill in the art. Typical pharmaceutical compositions comprise from about 0.5 to about 15 weight percent of disintegrant, preferably from about 1 to about 5 weight percent of disintegrant.

Disintegrants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, agar-agar, alginic acid, calcium carbonate, microcrystalline cellulose, croscarmellose sodium, crospovidone, polacrillin potassium, sodium starch glycolate, potato or tapioca starch, other starches, pre-gelatinized starch, other starches, clays, other alginates, other celluloses, gums, and mixtures thereof.

Lubricants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, light mineral oil, glycerin, sorbitol, mannitol, polyethylene glycol, other glycols, stearic acid, sodium lauryl sulfate, talc, hydrogenated vegetable oil (e.g., peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil), zinc stearate, ethyl oleate, ethyl laureate, agar, and mixtures thereof. Additional lubricants include, for example, a syloid silica gel (AEROSIL 200, manufactured by W.R. Grace Co. of Baltimore, MD), a coagulated aerosol of synthetic silica (marketed by Degussa Co. of Plano, TX), CAB-O-SIL (a pyrogenic silicon dioxide product sold by Cabot Co. of Boston, MA), and mixtures thereof. If used at all, lubricants are typically used in an amount of less than about 1 weight percent of the pharmaceutical compositions or dosage forms into which they are incorporated.

2) Controlled Release Dosage Forms

Active ingredients of the invention can be administered by controlled release means or by delivery devices that are well known to those of ordinary skill in the art. Examples include, but are not limited to, those described in U.S. Patent Nos.: 3,845,770; 3,916,899; 3,536,809; 3,598,123; and 4,008,719, 5,674,533, 5,059,595, 5,591,767, 5,120,548, 5,073,543, 5,639,476, 5,354,556, and 5,733,566, each of which is incorporated herein by reference. Such dosage forms can be used to provide slow or controlled-release of one or more active ingredients using, for example, hydropropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic systems, multilayer coatings, microparticles, liposomes, microspheres, or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled-release formulations known to those of ordinary skill in the art, including those described herein, can be readily selected for use with the active ingredients of the invention. The invention thus encompasses single unit dosage forms suitable for oral administration such as, but not limited to, tablets, capsules, gelcaps, and caplets that are adapted for controlled-release.

All controlled-release pharmaceutical products have a common goal of improving drug therapy over that achieved by their non-controlled counterparts. Ideally, the use of an optimally designed controlled-release preparation in medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations include extended activity of the drug, reduced dosage frequency, and increased patient compliance.

Most controlled-release formulations are designed to initially release an amount of drug (active ingredient) that promptly produces the desired therapeutic effect, and gradually and continually release of other amounts of drug to maintain this level of therapeutic or prophylactic effect over an extended period of time. In order to maintain this constant level of drug in the body, the drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body. Controlled-release of an active ingredient can be stimulated by various conditions including, but not limited to, pH, temperature, enzymes, water, or other physiological conditions or compounds.

A particular extended release formulation of this invention comprises a therapeutically or prophylactically effective amount of a compound of formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X), or Table 1, or a pharmaceutically acceptable salt, solvate, hydrate, clathrate, or prodrug thereof, in spheroids which further comprise microcrystalline cellulose and, optionally, hydroxypropylmethylcellulose coated with a mixture of ethyl cellulose and hydroxypropylmethylcellulose. Such extended release formulations can be prepared according to U.S. Patent No. 6,274,171, the entirety of which is incorporated herein by reference.

A specific controlled-release formulation of this invention comprises from about 6% to about 40% a compound of formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), (IX), (X), or Table 1, or a pharmaceutically acceptable salt, solvate, hydrate, clathrate, or prodrug thereof, by weight, about 50% to about 94% microcrystalline cellulose, NF, by weight, and optionally from about 0.25% to about 1% by weight of hydroxypropyl-methylcellulose, USP, wherein the spheroids are coated with a film coating composition comprised of ethyl cellulose and hydroxypropylmethylcellulose.

3) Parenteral Dosage Forms

Parenteral dosage forms can be administered to patients by various routes including, but not limited to, subcutaneous, intravenous (including bolus injection), intramuscular, and intraarterial. Because their administration typically bypasses patients' natural defenses against contaminants, parenteral dosage forms are preferably sterile or capable of being sterilized prior to administration to a patient. Examples of parenteral dosage forms include, but are not limited to, solutions ready for injection, dry products ready to be dissolved or suspended in a pharmaceutically acceptable vehicle for injection, suspensions ready for injection, and emulsions.

Suitable vehicles that can be used to provide parenteral dosage forms of the invention are well known to those skilled in the art. Examples include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles

such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

- Compounds that increase the solubility of one or more of the active
- 5 ingredients disclosed herein can also be incorporated into the parenteral dosage forms of the invention.

4) Transdermal, Topical, and Mucosal Dosage Forms

- Transdermal, topical, and mucosal dosage forms of the invention include, but are not limited to, ophthalmic solutions, sprays, aerosols, creams, lotions, ointments,
- 10 gels, solutions, emulsions, suspensions, or other forms known to one of skill in the art. *See, e.g.,* Remington's Pharmaceutical Sciences (1980 & 1990) 16th and 18th eds., Mack Publishing, Easton PA and Introduction to Pharmaceutical Dosage Forms (1985) 4th ed., Lea & Febiger, Philadelphia. Dosage forms suitable for treating mucosal tissues within the oral cavity can be formulated as mouthwashes or
- 15 as oral gels. Further, transdermal dosage forms include "reservoir type" or "matrix type" patches, which can be applied to the skin and worn for a specific period of time to permit the penetration of a desired amount of active ingredients.

- Suitable excipients (*e.g.*, carriers and diluents) and other materials that can be used to provide transdermal, topical, and mucosal dosage forms encompassed by
- 20 this invention are well known to those skilled in the pharmaceutical arts, and depend on the particular tissue to which a given pharmaceutical composition or dosage form will be applied. With that fact in mind, typical excipients include, but are not limited to, water, acetone, ethanol, ethylene glycol, propylene glycol, butane-1,3-diol, isopropyl myristate, isopropyl palmitate, mineral oil, and mixtures thereof to
- 25 form lotions, tinctures, creams, emulsions, gels or ointments, which are non-toxic and pharmaceutically acceptable. Moisturizers or humectants can also be added to pharmaceutical compositions and dosage forms if desired. Examples of such additional ingredients are well known in the art. *See, e.g.,* Remington's Pharmaceutical Sciences (1980 & 1990) 16th and 18th eds., Mack Publishing,
- 30 Easton PA.

Depending on the specific tissue to be treated, additional components may be used prior to, in conjunction with, or subsequent to treatment with active ingredients of the invention. For example, penetration enhancers can be used to assist in delivering the active ingredients to the tissue. Suitable penetration enhancers
5 include, but are not limited to: acetone; various alcohols such as ethanol, oleyl, and tetrahydrofuryl; alkyl sulfoxides such as dimethyl sulfoxide; dimethyl acetamide; dimethyl formamide; polyethylene glycol; pyrrolidones such as polyvinylpyrrolidone; Kollidon grades (Povidone, Polyvidone); urea; and various
10 water-soluble or insoluble sugar esters such as Tween 80 (polysorbate 80) and Span 60 (sorbitan monostearate).

The pH of a pharmaceutical composition or dosage form, or of the tissue to which the pharmaceutical composition or dosage form is applied, may also be adjusted to improve delivery of one or more active ingredients. Similarly, the polarity of a solvent carrier, its ionic strength, or tonicity can be adjusted to improve
15 delivery. Compounds such as stearates can also be added to pharmaceutical compositions or dosage forms to advantageously alter the hydrophilicity or lipophilicity of one or more active ingredients so as to improve delivery. In this regard, stearates can serve as a lipid vehicle for the formulation, as an emulsifying agent or surfactant, and as a delivery-enhancing or penetration-enhancing agent.
20 Different salts, hydrates or solvates of the active ingredients can be used to further adjust the properties of the resulting composition.

5) Dosage & Frequency of Administration

The amount of the compound or composition of the invention which will be effective in the prevention, treatment, management, or amelioration of a proliferative
25 disorders, such as cancer, or one or more symptoms thereof, will vary with the nature and severity of the disease or condition, and the route by which the active ingredient is administered. The frequency and dosage will also vary according to factors specific for each patient depending on the specific therapy (*e.g.*, therapeutic or prophylactic agents) administered, the severity of the disorder, disease, or
30 condition, the route of administration, as well as age, body, weight, response, and the past medical history of the patient. Effective doses may be extrapolated from

dose-response curves derived from *in vitro* or animal model test systems. Suitable regimens can be selected by one skilled in the art by considering such factors and by following, for example, dosages reported in the literature and recommended in the *Physician's Desk Reference* (57th ed., 2003).

5 Exemplary doses of a small molecule include milligram or microgram amounts of the small molecule per kilogram of subject or sample weight (*e.g.*, about 1 microgram per kilogram to about 500 milligrams per kilogram, about 100 micrograms per kilogram to about 5 milligrams per kilogram, or about 1 microgram per kilogram to about 50 micrograms per kilogram).

10 In general, the recommended daily dose range of a compound of the invention for the conditions described herein lie within the range of from about 0.01 mg to about 1000 mg per day, given as a single once-a-day dose preferably as divided doses throughout a day. In one embodiment, the daily dose is administered twice daily in equally divided doses. Specifically, a daily dose range should be from
15 about 5 mg to about 500 mg per day, more specifically, between about 10 mg and about 200 mg per day. In managing the patient, the therapy should be initiated at a lower dose, perhaps about 1 mg to about 25 mg, and increased if necessary up to about 200 mg to about 1000 mg per day as either a single dose or divided doses, depending on the patient's global response. It may be necessary to use dosages of
20 the active ingredient outside the ranges disclosed herein in some cases, as will be apparent to those of ordinary skill in the art. Furthermore, it is noted that the clinician or treating physician will know how and when to interrupt, adjust, or terminate therapy in conjunction with individual patient response.

Different therapeutically effective amounts may be applicable for different
25 proliferative disorders, as will be readily known by those of ordinary skill in the art. Similarly, amounts sufficient to prevent, manage, treat or ameliorate such proliferative disorders, but insufficient to cause, or sufficient to reduce, adverse effects associated with the compounds of the invention are also encompassed by the above described dosage amounts and dose frequency schedules. Further, when a
30 patient is administered multiple dosages of a compound of the invention, not all of the dosages need be the same. For example, the dosage administered to the patient may be increased to improve the prophylactic or therapeutic effect of the compound

or it may be decreased to reduce one or more side effects that a particular patient is experiencing.

In a specific embodiment, the dosage of the composition of the invention or a compound of the invention administered to prevent, treat, manage, or ameliorate a proliferative disorders, such as cancer, or one or more symptoms thereof in a patient is 150 µg/kg, preferably 250 µg/kg, 500 µg/kg, 1 mg/kg, 5 mg/kg, 10 mg/kg, 25 mg/kg, 50 mg/kg, 75 mg/kg, 100 mg/kg, 125 mg/kg, 150 mg/kg, or 200 mg/kg or more of a patient's body weight. In another embodiment, the dosage of the composition of the invention or a compound of the invention administered to prevent, treat, manage, or ameliorate a proliferative disorders, such as cancer, or one or more symptoms thereof in a patient is a unit dose of 0.1 mg to 20 mg, 0.1 mg to 15 mg, 0.1 mg to 12 mg, 0.1 mg to 10 mg, 0.1 mg to 8 mg, 0.1 mg to 7 mg, 0.1 mg to 5 mg, 0.1 to 2.5 mg, 0.25 mg to 20 mg, 0.25 to 15 mg, 0.25 to 12 mg, 0.25 to 10 mg, 0.25 to 8 mg, 0.25 mg to 7m g, 0.25 mg to 5 mg, 0.5 mg to 2.5 mg, 1 mg to 20 mg, 1 mg to 15 mg, 1 mg to 12 mg, 1 mg to 10 mg, 1 mg to 8 mg, 1 mg to 7 mg, 1 mg to 5 mg, or 1 mg to 2.5 mg.

The dosages of prophylactic or therapeutic agents other than compounds of the invention, which have been or are currently being used to prevent, treat, manage, or proliferative disorders, such as cancer, or one or more symptoms thereof can be used in the combination therapies of the invention. Preferably, dosages lower than those which have been or are currently being used to prevent, treat, manage, or ameliorate a proliferative disorders, or one or more symptoms thereof, are used in the combination therapies of the invention. The recommended dosages of agents currently used for the prevention, treatment, management, or amelioration of a proliferative disorders, such as cancer, or one or more symptoms thereof, can be obtained from any reference in the art including, but not limited to, Hardman *et al.*, eds., 1996, Goodman & Gilman's The Pharmacological Basis Of Basis Of Therapeutics 9th Ed, Mc-Graw-Hill, New York; Physician's Desk Reference (PDR) 57th Ed., 2003, Medical Economics Co., Inc., Montvale, NJ, which are incorporated herein by reference in its entirety.

In certain embodiments, when the compounds of the invention are administered in combination with another therapy, the therapies (*e.g.*, prophylactic

or therapeutic agents) are administered less than 5 minutes apart, less than 30 minutes apart, 1 hour apart, at about 1 hour apart, at about 1 to about 2 hours apart, at about 2 hours to about 3 hours apart, at about 3 hours to about 4 hours apart, at about 4 hours to about 5 hours apart, at about 5 hours to about 6 hours apart, at about 5 6 hours to about 7 hours apart, at about 7 hours to about 8 hours apart, at about 8 hours to about 9 hours apart, at about 9 hours to about 10 hours apart, at about 10 hours to about 11 hours apart, at about 11 hours to about 12 hours apart, at about 12 hours to 18 hours apart, 18 hours to 24 hours apart, 24 hours to 36 hours apart, 36 hours to 48 hours apart, 48 hours to 52 hours apart, 52 hours to 60 hours apart, 60 10 hours to 72 hours apart, 72 hours to 84 hours apart, 84 hours to 96 hours apart, or 96 hours to 120 hours part. In one embodiment, two or more therapies (*e.g.*, prophylactic or therapeutic agents) are administered within the same patent visit.

In certain embodiments, one or more compounds of the invention and one or more other the therapies (*e.g.*, prophylactic or therapeutic agents) are cyclically 15 administered. Cycling therapy involves the administration of a first therapy (*e.g.*, a first prophylactic or therapeutic agents) for a period of time, followed by the administration of a second therapy (*e.g.*, a second prophylactic or therapeutic agents) for a period of time, followed by the administration of a third therapy (*e.g.*, a third prophylactic or therapeutic agents) for a period of time and so forth, and repeating 20 this sequential administration, *i.e.*, the cycle in order to reduce the development of resistance to one of the agents, to avoid or reduce the side effects of one of the agents, and/or to improve the efficacy of the treatment.

In certain embodiments, administration of the same compound of the invention may be repeated and the administrations may be separated by at least 1 25 day, 2 days, 3 days, 5 days, 10 days, 15 days, 30 days, 45 days, 2 months, 75 days, 3 months, or 6 months. In other embodiments, administration of the same prophylactic or therapeutic agent may be repeated and the administration may be separated by at least at least 1 day, 2 days, 3 days, 5 days, 10 days, 15 days, 30 days, 45 days, 2 months, 75 days, 3 months, or 6 months.

30 In a specific embodiment, the invention provides a method of preventing, treating, managing, or ameliorating a proliferative disorders, such as cancer, or one or more symptoms thereof, said methods comprising administering to a subject in

need thereof a dose of at least 150 µg/kg, preferably at least 250 µg/kg, at least 500 µg/kg, at least 1 mg/kg, at least 5 mg/kg, at least 10 mg/kg, at least 25 mg/kg, at least 50 mg/kg, at least 75 mg/kg, at least 100 mg/kg, at least 125 mg/kg, at least 150 mg/kg, or at least 200 mg/kg or more of one or more compounds of the invention once every day, preferably, once every 2 days, once every 3 days, once every 4 days, once every 5 days, once every 6 days, once every 7 days, once every 8 days, once every 10 days, once every two weeks, once every three weeks, or once a month.

10 F. Other Embodiments

The compounds of the invention may be used as research tools (for example, to evaluate the mechanism of action of new drug agents, to isolate new drug discovery targets using affinity chromatography, as antigens in an ELISA or ELISA-like assay, or as standards in *in vitro* or *in vivo* assays). These and other uses and
15 embodiments of the compounds and compositions of this invention will be apparent to those of ordinary skill in the art.


The invention is further defined by reference to the following examples describing in detail the preparation of compounds of the invention. It will be apparent to those skilled in the art that many modifications, both to materials and
20 methods, may be practiced without departing from the purpose and interest of this invention. The following examples are set forth to assist in understanding the invention and should not be construed as specifically limiting the invention described and claimed herein. Such variations of the invention, including the substitution of all equivalents now known or later developed, which would be within
25 the purview of those skilled in the art, and changes in formulation or minor changes in experimental design, are to be considered to fall within the scope of the invention incorporated herein.

EXAMPLES

30 Reagents and solvents used below can be obtained from commercial sources such as Aldrich Chemical Co. (Milwaukee, Wisconsin, USA). ¹H-NMR and ¹³C-NMR spectra were recorded on a Varian 300MHz NMR spectrometer. Significant

peaks are tabulated in the order: δ (ppm): chemical shift, multiplicity (s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br s, broad singlet), coupling constant(s) in Hertz (Hz) and number of protons.

5 **Example 1: Synthesis of Compound 76**



The hydrazide (M) (1.45 g, 7.39 mmol) and the isothiocyanate (N) (1.59 g, 7.39 mmol) were dissolved in ethanol (20 ml) with heating. When the starting materials were dissolved the solution was allowed to cool to room temperature and a precipitate formed. This precipitate was filtered then washed with ether to provide the intermediate (P) as a white solid (2.85 g, 97%). The intermediate (VII) (1.89 g, 4.77 mmol) was heated in a solution of sodium hydroxide (0.38 g, 9.54 mmol) in water (20 mL) at 110°C for 2 hours. The solution was allowed to cool to room temperature then acidified with conc. HCl. The resulting precipitate was filtered then washed with water (100 mL) and dried. The crude product was recrystallized from ethanol to produce compound 76 as a white solid (1.4 g, 75%).

^1H NMR (DMSO- d_6) δ 9.43-9.53 (bs, 2H), 8.11-8.16 (m, 1H), 7.47-7.55 (m, 2H), 7.38 (d, $J=8.1$ Hz, 1H), 7.31-7.36 (m, 1H), 6.98 (d, $J=8.1$ Hz, 1H), 6.71 (s, 1H), 6.17 (s, 1H), 3.98 (s, 3H), 2.17 (q, $J=7.5$ Hz, 2H), 0.73 (t, $J=7.5$ Hz, 3H);

ESMS calculated for (C₂₁H₁₉N₃O₃S) 393.11; Found 394.1(M+1)⁺.


Example 2: Synthesis of Compound 124

3-(2,4-Dihydroxy-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole (505 mg,
5 1.5 mmol), which is commercially available from Scientific Exchange, Inc., Center
Ossipee, NH 03814, and Et₃N (0.84ml, 6.0 mmol) in 15ml CH₂Cl₂ were treated
dropwise with ethyl isocyanate (360mg, 5.0 mmol) at 0°C. The mixture was then
warmed to room temperature and stirred for 3h. The reaction mixture was diluted
with CH₂Cl₂, washed with H₂O and saturated brine, dried with Na₂SO₄, and
10 concentrated *in vacuo*. The residue was chromatographed (Hexane/ EtOAc 3:1) to
give Compound 124 as a white solid (480 mg, 58%).

¹H-NMR (CDCl₃) δ 10.13 (s, 1H), 7.96 (d, J=9.0 Hz, 2H), 7.61-7.57 (m,
3H), 7.49-7.36(m, 2H), 7.01(s, 1H), 6.88 (d, J=8.4Hz, 1H), 6.70 (d, J=8.4Hz, 1H),
4.98-4.96(m, 2H), 3.56(q, J=7.2 Hz, J=12.6 Hz, 2H), 3.28-3.10(m, 4H), 1.33(t,
15 J=7.2 Hz, 3H), 1.13 (q, J=15.0 Hz, J=7.2Hz, 6H);

ESMS calculated for C₂₇H₂₈N₆O₅S: 548.18; Found: 549.1 (M+1)⁺.

Example 3: Synthesis of Compound 188



1-Benzenesulfonyl-7-methoxy-1H-indole (Q)

- 5 To a solution of 7-methoxyindole (1 eq) in DMF cooled in an ice bath was added NaH (60% dispersion in oil, 1.2 eq). The reaction was stirred for 1 hr at room temperature then recooled in an ice bath. Benzenesulfonyl chloride (1.1 eq) was added then the reaction was stirred for 2 hrs at room temperature. Water/ethyl acetate were added and the ethyl acetate layer was washed repeatedly (3x) with
- 10 water. The ethyl acetate layer was concentrated and evaporated to dryness.

1-Benzenesulfonyl-7-methoxy-4-nitro-1H-indole (R)

To a solution of 1-benzenesulfonyl-7-methoxy-1H-indole (Q) (1 eq) in dichloromethane cooled in an ice bath was added SiO₂-HNO₃ (2 wt eq) in small

portions. The reaction was stirred for 1 hr at room temperature. Activated carbon (2 wt eq) was added then the entire mixture was stirred for 1 hr. The mixture was then filtered and evaporated to dryness. Separation of the isomers was achieved by column chromatography.

5

7-Methoxy-4-nitro-1H-indole (S)

To a solution of 1-benzenesulfonyl-7-methoxy-4-nitro-1H-indole (**R**) (1 eq) in methanol was added a solution of sodium hydroxide (5 eq) in water. The solution was heated to reflux for 3 hrs. Methanol was removed under reduced pressure then water and ethyl acetate were added. The ethyl acetate layer separated and washed repeatedly (3x) with water. The ethyl acetate layer was concentrated and evaporated to dryness to produce the desired product.

10

1-Isopropyl-7-methoxy-4-nitro-1H-indole (T)

To a solution of 7-methoxy-4-nitro-1H-indole (**S**) (1 eq) in DMF cooled in an ice bath was added NaH (60% dispersion in oil, 1.2 eq). The reaction was stirred for 1 hr at room temperature then recooled in an ice bath. 2-Iodopropane (1.1 eq) was added then the reaction was stirred for 2 hrs at room temperature. Water and ethyl acetate were added. The ethyl acetate layer was separated and washed repeatedly (3x) with water. The ethyl acetate layer was concentrated then evaporated to dryness. Further purification by column chromatography produced the pure desired product.

15

20

1-Isopropyl-7-methoxy-1H-indol-4-ylamine (U)

A solution of 1-isopropyl-7-methoxy-4-nitro-1H-indole (**T**) (1eq) and palladium 10% on activated carbon (0.1 wt eq) in methanol/ethyl acetate (1:1) was shaken on a Parr hydrogenation apparatus under hydrogen for 1 hr. The reaction was then filtered through Celite and evaporated to dryness to produce the desired product.

25

30

1-Isopropyl-4-isothiocyanato-7-methoxy-1H-indole (V)

To a solution of 1-isopropyl-7-methoxy-1H-indol-4-ylamine (U) (1eq) in dichloromethane was added 1,1'-thiocarbonyldiimidazole (1.2 eq). The reaction was stirred for 2 hrs at room temperature then evaporated to dryness. Further purification by column chromatography produced the pure desired product.

5


3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-5-mercapto-[1,2,4] triazole (Compound 188)

5-Ethyl-2,4-dihydroxy-benzoic acid hydrazide (W) (1eq) and 1-isopropyl-4-isothiocyanato-7-methoxy-1H-indole (V) (1.01 eq) were heated in ethanol (0.02 M based on isothiocyanate) at 80°C for 1 hr. The solution was allowed to cool to room temperature overnight. The resulting precipitate was filtered, washed with ether, dried and used without further purification (yield 80%). The precipitate was suspended in aqueous NaOH solution (2 eq NaOH) and nitrogen was bubbled through this suspension for 10 min. The reaction was then heated to 110°C for 1 hr under a nitrogen atmosphere then allowed to cool to room temperature. Neutralisation with conc. HCl produced a white precipitate which was filtered and washed with water. Repeated recrystallisation from EtOH/water produced the desired product (purity >95%, yield 50-70%)

¹H-NMR (DMSO-d₆) δ (ppm), 9.52 (s, 1H), 9.42 (s, 1H), 7.40 (d, J=3.3Hz, 1H), 6.82 (d, J=8.4Hz, 1H), 6.61 (s, 1H), 6.20 (s, 1H), 6.05 (d, J=3.3 Hz, 1H), 5.30 (qn, J=6.6Hz, 1H), 3.89 (s, 3H), 2.14 (q, J=7.5Hz, 2H), 1.41-1.47 (m, 6H), 0.68 (t, J=7.5Hz, 3H);

ESMS CALCULATED. FOR C₂₂H₂₄N₄O₃S: 424.16; FOUND: 425.1 (M+1)⁺.

25

Example 4: Synthesis of Compound 223

- 5 2,4-Dimethoxy-5-isopropylbenzoic acid (2.24 g, 10.0 mmol, 1.00 equiv.) in 50 mL CH_2Cl_2 at room temperature was treated with $(\text{COCl})_2$ (1.40 g, 11.0 mmol, 1.10 equiv.) and catalytic amount of DMF (0.1 mL) for 1 hour. Solvent and excess $(\text{COCl})_2$ were removed *in vacuo*. The residue was dissolved in 100 mL CH_2Cl_2 , and
- 10 treated with 1,3-dimethyl-5-aminoindole (1.60 g, 10.0 mmol, 1.00 equiv.) and triethylamine (1.55 g, 15.0 mmol, 1.50 equiv.) at 0°C for one hour. Aqueous workup and removal of solvent gave a light brown solid which was washed with ether to yield off-white solid (2.28g, 6.22 mmol, 62%).

^1H NMR (CDCl_3) δ (ppm) 9.78 (br s, 1H), 8.21 (s, 1H), 8.09 (d, $J = 2.1$ Hz, 1H), 7.31 (dd, $J = 8.7$ Hz, 2.1 Hz, 1H), 7.22 (d, $J = 8.7$ Hz, 1H), 6.82 (s, 1H), 6.50 (s, 1H), 4.09 (s, 3H), 3.92 (s, 3H), 3.73 (s, 3H), 3.26 (hept, $J = 6.9$ Hz, 1H), 2.32 (s, 3H), 1.24 (d, $J = 6.9$ Hz, 6H).

- The off-white solid obtained above was treated with Lawesson's reagent
- 20 (1.51 g, 3.74 mmol, 0.6 equiv.) in 50 mL toluene at 110°C for three hours. Toluene was removed on rotary evaporator and vacuum pump, and the residue was treated

with hydrazine (anhydrous, 3.0 g, 94 mmol, 15.0 equiv.) in 20 mL dioxane at 80°C for 30 minutes. The reaction mixture was extracted with ethyl acetate and water to remove excess hydrazine. The organic layer was dried over MgSO₄, and filtered to remove drying agent. Carbodiimidazole (CDI)(3.02 g, 18.7 mmol, 3.00 equiv.) was added to the solution, and the solution was refluxed (65°C) for 2 hours. Solvent was removed, and the residue was treated with 20 mL THF and 10 mL NaOH (2M) to destroy excess CDI. Extraction with ethyl acetate (EtOAc) and water, followed by chromatography purification gave the desired product 3-(2,4-methoxy-5-isopropyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-hydroxy-[1,2,4] triazole as light brown solid (2.20 g, 5.42 mmol, 87%).

¹H NMR (CDCl₃), δ (ppm) 9.63 (br s, 1H), 7.34 (d, *J* = 2.1 Hz, 1H), 7.20 (s, 1H), 7.18 (d, *J* = 8.4 Hz, 1H), 7.00 (dd, *J* = 8.4 Hz, 2.1 Hz, 1H), 6.80 (s, 1H), 6.19 (s, 1H), 3.76 (s, 3H), 3.69 (s, 3H), 3.40 (s, 3H), 3.15 (hept, *J* = 6.9 Hz, 1H), 2.20 (s, 3H), 1.10 (d, *J* = 6.9 Hz, 6H).

3-(2,4-methoxy-5-isopropyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-hydroxy-[1,2,4] triazole obtained above was treated with pyridine hydrochloride (12.53 g, 108.3 mmol, 20.0 equiv.), NaI (0.812 g, 5.42 mmol, 1.0 equiv.) and 0.5 mL water at 205°C under nitrogen protection for 1 hour. The reaction mixture was treated with 200 mL water. The solid was collected by filtration, washed with 3 x 20 mL water, and dissolved in 50 mL 2M NaOH solution. The aqueous solution was extracted with 100 mL EtOAc, and the EtOAc layer was extracted with 2 x 20 mL 0.5M NaOH. EtOAc layer was discarded. The aqueous layer were combined, neutralized with HCl to PH around 5, and extracted with 3 x 100 mL EtOAc. The combined EtOAc layer was diluted with 50 mL THF, dried over MgSO₄, and filtered through silica gel plug. Most of solvents were removed to form a slurry with around 2 mL of solvent left. Solid was collected by filtration, washed with 2 mL EtOAc, and dried. The desired product 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-hydroxy-[1,2,4] triazole (**Compound 223**) was obtained as an off-white solid (1.75g, 4.63mmol, 85%).

^1H NMR (CD_3OD), δ (ppm) 7.46 (d, $J = 1.8$ Hz, 1H), 7.41 (d, $J = 8.4$ Hz, 1H), 7.04 (dd, $J = 8.4$ Hz, 1.8 Hz, 1H), 7.02 (s, 1H), 6.53 (s, 1H), 6.26 (s, 1H), 3.74 (s, 3H), 2.88 (sept, $J = 6.9$ Hz, 1H), 2.24 (s, 3H), 0.62 (d, $J = 6.9$ Hz, 6H);

5 ESMS calculated. for $\text{C}_{21}\text{H}_{23}\text{N}_4\text{O}_3$: 378.1; Found: 379.1 ($\text{M} + 1$) $^+$.

10 The following compounds were prepared as described above in the section entitled "Methods of Making the Compounds of the Invention" and as exemplified in Examples 1 through 4.

Example 5: Compound 1

15 ESMS calcd for $\text{C}_{18}\text{H}_{13}\text{N}_3\text{OS}$: 319.1; Found: 320.0 ($\text{M} + 1$) $^+$.

Example 6: Compound 2

ESMS calcd for $\text{C}_{21}\text{H}_{19}\text{N}_3\text{O}_4\text{S}$: 409.11; Found: 410.0 ($\text{M} + \text{H}$) $^+$.

Example 7: Compound 5

20 ESMS calcd for $\text{C}_{19}\text{H}_{15}\text{N}_3\text{O}_2\text{S}$: 365.08; Found: 266.0 ($\text{M} + \text{H}$) $^+$.

Example 8: Compound 6

ESMS calcd for $\text{C}_{20}\text{H}_{17}\text{N}_3\text{O}_2\text{S}$: 379.10; Found: 380.0 ($\text{M} + \text{H}$) $^+$.

25 **Example 9: Compound 7**

ESMS calcd for $\text{C}_{21}\text{H}_{19}\text{N}_3\text{O}_2\text{S}$: 393.11; Found: 394.0 ($\text{M} + \text{H}$) $^+$.

Example 10: Compound 8

30 ESMS calcd for $\text{C}_{21}\text{H}_{19}\text{N}_3\text{O}_3\text{S}$: 393.11; Found: 394.0 ($\text{M} + \text{H}$) $^+$.

Example 11: Compound 9

ESMS calcd for $\text{C}_{21}\text{H}_{19}\text{N}_3\text{O}_2\text{S}$: 393.11; Found: 394.0 ($\text{M} + \text{H}$) $^+$.

NOT FURNISHED UPON FILING

ESMS calcd for $C_{15}H_{13}N_3O_3S$: 315.07; Found: 316.0 (M+H)⁺.

Example 21: Compound 22

ESMS calcd for $C_{15}H_{13}N_3O_3S$: 315.07; Found: 316.0 (M+H)⁺.

5

Example 22: Compound 23

ESMS calcd for $C_{14}H_{10}FN_3O_2S$: 303.05; Found: 304.0 (M+H)⁺.

Example 23: Compound 23

10 1H NMR (DMSO- d_6) δ 9.69 (s, 1H), 9.65 (s, 1H), 7.16 (d, $J=7.2$ Hz, 1H),
7.05 (t, $J=7.2$ Hz, 1H), 6.93 (d, $J=8.1$ Hz, 2H), 6.11-6.16 (m, 2H), 2.21 (s, 3H), 1.89
(s, 3H);

ESMS Calcd $C_{16}H_{15}N_3O_2S$: 313.09, Found 314.1(M+1)⁺.

15 **Example 24: Compound 24**

ESMS calcd for $C_{16}H_{15}N_3O_2S$: 313.09; Found: 314.0 (M+H)⁺.

Example 25: Compound 25

20 1H NMR (DMSO- d_6) δ 10.44 (m, 1H), 8.00-7.95 (m, 2H), 7.55-7.37 (m, 5H),
6.61 (d, $J=7.8$ and 1.8 Hz, 1H), 6.51 (t, $J=8.6$ Hz, 1H), 6.41(d, $J=10.8$ Hz, 1H);

ESMS calcd for $C_{18}H_{12}FN_3OS$: 337.07; Found: 338.0 (M+1)⁺.

Example 26: Compound 26

25 1H NMR (DMSO- d_6) δ 9.57 (s, 1H), 7.99 (d, $J=8.4$ Hz, 1H), 7.96 (d, $J=6.9$
Hz, 1H), 7.55-7.37 (m, 5H), 6.61 (d, $J=8.1$ Hz, 1H), 5.83 (d, $J=2.1$ Hz, 1H),
5.73(dd, $J=8.1$ and 1.8 Hz, 1H), 5.24 (s, 2H);

ESMS calcd for $C_{18}H_{14}N_4OS$: 334.09; Found: 335.0 (M+1)⁺.

Example 27: Compound 27

30 ESMS calcd for $C_{18}H_{19}N_3O_2S$: 341.12; Found: 342.0 (M+H)⁺.

Example 28: Compound 28

ESMS calcd for $C_{16}H_{15}N_3O_2S$: 313.09; Found: 314.0 (M+H)⁺.

Example 29: Compound 29

ESMS calcd for $C_{16}H_{15}N_3O_2S$: 313.09; Found: 314.0 (M+H)⁺.

5

Example 30: Compound 30

ESMS calcd for $C_{16}H_{15}N_3O_2S$: 313.09; Found: 314.0 (M+H)⁺.

Example 31: Compound 31

10

ESMS calcd for $C_{14}H_{10}FN_3O_2S$: 303.05; Found: 304.0 (M+H)⁺.

Example 32: Compound 32

ESMS calcd for $C_{15}H_{13}N_3O_2S$: 331.04; Found: 332.0 (M+H)⁺.

15

Example 33: Compound 33

ESMS calcd for $C_{18}H_{13}N_3O_2S$: 335.07; Found: 336.0 (M+H)⁺.

Example 34: Compound 34

ESMS calcd for $C_{16}H_{15}N_3O_2S$: 313.09; Found: 314.0 (M+H)⁺.

20

Example 35: Compound 35

ESMS calcd for $C_{15}H_{12}FN_3O_2S$: 317.06; Found: 317.0 (M+H)⁺.

Example 36: Compound 36

25

ESMS calcd for $C_{20}H_{15}N_3O_2S$: 361.1; Found: 362.0 (M+1)⁺.

Example 37: Compound 37

¹H NMR (DMSO-d₆) δ 10.03 (s, 1H), 8.00-7.96 (m, 2H), 7.55-7.37 (m, 5H),
7.00 (d, J = 8.1 Hz, 1H), 6.20 (m, 2H), 3.57 (s, 3H);

30

ESMS calcd for $C_{19}H_{15}N_3O_2S$: 349.09; Found: 350.0 (M+1)⁺.

Example 38: Compound 38

ESMS calcd for $C_{14}H_9Cl_2N_3O_2S$: 352.98; Found: 353.9 (M+H)⁺.

Example 39: Compound 39

- 5 ¹H NMR (DMSO-d₆) δ 9.74 (s, 1H), 9.63 (s, 1H), 8.14 (m, 1H), 7.52-7.48 (m, 2H), 7.37 (d, *J* = 8.4 Hz, 1H), 7.32 (m, 1H), 6.96 (d, = 8.1 Hz, 1H), 6.90 (d, = 8.4 Hz, 1H), 6.08 (d, = 1.9 Hz, 1H), 6.01 (d, = 8.4 Hz, 1H), 3.98 (s, 3H);

ESMS calcd for $C_{19}H_{15}N_3O_3S$: 365.08; Found: 366.0 (M+1)⁺.

10 **Example 40: Compound 40**

ESMS calcd for $C_{25}H_{16}N_3O_2S$: 409.09; Found: 410.0 (M+1)⁺.

Example 41: Compound 42

- 15 ¹H NMR (DMSO-d₆) δ 9.75(s, 1H), 9.67(s, 1H), 7.08(s, 2H), 6.96-6.94(m, 2H), 6.18-6.13(m, 2H), 2.72-2.50(m, 3H), 2.35-2.28(m, 1H), 1.64-1.60(m, 4H);

ESMS calcd for $C_{18}H_{17}N_3O_2S$: 339.10; Found: 340.0 (M+1)⁺.

Example 42: Compound 43

- 20 ESMS calcd for $C_{22}H_{15}N_3O_2S$: 385.09; Found: 386.0 (M+1)⁺.

Example 43: Compound 44

ESMS calcd for $C_{20}H_{15}N_3O_2S$: 361.09; Found: 362.0 (M+1)⁺.

Example 44: Compound 45

- 25 ESMS calcd for $C_{19}H_{15}N_3O_2S$: 349.09; Found: 350.0 (M+1)⁺.

Example 45: Compound 46

ESMS calcd for $C_{19}H_{21}N_3O_3S$: 371.13; Found: 372.0 (M+1)⁺.

30 **Example 46: Compound 47**

ESMS calcd for $C_{22}H_{27}N_3O_3S$: 413.18; Found: 414.1 (M+1)⁺.

Example 47: Compound 48

ESMS calcd for $C_{18}H_{12}ClN_3O_2S$: 369.03; Found: 370.0 (M+H)⁺.

Example 48: Compound 49

5 ¹H NMR (DMSO-d₆) δ 9.49 (s, 1H), 9.40 (s, 1H), 7.94-7.99 (m, 2H), 7.38-7.56 (m, 5H), 6.70 (s, 1H), 6.13 (s, 1H), 2.12 (q, *J*=7.2 Hz, 2H), 0.71 (t, *J*=7.2 Hz, 3H);

ESMS Calcd for $C_{20}H_{17}N_3O_2S$: 363.10, Found 364.1(M+1)⁺.

10 **Example 49: Compound 50**

ESMS calcd for $C_{20}H_{15}N_3O_5S$: 409.07; Found: 410.0 (M+H)⁺.

Example 50: Compound 51

ESMS calcd for $C_{18}H_{14}N_4O_2S$: 350.08; Found: 351.0 (M+H)⁺.

15

Example 51: Compound 52

ESMS calcd for $C_{17}H_{12}N_4OS$: 320.07; Found: 320.9 (M+H)⁺.

Example 52: Compound 53

20 ¹H NMR (CDCl₃) δ 12.0 (br s, 1H), 9.87 (br s, 1H), 9.83 (br s, 1H), 7.97 (d, *J* = 8.1 Hz, 2H), 7.41-7.56 (m, 5H), 7.13 (d, *J* = 1.5 Hz, 1H), 7.07 (d, *J* = 8.7 Hz, 1H), 6.71 (dd, *J* = 1.8 Hz, 8.1 Hz, 1H), 1.93 (s, 3H);

ESMS calcd for $C_{20}H_{17}N_4O_2S$: 376.1; Found: 377.0(M+1)⁺.

25 **Example 53: Compound 56**

ESMS calcd for $C_{16}H_{15}N_3O_4S$: 345.08; Found: 346.0 (M+1)⁺.

Example 54: Compound 57

ESMS calcd for $C_{18}H_{16}N_4O_2S$: 352.10; Found: 353.0 (M+1)⁺.

30

Example 55: Compound 61

^1H NMR (DMSO- d_6) δ 9.66(s, 1H), 9.60(s, 1H), 7.29-7.27(m, 1H), 7.12-7.10(m, 2H), 7.03-7.00(m, 1H), 6.19-6.17(m, 2H), 1.18(s, 18H);

ESMS calcd for $\text{C}_{22}\text{H}_{27}\text{N}_3\text{O}_2\text{S}$: 397.18; Found: 398.1 (M+1) $^+$.

5 **Example 56: Compound 64**

ESMS calcd for $\text{C}_{21}\text{H}_{15}\text{N}_3\text{O}_3\text{S}$: 389.08; Found: 390.0 (M+H) $^+$.

Example 57: Compound 65

ESMS calcd for $\text{C}_{19}\text{H}_{13}\text{N}_3\text{O}_4\text{S}$: 379.06; Found: 380.0 (M+1) $^+$.

10

Example 58: Compound 66

ESMS calcd for $\text{C}_{21}\text{H}_{18}\text{N}_4\text{O}_3\text{S}$: 406.11; Found: 407.0 (M+1) $^+$.

Example 59: Compound 67

15

ESMS calcd for $\text{C}_{21}\text{H}_{19}\text{N}_3\text{O}_3\text{S}$: 393.11; Found: 394.0 (M+1) $^+$.

Example 60: Compound 68

ESMS calcd for $\text{C}_{21}\text{H}_{19}\text{N}_3\text{O}_3\text{S}$: 393.11; Found: 394.0 (M+1) $^+$.

20

Example 61: Compound 69

ESMS calcd for $\text{C}_{21}\text{H}_{19}\text{N}_3\text{O}_3\text{S}$: 393.11; Found: 394.0 (M+1) $^+$.

Example 62: Compound 70

ESMS calcd for $\text{C}_{17}\text{H}_{12}\text{N}_4\text{O}_2\text{S}$: 336.07; Found: 337.0 (M+H) $^+$.

25

Example 63: Compound 71

ESMS calcd for $\text{C}_{21}\text{H}_{19}\text{N}_3\text{O}_3\text{S}$: 393.11; Found: 394.0 (M+1) $^+$.

Example 64: Compound 72

30

^1H NMR (DMSO- d_6) δ 10.3 (br s, 1H), 7.95-8.19 (m, 2H), 7.48-7.72 (m, 5H), 7.17 (d, J = 8.4 Hz, 1H), 6.44 (d, J = 8.4 Hz, 1H), 5.95 (d, J = 2.1 Hz, 1H), 5.73 (dd, J = 2.1 Hz, 8.4 Hz, 1H), 5.47 (br s, 1H), 3.62 (s, 3H);

ESMS calcd for $C_{19}H_{17}N_4O_2S_2$: 412.1; Found: 413.0(M+1)⁺.

Example 65: Compound 73

¹H NMR (DMSO-d₆) δ 9.37 (s, 1H), 8.94 (s, 1H), 7.94-7.98 (m, 2H), 7.43-
5 7.60 (m, 5H), 5.97 (s, 1H), 1.85 (s, 3H), 1.81 (s, 3H);

ESMS calcd for $C_{20}H_{18}N_3O_2S$: 363.1; Found: 364.0(M+1)⁺.

Example 66: Compound 74

ESMS calcd for $C_{21}H_{19}N_3O_4S$: 409.11; Found: 410.0 (M+H)⁺.
10

Example 67: Compound 75

¹H NMR (DMSO-d₆) δ 9.46 (s, 1H), 9.45 (s, 1H), 7.95-8.00 (m, 2H), 7.38-
7.56 (m, 5H), 6.65 (s, 1H), 6.15 (s, 1H), 2.07-2.14 (m, 2H), 0.81-1.18 (m, 11H);

ESMS calcd for $C_{24}H_{26}N_3O_2S$: 419.1; Found: 420.1(M+1)⁺.
15

Example 68: Compound 76

ESMS calcd for $C_{21}H_{19}N_3O_3S$: 393.11; Found: 394.0 (M+H)⁺.

Example 69: Compound 77

ESMS calcd for $C_{21}H_{19}N_3O_3S$: 393.11; Found: 394.0 (M+H)⁺.
20

Example 70: Compound 78

¹H NMR (DMSO-d₆) δ 9.71 (s, 1H), 9.35 (s, 1H), 7.98-8.04 (m, 2H), 7.50-
7.62 (m, 5H), 6.58 (s, 1H), 2.15 (q, *J* = 7.5 Hz, 2H), 0.58 (t, *J* = 7.5 Hz, 3H);

ESMS calcd for $C_{20}H_{17}ClN_3O_2S$: 397.0; Found: 398.0(M+1)⁺.
25

Example 71: Compound 79

ESMS calcd for $C_{19}H_{21}N_3O_3S$: 371.13; Found: 372.0 (M+H)⁺.

Example 72: Compound 80

ESMS calcd for $C_{21}H_{19}N_3O_2S$: 393.11; Found: 394.0 (M+H)⁺.
30

Example 73: Compound 81

ESMS calcd for $C_{20}H_{17}N_3O_2S$: 379.10; Found: 380.0 (M+H)⁺.

Example 74: Compound 82

5 ESMS calcd for $C_{21}H_{19}N_3O_2S$: 393.11; Found: 394.0 (M+H)⁺.

Example 75: Compound 83

ESMS calcd for $C_{20}H_{17}N_3O_3S$: 379.10; Found: 380.0 (M+H)⁺.

Example 76: Compound 84

10 ESMS calcd for $C_{20}H_{17}N_3O_3S$: 379.10; Found: 380.0 (M+H)⁺.

Example 77: Compound 85

ESMS calcd for $C_{19}H_{15}N_3O_2S$: 365.08; Found: 266.0 (M+H)⁺.

15

Example 78: Compound 86

¹H NMR (DMSO-d₆) δ 9.68 (s, 1H), 9.58 (s, 1H), 8.2 (dd, J = 7.0 and 2.4 Hz, 1H), 7.50 (m, 2H), 7.40 (tr, J = 8.1 Hz, 1H), 7.32 (m, 1H), 6.97 (d, J = 7.5 Hz, 1H), 6.95 (m, 1H), 6.89 (d, = 8.4 Hz, 1H), 6.08 (d, = 2.1 Hz, 1H), 6.0 (dd, = 7.4 and 2.1 Hz, 1H), 3.96 (s, 3H);

20

ESMS calcd for $C_{19}H_{15}N_3O_3S$: 365.08; Found: 366.0 (M+1)⁺.

Example 79: Compound 87

¹H NMR (MeOH-d₄) δ 8.25 (m, 1H), 7.96 (s, 1H), 7.46-7.44 (m, 2H), 7.26 (d, J = 8.4 Hz, 1H), 6.83 (d, J = 8.1 Hz, 1H), 6.70 (d, J = 8.7 Hz, 1H), 6.17 (d, J = 2.1 Hz, 1H), 5.98 (dd, J = 8.4 and 2.4 Hz, 1H);

25

ESMS calcd for $C_{18}H_{13}N_3O_3S$: 351.07; Found: 352.0 (M+1)⁺.

Example 80: Compound 88

¹H-NMR (DMSO-d₆) δ 9.69 (s, 1H), 9.59 (s, 1H), 7.54 (d, J=8.1Hz, 1H), 7.46 (d, J=3Hz, 1H), 7.14 (t, J=7.8 Hz, 1H), 6.97 (d, J=7.2 Hz, 1H), 6.89 (d,

30

$J=8.7\text{Hz}$, 1H), 6.12-6.13 (m, 2H), 6.02 (dd, $J_1=2.4\text{ Hz}$, $J_2=8.4\text{ Hz}$, 1H), 4.74 (qn, $J=6.6\text{Hz}$, 1H), 1.40-1.46 (m, 6H);

ESMS calcd for $\text{C}_{19}\text{H}_{18}\text{N}_4\text{O}_2\text{S}$: 366.12; Found: 367.1 (M+1)⁺.

5 **Example 81: Compound 89**

ESMS calcd for $\text{C}_{22}\text{H}_{21}\text{N}_3\text{O}_2\text{S}$: 391.14; Found: 392.0 (M+H)⁺.

Example 82: Compound 90

10 ^1H NMR (DMSO- d_6) δ 9.47 (s, 1H), 9.43 (s, 1H), 7.94-8.00 (m, 2H), 7.39-7.57 (m, 5H), 6.68 (s, 1H), 6.15 (s, 1H), 2.05-2.15 (m, 2H), 1.05-1.17 (m, 2H), 0.50 (t, $J = 7.5\text{ Hz}$, 3H); ESMS calcd for $\text{C}_{21}\text{H}_{20}\text{N}_3\text{O}_2\text{S}$: 377.1; Found: 378.0(M+1)⁺.

Example 83: Compound 91

15 ^1H NMR (DMSO- d_6) δ 9.15 (s, 1H), 8.50 (s, 1H), 8.00 -8.07 (m, 2H), 7.47-7.63 (m, 5H), 6.27 (s, 1H), 2.06 (q, $J = 7.5\text{ Hz}$, 2H), 1.93 (s, 3H), 0.45 (t, $J = 7.5\text{ Hz}$, 3H);

ESMS calcd for $\text{C}_{21}\text{H}_{20}\text{N}_3\text{O}_2\text{S}$: 377.1; Found: 378.0(M+1)⁺.

Example 84: Compound 93

20 ESMS calcd for $\text{C}_{16}\text{H}_{15}\text{N}_3\text{O}_4\text{S}$: 345.08; Found: 346.0 (M+H)⁺.

Example 85: Compound 95

ESMS calcd for $\text{C}_{16}\text{H}_{12}\text{N}_4\text{O}_2\text{S}$: 324.07; Found: 325.0 (M+H)⁺.

25 **Example 86: Compound 96**

ESMS calcd for $\text{C}_{19}\text{H}_{18}\text{N}_4\text{O}_3\text{S}$: 382.11; Found: 383.0 (M+H)⁺.

Example 87: Compound 98

30 ESMS calcd for $\text{C}_{17}\text{H}_{12}\text{N}_4\text{O}_2\text{S}$: 336.07; Found: 337.0 (M+H)⁺.

Example 88: Compound 99

ESMS calcd for $\text{C}_{19}\text{H}_{13}\text{N}_3\text{O}_4\text{S}$: 379.06; Found: 379.9 (M+H)⁺.

Example 89: Compound 100

¹H-NMR (DMSO-d₆) δ 9.52 (s, 1H), 9.42 (s, 1H), 7.56 (d, *J*=8.7Hz, 1H), 7.49 (d, *J*=3.3Hz, 1H), 7.14 (t, *J*=7.5 Hz, 1H), 6.95 (d, *J*=8.4Hz, 1H), 6.61 (s, 1H),
5 6.21 (s, 1H), 6.14 (dd, *J*=3.3Hz, 1H), 4.76 (qn, *J*=6.6Hz, 1H), 2.14 (q, *J*=7.5Hz, 2H), 1.41-1.47 (m, 6H), 0.66 (t, *J*=7.5Hz, 3H);

ESMS calcd for C₂₁H₂₂N₄O₂S: 394.15; Found: 395.1 (M+1)⁺.

Example 90: Compound 101

10 ESMS calcd for C₁₉H₁₇N₅O₃S: 395.11; Found: 396.0 (M+H)⁺.

Example 91: Compound 102

ESMS calcd. for C₁₉H₂₀N₅O₂S: 381.1; Found: 382.0 (M + 1)⁺.

Example 92: Compound 103

15 ¹H NMR (DMSO-d₆) δ 9.48 (s, 1H), 9.38 (s, 1H), 7.29(d, *J* = 8.4 Hz, 1H), 7.25(d, *J* = 1.8 Hz, 1H), 6.85-6.89 (m, 2H), 6.18 (s, 1H), 3.61 (s, 3H), 2.30 (s, 3H), 2.29 (q, *J* = 7.5 Hz, 2H), 2.09 (s, 3H), 0.94 (t, *J* = 7.5 Hz, 3H);

ESMS calcd for C₂₁H₂₃N₄O₂S: 394.1; Found: 395.0(M+1)⁺.

20

Example 93: Compound 104

ESMS calcd for C₁₉H₁₅N₃O₃S: 365.08; Found: 366.0 (M+H)⁺.

Example 94: Compound 106

25 ESMS calcd for C₂₀H₁₇N₄O₂S: 377.1; Found: 378.0(M+H)⁺.

Example 95: Compound 107

ESMS calcd for C₁₈H₁₃ClN₃O₂S: 369.0; Found: 370.0(M+H)⁺.

Example 96: Compound 116

30 ¹H NMR (DMSO-d₆) δ 7.98-7.56 (m, 2H), 7.55-7.30 (m, 6H), 6.43 (dd, *J* = 8.1 and 1.8 Hz, 1H), 6.29 (m, 1H), 3.65 (s, 3H), 3.16 (s, 3H);

ESMS calcd for $C_{20}H_{17}N_3O_2S$: 363.10; Found: 364.0 (M+1)⁺.

Example 97: Compound 117

¹H-NMR (CDCl₃) δ 7.83(d, J=8.1 Hz, 2H), 7.48-7.34(m, 4H), 7.28-7.20(m, 1H), 6.99 (d, J=1.8Hz, 1H), 6.80(d, J=8.7Hz, 1H), 6.62-6.58(m, 1H), 2.94(s, 3H), 2.89(s, 3H), 2.84(s, 3H), 2.81(s, 3H), 2.75-2.69(m, 6H);

ESMS calcd for $C_{27}H_{28}N_6O_5S$: 548.18; Found: 549.2 (M+1)⁺.

Example 98: Compound 122

¹H-NMR (CDCl₃) δ 7.98(m, 2H), 7.60-7.55(m, 3H), 7.51-7.45(m, 1H), 7.36-7.33(m, 1H), 6.98-6.97(m, 1H), 6.86(d, J=9.9Hz, 1H), 6.70-6.67(m, 1H), 2.86(s, 3H), 2.26(s, 3H), 2.21(s, 3H);

ESMS calcd for $C_{24}H_{19}N_3O_5S$: 461.10; Found: 462.0 (M+1)⁺.

Example 99: Compound 125

ESMS calcd for $C_{20}H_{17}N_3O_3S$: 379.10; Found: 380.0 (M+H)⁺.

Example 100: Compound 126

ESMS calcd for $C_{10}H_{11}N_3O_2S$: 237.06; Found: 238.0 (M+H)⁺.

Example 101: Compound 127

ESMS calcd for $C_{11}H_{13}N_3O_2S$: 251.07; Found: 252.0 (M+H)⁺.

Example 102: Compound 128

ESMS calcd for $C_{11}H_{13}N_3O_2S$: 251.07; Found: 252.0 (M+H)⁺.

Example 103: Compound 129

ESMS calcd for $C_{11}H_{11}N_3O_2S$: 249.06; Found: 250.0 (M+H)⁺.

Example 104: Compound 130

ESMS calcd for $C_{12}H_{15}N_3O_2S$: 265.09; Found: 266.0 (M+H)⁺.

Example 105: Compound 131

ESMS calcd for $C_{20}H_{15}N_3O_4S$: 393.08; Found: 394.1 (M+H)⁺.

Example 106: Compound 177

5 ¹H NMR (DMSO-d₆) δ 9.34(s, 1H), 9.22 (s, 1H), 8.01-7.96 (m, 2H), 7.58-7.44 (m, 5H), 6.56 (s, 1H), 6.14 (s, 1H), 3.29 (s, 3H);

ESMS calcd for $C_{19}H_{15}N_3O_3S$: 365.08; Found: 366.0(M+1)⁺.

Example 107: Compound 178

10 ¹H NMR (DMSO-d₆) δ 10.29 (s, 1H), 9.49 (s, 1H), 9.42 (s, 1H), 8.16 (t, *J* = 5.1 Hz, 1H), 7.45-7.43 (m, 2H), 7.26 (t, *J* = 8.0 Hz, 1H), 6.84 (d, *J* = 7.8 Hz, 1H), 6.75 (d, *J* = 8.7 Hz, 1H), 6.66 (s, 1H), 6.14 (s, 1H), 2.12 (q, *J* = 7.5 Hz, 2H), 0.70 (t, *J* = 7.2 Hz, 3H);

ESMS calcd for $C_{20}H_{17}N_3O_3S$: 379.10; Found: 379.9 (M+1)⁺.

15

Example 108: Compound 179

ESMS calcd for $C_{19}H_{15}N_3O_2S$: 349.09; Found: 350.0 (M+1)⁺.

Example 109: Compound 180

20 ESMS calcd for $C_{19}H_{15}N_3O_2S$: 349.09; Found: 350.0 (M+H)⁺.

Example 110: Compound 181

ESMS calcd for $C_{20}H_{15}N_3O_2S$: 361.09; Found: 362.0 (M+H)⁺.

Example 111: Compound 182

25 ESMS calcd for $C_{16}H_{15}N_3O_3S$: 329.08; Found: 330.0 (M+H)⁺.

Example 112: Compound 183

ESMS calcd for $C_{20}H_{17}N_3O_2S$: 363.10; Found: 364.0 (M+H)⁺.

30

Example 113: Compound 184

ESMS calcd for $C_{18}H_{13}N_3O_3S$: 350.38; Found: 351.9(M+H)⁺.

Example 114: Compound 185

ESMS calcd. for $C_{20}H_{21}N_4O_2S$: 380.1; Found: 381.0 ($M + 1$)⁺.

5 Example 115: Compound 187

ESMS calcd. for $C_{19}H_{20}N_5O_2S$: 381.1; Found: 382.0 ($M + 1$)⁺.

Example 116: Compound 190

ESMS CALCD. FOR $C_{21}H_{22}N_4O_2S$: 394.15; FOUND: 395.0 ($M+1$)⁺.

10

Example 117: Compound 191

ESMS calcd. for $C_{22}H_{23}N_4O_4S$: 438.1; Found: 439.0 ($M + 1$)⁺.

Example 118: Compound 192

15 ESMS calcd. for $C_{20}H_{22}N_5O_2S$: 395.1; Found: 396.0 ($M + 1$)⁺.

Example 119: Compound 193

ESMS calcd. for $C_{20}H_{22}N_5O_2S$: 395.1; Found: 396.0 ($M + 1$)⁺.

20 Example 120: Compound 194

ESMS calcd. for $C_{23}H_{27}N_4O_2S$: 422.1; Found: 423.0 ($M + 1$)⁺.

Example 121: Compound 195

ESMS calcd. for $C_{23}H_{25}N_4O_2S$: 420.1; Found: 421.0 ($M + 1$)⁺.

25

Example 122: Compound 196

ESMS calcd. for $C_{25}H_{29}N_4O_2S$: 448.1; Found: 449.3 ($M + 1$)⁺.

Example 123: Compound 197

30 ESMS calcd. for $C_{22}H_{24}N_4O_2S$: 408.16; Found: 409.2 ($M+1$)⁺.

Example 124: Compound 198

ESMS calcd. for $C_{23}H_{26}N_4O_2S$: 422.18; Found: 423.3 (M+1)⁺.

Example 125: Compound 199

ESMS calcd. for $C_{24}H_{28}N_4O_2S$: 436.19; Found: 437.3 (M+1)⁺.

5

Example 126: Compound 200

ESMS calcd. for $C_{22}H_{22}N_4O_2S$: 406.15; Found: 407.2 (M+1)⁺.

Example 127: Compound 201

10 ESMS calcd. for $C_{23}H_{24}N_4O_3S$: 436.16; Found: 437.3 (M+1)⁺.

Example 128: Compound 202

ESMS calcd. for $C_{22}H_{23}N_4O_2S$: 406.1; Found: 407.0 (M + H)⁺.

15 **Example 129: Compound 204**

ESMS calcd. for $C_{24}H_{28}N_4O_3S$: 452.19; Found: 453.2 (M+1)⁺.

Example 130: Compound 205

ESMS calcd. for $C_{23}H_{24}N_4O_3S$: 436.16; Found: 437.1 (M+1)⁺.

20

Example 131: Compound 206

ESMS calcd. for $C_{21}H_{23}N_4O_2S$: 394.1; Found: 395.1 (M + 1)⁺.

Example 132: Compound 207

25 ESMS calcd. for $C_{20}H_{21}N_4O_2S$: 380.1; Found: 381.1 (M + 1)⁺.

Example 133: Compound 208

ESMS calcd. for $C_{23}H_{26}N_4O_3S$: 438.17; Found: 439.1 (M+1)⁺.

30 **Example 134: Compound 209**

ESMS calcd. for $C_{22}H_{24}N_4O_2S$: 408.1; Found: 409.1 (M + 1)⁺.

Example 135: Compound 210

ESMS calcd. for $C_{24}H_{23}N_4O_2S$: 430.1; Found: 431.1 (M + 1)⁺.

Example 136: Compound 211

5 ESMS calcd. for $C_{21}H_{22}N_4O_3S$: 410.14; Found: 411.1 (M+1)⁺.

Example 137: Compound 212

ESMS calcd. for $C_{23}H_{26}N_4O_3S$: 438.17; Found: 439.1 (M+1)⁺.

10 **Example 138: Compound 213**

ESMS calcd. for $C_{20}H_{21}N_4O_2S$: 380.1; Found: 381.1 (M + 1)⁺.

Example 139: Compound 214

ESMS calcd. for $C_{19}H_{19}N_4O_2S$: 366.1; Found: 367.1 (M + 1)⁺.

15

Example 140: Compound 215

ESMS calcd. for $C_{20}H_{19}N_3O_4S$: 397.1; Found: 398.1 (M+1)⁺.

Example 141: Compound 216

20 ¹H NMR (DMSO-d₆): δ (ppm) 9.56 (s, 1H), 9.40 (s, 1H), 8.03 (d, J = 2.4 Hz, 1H), 7.58 (d, J = 8.4 Hz, 1H), 7.54 (d, J = 2.1 Hz, 1H), 7.11 (dd, J = 8.4, 2.1 Hz, 1H), 6.97 (d, J = 2.4 Hz, 1H), 6.89 (s, 1H), 6.17 (s, 1H), 2.23 (q, J = 7.2 Hz, 2H), 0.93 (t, J = 7.2 Hz, 3H);

ESMS calcd. for $C_{18}H_{15}N_3O_3S$: 353.08; Found: 354.0 (M+1)⁺.

25

Example 142: Compound 217

¹H NMR (DMSO-d₆): δ (ppm) 9.59 (s, 1H), 9.43 (s, 1H), 7.67 (d, J = 8.7 Hz, 1H), 7.54 (d, J = 2.1 Hz, 1H), 7.20 (dd, J = 8.4, 2.1 Hz, 1H), 6.96 (s, 1H), 6.18 (s, 1H), 2.60 (s, 3H), 2.34 (q, J = 7.2 Hz, 2H), 0.98 (t, J = 7.2 Hz, 3H);

30 ESMS calcd. for $C_{18}H_{16}N_4O_3S$: 368.09; Found: 369.0 (M+1)⁺.

Example 143: Compound 218

ESMS calcd. for $C_{21}H_{23}N_4O_2S$: 394.1; Found: 395.1 ($M + 1$)⁺.

Example 144: Compound 219

ESMS calcd. for $C_{21}H_{21}N_4O_2S$: 392.1; Found: 393.1 ($M + 1$)⁺.

5

Example 145: Compound 220

ESMS calcd. for $C_{20}H_{21}N_4O_3$: 364.1; Found: 365.1 ($M + 1$)⁺.

Example 146: Compound 221

10 ESMS calcd. for $C_{20}H_{21}N_4O_2S$: 379.1; Found: 381.1 ($M + 1$)⁺.

Example 147: Compound 222

ESMS calcd. for $C_{21}H_{23}N_4O_2S$: 394.1; Found: 395.1($M + 1$)⁺.

15 **Example 148: Compound 224**

ESMS calcd. for $C_{19}H_{21}N_4O_2S$: 368.1; Found: 369.1 ($M + 1$)⁺.

Example 149: Compound 225

ESMS calcd. for $C_{19}H_{19}N_4O_2S$: 366.1; Found: 367.1($M + 1$)⁺.

20

Example 150: Compound 226

ESMS calcd. for $C_{20}H_{21}N_4O_3$: 364.1; Found: 365.1 ($M + 1$)⁺.

Example 151: Compound 227

25 ESMS calcd. for $C_{21}H_{22}N_4O_2S$: 394.15; Found: 395.1 ($M+1$)⁺.

Example 152: Compound 228

ESMS calcd. for $C_{22}H_{24}N_4O_2S$: 408.16; Found: 409.1 ($M+1$)⁺.

30 **Example 153: Compound 229**

ESMS calcd. for $C_{20}H_{18}F_3N_5O_2S$: 449.11; Found: 450.1 ($M+1$)⁺.

Example 154: Compound 230

ESMS calcd. for $C_{19}H_{19}N_5O_2S$: 381.13; Found: 382.1 (M+1)⁺.

Example 155: Compound 231

5 ESMS calcd. for $C_{19}H_{19}N_5O_2S$: 381.13; Found: 382.1 (M+1)⁺.

Example 156: Compound 232

ESMS calcd. for $C_{22}H_{24}N_4O_3S$: 392.18; Found: 393.1 (M+1)⁺.

10 **Example 157: Compound 233**

ESMS calcd. for $C_{18}H_{17}N_3O_4S$: 371.09; Found: 372.1 (M+1)⁺.

Example 158: Compound 234

ESMS calcd. for $C_{20}H_{21}N_3O_2S$: 367.14; Found: 368.1 (M+1)⁺.

15

Example 159: Compound 235

ESMS calcd. for $C_{19}H_{19}N_5O_2S$: 381.13; Found: 382.1 (M+1)⁺.

Example 160: Compound 239

20 ESMS clcd for $C_{19}H_{21}N_4O_2S$: 368.1; Found: 369.1 (M + H)⁺.

Example 161: Compound 240

ESMS clcd for $C_{18}H_{16}N_4O_3S$: 368.09.10; Found: 369.1 (M+H)⁺.

25 **Example 162: Compound 241**

ESMS clcd for $C_{17}H_{15}N_5O_3S$: 369.09; Found: 370.1 (M+H)⁺.

Example 163: Compound 242

ESMS clcd for $C_{19}H_{18}N_4O_3S$: 382.11; Found: 383.1 (M+H)⁺.

30

Example 164: Compound 243

ESMS clcd for $C_{22}H_{26}N_4O_3S$: 426.17; Found: 427.1 (M+H)⁺.

Example 165: Compound 244

ESMS clcd for $C_{18}H_{16}N_4O_4S$: 384.09; Found: 385.1 (M+H)⁺

Example 166: Compound 245

ESMS clcd for $C_{18}H_{16}N_4O_3S_2$: 400.07; Found: 401.1 (M+H)⁺

Example 167: Compound 245

ESMS clcd for $C_{17}H_{14}N_4O_3S_2$: 386.05; Found: 387.0 (M+H)⁺.

10

Example 168: Inhibition of Hsp90

Hsp90 protein was obtained from Stressgen (Cat#SPP-770). Assay buffer: 100 mM Tris-HCl, Ph7.4, 20 mM KCl, 6 mM MgCl₂. Malachite green (0.0812% w/v) (M9636) and polyvinyl alcohol USP (2.32% w/v) (P1097) were obtained from
15 Sigma. A Malachite Green Assay (see Methods Mol Med, 2003, 85:149 for method details) was used for examination of ATPase activity of Hsp90 protein. Briefly, Hsp90 protein in assay buffer (100 mM Tris-HCl, Ph7.4, 20 mM KCl, 6 mM MgCl₂) was mixed with ATP alone (negative control) or in the presence of Geldanamycin (a positive control) or Compound 108 in a 96-well plate. Malachite green reagent was
20 added to the reaction. The mixtures were incubated at 37°C for 4 hours and sodium citrate buffer (34% w/v sodium citrate) was added to the reaction. The plate was read by an ELISA reader with an absorbance at 620 nm.

As can be seen in Figure 1, 40 μM of geldanamycin, a natural product known to inhibit Hsp90 activity, the ATPase activity of Hsp90 was only slightly higher than
25 background. 40 μM Compound 108 showed an even greater inhibition of ATPase activity of Hsp90 than geldanamycin, and even at 4 μM Compound 108 showed significant inhibition of ATPase activity of Hsp90 protein.

Example 169: Degradation of Hsp90 Client Proteins via Inhibition of Hsp90
30 **Activity****A. Cells and Cell Culture**

Human high-Her2 breast carcinoma BT474 (HTB-20), SK-BR-3 (HTB-30) and MCF-7 breast carcinoma (HTB-22) from American Type Culture Collection, VA, USA were grown in Dulbecco's modified Eagle's medium with 4 mM L-glutamine and antibiotics (100IU/ml penicillin and 100 ug/ml streptomycin; GibcoBRL). To obtain exponential cell growth, cells were trypsinized, counted and seeded at a cell density of 0.5×10^6 cells /ml regularly, every 3 days. All experiments were performed on day 1 after cell passage.

10 B. Degradation of Her2 in Cells after Treatment with a Compound of the Invention

BT-474 cells were treated with 0.5 μ M, 2 μ M, or 5 μ M of 17AAG (a positive control) or 0.5 μ M, 2 μ M, or 5 μ M of Compound 108 or Compound 49 overnight in DMEM medium. After treatment, each cytoplasmic sample was prepared from 1×10^6 cells by incubation of cell lysis buffer (#9803, Cell Signaling Technology) on ice for 10 minutes. The resulting supernatant used as the cytosol fractions were dissolved with sample buffer for SDS-PAGE and run on a SDS-PAGE gel, blotted onto a nitrocellulose membrane by using semi-dry transfer. Non-specific binding to nitrocellulose was blocked with 5% skim milk in TBS with 0.5% Tween at room temperature for 1 hour, then probed with anti-Her2/ErB2 mAb (rabbit IgG, #2242, Cell Signaling) and anti-Tubulin (T9026, Sigma) as housekeeping control protein. HRP-conjugated goat anti-rabbit IgG (H+L) and HRP-conjugated horse anti-mouse IgG (H+L) were used as secondary Ab (#7074, #7076, Cell Signaling) and LumiGLO reagent, 20x Peroxide (#7003, Cell Signaling) was used for visualization.

As can be seen from Figure 2, Her2, an Hsp90 client protein, is almost completely degraded when cells are treated with 5 μ M of Compound 108 and partially degraded when cells are treated with 2 μ M and 0.5 μ M of Compound 108. Compound 49 which is even more active than Compound 108 causes complete degradation of Her2 when cells are treated with 2 μ M and 5 μ M and causes partial degraded when cells are treated with 0.5 μ M. 17AAG is a known Hsp90 inhibitor and is used as a positive control.

C. Fluorescent Staining of Her2 on the Surface of Cells Treated with a Compound of the Invention

After treatment with a compound of the invention, cells were washed twice with 1xPBS/1%FBS, and then stained with anti-Her2- FITC (#340553, BD) for 30 min at 4°C. Cells were then washed three times in FACS buffer before the fixation in 0.5 ml 1% paraformaldehyde. Data was acquired on a FACSCalibur system. Isotype-matched controls were used to establish the non-specific staining of samples and to set the fluorescent markers. A total 10,000 events were recorded from each sample. Data were analysed by using CellQuest software (BD Biosciences). The IC₅₀ range for Hsp90 inhibition by compounds of the invention are listed below in Table 2.

Table 2: IC₅₀ range of compounds of the invention for inhibition of Hsp90

IC ₅₀ Range	Compound Number
< 3μM	8, 13, 39, 49, 63, 76, 77, 79, 87, 88, 95, 96, 100, 103, 177, 178, 185, 188, 189, 195, 197, 198, 201, 202, 203, 204, 205, 206, 207, 208, 209, 211, 212, 213, 214, 215, 216, 218, 219, 220, 221, 222, 223
3μM to 10μM	2, 5, 6, 7, 9, 14, 27, 28, 34, 36, 38, 42, 48, 64, 70, 93, 97, 108, 122, 183, 184, 194, 196, 217
10μM to 100μM	21, 22, 30, 51, 59, 60, 61, 62, 94, 98, 99, 102, 104, 123, 181, 182, 186, 187, 191, 192, 193, 199, 210

D. Apoptosis analysis

After treatment with the compounds of the invention, cells were washed once with 1xPBS/1%FBS, and then stained in binding buffer with FITC-conjugated Annexin V and Propidium iodide (PI) (all obtained from BD Biosciences) for 30 min at 4°C. Flow cytometric analysis was performed with FACSCalibur (BD Biosciences) and a total 10,000 events were recorded from each sample. Data were analyzed by using CellQuest software (BD Biosciences). The relative fluorescence was calculated after subtraction of the fluorescence of control.

E. Degradation of c-Kit in Cells after Treatment with a Compound of the Invention

Two leukemia cell lines, HEL92.1.7 and Kasumi-1, were used for testing c-kit degradation induced by Hsp90 inhibitors of the invention. The cells (3×10^5 per well) were treated with 17AAG (0.5 μM), Compound 188 or Compound 221 for about 18 h (see Figs. 3 and 4 for concentrations). The cells were collected and centrifuged (SORVALL RT 6000D) at 1200 rpm for 5 min. The supernatants were discarded, and the cells were washed one time with 1X PBS. After centrifugation the cells were stained with FITC conjugated c-kit antibody (MBL International, Cat# K0105-4) in 100 ml 1X PBS at 4°C for 1 h. The samples were read and analyzed with FACSCalibur flow cytometer (Becton Dickinson).

c-Kit, a tyrosine kinase receptor and one of the Hsp90 client proteins, was selected and used in a FACS-based degradation assay. The results of the assay showed that Compound 188 and Compound 221, induced c-kit degradation at 0.5 and 0.05 μM in a dose-dependent manner. Surprisingly, 17-AAG, which is a potent Hsp90 inhibitor and is in phase 2 clinical trials, could not induce c-kit degradation at 0.5 μM in two leukemia cell lines, HEL92.1.7 (see Fig. 3) and Kasumi-1 (see Fig. 4). Since the compounds of the invention cause c-kit degradation more efficiently than other Hsp90 inhibitors, the compounds of the invention are expected to be more effective in the treatment of c-kit associated tumors, such as leukemias, mast cell tumors, small cell lung cancer, testicular cancer, some cancers of the gastrointestinal tract (including GIST), and some central nervous system.

The results of the FACS analysis were confirmed with Western blot analysis (see Fig. 5). In Kasumi-1 cells (myelogenous leukemia), Compound 221 (100 nM and 400 nM) induced the degradation of c-Kit. In contrast, 17-AAG had no effect of c-Kit protein levels.

F. Degradation of c-Met in Cells after Treatment with a Compound of the Invention

We examined the ability of the Hsp90 inhibitors of the invention to induce the degradation of c-Met, an Hsp90 client protein that is expressed at high levels in several types of non-small cell lung cancer. NCI-H1993 (ATCC, cat# CRL-5909)

were seeded in 6-well plates at 5×10^5 cells/well. The cells were treated with 17AAG (100 nM or 400 nM) or Compound 221 (100 nM or 400 nM), and cell lysis was prepared 24 h after treatment. Equal amount of proteins were used for Western blot analysis. The compounds of the invention potently induced degradation of c-Met in this cell line due to inhibition of Hsp90 (see Fig. 6).

Example 170: Compound 49 Displays Anti-tumor Activity Against the Human Tumor Cell Line MDA-MB-435S in a nude Mouse Xenograft Model

The human tumor cell line, MDA-MB-435S (ATCC #HTB-129; G. Ellison, et al., *Mol. Pathol.* 55:294-299, 2002), was obtained from the American Type Culture Collection (Manassus, Virginia, USA). The cell line was cultured in growth media prepared from 50% Dulbecco's Modified Eagle Medium (high glucose), 50% RPMI Media 1640, 10% fetal bovine serum (FBS), 1% 100X L-glutamine, 1% 100X Penicillin-Streptomycin, 1% 100X sodium pyruvate and 1% 100X MEM non-essential amino acids. FBS was obtained from Sigma-Aldrich Corp. (St. Louis, Missouri, USA), and all other reagents were obtained from Invitrogen Corp. (Carlsbad, California, USA). Approximately $4-5 \times 10^6$ cells that had been cryopreserved in liquid nitrogen were rapidly thawed at 37°C and transferred to a 175 cm^2 tissue culture flask containing 50 ml of growth media and then incubated at 37°C in a 5% CO_2 incubator. The growth media was replaced every 2-3 days until the flask became 90% confluent, typically in 5-7 days. To passage and expand the cell line, a 90% confluent flask was washed with 10 ml of room temperature phosphate buffered saline (PBS) and the cells were disassociated by adding 5 ml 1X Trypsin-EDTA (Invitrogen) and incubating at 37°C until the cells detached from the surface of the flask. To inactivate the trypsin, 5 ml of growth media was added and then the contents of the flask were centrifuged to pellet the cells. The supernatant was aspirated and the cell pellet was resuspended in 10 ml of growth media and the cell number determined using a hemocytometer. Approximately $1-3 \times 10^6$ cells per flask were seeded into 175 cm^2 flasks containing 50 ml of growth media and incubated at 37°C in a 5% CO_2 incubator.

When the flasks reached 90% confluence, the above passaging process was repeated until sufficient cells had been obtained for implantation into mice.

Six to eight week old, female Crl:CD-1-*nu*BR (*nude*) mice were obtained from Charles River Laboratories (Wilmington, Massachusetts, USA). Animals were housed 4-5/cage in micro-isolators, with a 12hr/12hr light/dark cycle, acclimated for at least 1 week prior to use and fed normal laboratory chow *ad libitum*. Studies were conducted on animals between 7 and 12 weeks of age at implantation. To implant tumor cells into *nude* mice, the cells were trypsinized as above, washed in PBS and resuspended at a concentration of 50×10^6 cells/ml in PBS. Using a 27 gauge needle and 1 cc syringe, 0.1 ml of the cell suspension was injected into the corpus adiposum of *nude* mice. The corpus adiposum is a fat body located in the ventral abdominal viscera in the right quadrant of the abdomen at the juncture of the os coxae (pelvic bone) and the os femoris (femur). Tumors were then permitted to develop *in vivo* until they reached approximately 150 mm³ in volume, which typically required 2-3 weeks following implantation. Tumor volumes (V) were calculated by caliper measurement of the width (W), length (L) and thickness (T) of tumors using the following formula: $V = 0.5326 \times (L \times W \times T)$. Animals were randomized into treatment groups so that the average tumor volumes of each group were similar at the start of dosing.

Stock solutions of test compounds were prepared by dissolving the appropriate amounts of each compound in dimethyl sulfoxide (DMSO) by sonication in an ultrasonic water bath. Stock solutions were prepared at the start of the study, stored at -20°C and diluted fresh each day for dosing. A solution of 20% Cremophore RH40 (polyoxyl 40 hydrogenated castor oil; BASF Corp., Aktiengesellschaft, Ludwigshafen, Germany) in 80% D5W (5% dextrose in water; Abbott Laboratories, North Chicago, Illinois, USA) was also prepared by first heating 100% Cremophore RH40 at 50-60°C until liquefied and clear, diluting 1:5 with 100% D5W, reheating again until clear and then mixing well. This solution was stored at room temperature for up to 3 months prior to use. To prepare formulations for daily dosing, DMSO stock solutions were diluted 1:10 with 20% Cremophore RH40. The final formulation for dosing contained 10% DMSO, 18% Cremophore RH40, 3.6% dextrose and 68.4% water and the appropriate amount of

test article. Animals were intraperitoneal (IP) injected with this solution at 10 ml per kg body weight on a schedule of 5 days per week (Monday thru Friday, with no dosing on Saturday and Sunday) for 3 weeks.

As shown in Figure 7, treatment with 300 mg/kg body weight of Compound 49 decreased the growth rate of MDA-MB-435S cells in *nude* mice to a greater extent than did a dose of 100 mg/kg body weight of the Hsp90 inhibitor 17-AAG. This effect was not associated with significant toxicity, as shown by the lack of an effect on body weights (Figure 8).

Example 171: Compound 188 Displays Anti-tumor Activity Against Human Tumor Cells in a *nude* Mouse Xenograft Model

The human squamous non-small cell lung cancer cell line, RERF-LC-AI (RCB0444; S. Kyoizumi, et al., *Cancer. Res.* 45:3274-3281, 1985), was obtained from the Riken Cell Bank (Tsukuba, Ibaraki, Japan). The cell line was cultured in growth media prepared from 50% Dulbecco's Modified Eagle Medium (high glucose), 50% RPMI Media 1640, 10% fetal bovine serum (FBS), 1% 100X L-glutamine, 1% 100X penicillin-streptomycin, 1% 100X sodium pyruvate and 1% 100X MEM non-essential amino acids. FBS was obtained from American Type Culture Collection (Manassas, Virginia, USA) and all other reagents were obtained from Invitrogen Corp. (Carlsbad, California, USA). Approximately 4-5 x 10⁶ cells that had been cryopreserved in liquid nitrogen were rapidly thawed at 37°C and transferred to a 175 cm² tissue culture flask containing 50 ml of growth media and then incubated at 37°C in a 5% CO₂ incubator.

The growth media was replaced every 2-3 days until the flask became 90% confluent, typically in 5-7 days. To passage and expand the cell line, a 90% confluent flask was washed with 10 ml of room temperature phosphate buffered saline (PBS) and the cells were disassociated by adding 5 ml 1X trypsin-EDTA (Invitrogen) and incubating at 37°C until the cells detached from the surface of the flask. To inactivate the trypsin, 5 ml of growth media was added and then the contents of the flask were centrifuged to pellet the cells. The supernatant was aspirated and the cell pellet was resuspended in 10 ml of growth media and the cell number determined using a hemocytometer. Approximately 1-3 x 10⁶ cells per

flask were seeded into 175 cm² flasks containing 50 ml of growth media and incubated at 37°C in a 5% CO₂ incubator. When the flasks reached 90% confluence, the above passaging process was repeated until sufficient cells had been obtained for implantation into mice.

- 5 Seven to eight week old, female Crl:CD-1-*nu*BR (*nude*) mice were obtained from Charles River Laboratories (Wilmington, Massachusetts, USA). Animals were housed 4-5/cage in micro-isolators, with a 12hr/12hr light/dark cycle, acclimated for at least 1 week prior to use and fed normal laboratory chow *ad libitum*. Studies were conducted on animals between 8 and 12 weeks of age at
- 10 implantation. To implant RERF-LC-AI tumor cells into nude mice, the cells were trypsinized as above, washed in PBS and resuspended at a concentration of 50 x 10(6) cells/ml in 50% non-supplemented RPMI Media 1640 and 50% Matrigel Basement Membrane Matrix (#354234; BD Biosciences; Bedford, Massachusetts, USA). Using a 27 gauge needle and 1 cc syringe, 0.1 ml of the cell suspension was
- 15 injected subcutaneously into the flank of each *nude* mouse. Tumor volumes (V) were calculated by caliper measurement of the width (W), length (L) and thickness (T) of tumors using the following formula: $V = 0.5236 \times (L \times W \times T)$.

- In vivo* passaged RERF-LC-AI tumor cells (RERF-LC-AI^{IVP}) were isolated to improve the rate of tumor implantation relative to the parental cell line in *nude*
- 20 mice. RERF-LC-AI tumors were permitted to develop *in vivo* until they reached approximately 250 mm³ in volume, which required approximately 3 weeks following implantation. Mice were euthanized via CO₂ asphyxiation and their exteriors sterilized with 70% ethanol in a laminar flow hood. Using sterile technique, tumors were excised and diced in 50 ml PBS using a scalpel blade. A
- 25 single cell suspension was prepared using a 55 ml Wheaton Safe-Grind tissue grinder (catalog #62400-358; VWR International, West Chester, Pennsylvania, USA) by plunging the pestle up and down 4-5 times without twisting. The suspension was strained through a 70 μM nylon cell strainer and then centrifuged to pellet the cells. The resulting pellet was resuspended in 0.1 M NH₄Cl to lyse
- 30 contaminating red blood cells and then immediately centrifuged to pellet the cells. The cell pellet was resuspended in growth media and seeded into 175 cm² flasks

containing 50 ml of growth media at 1-3 tumors/flask or approximately 10×10^6 cells/flask. After overnight incubation at 37°C in a 5% CO₂ incubator, non-adherent cells were removed by rinsing two times with PBS and then the cultures were fed with fresh growth media. When the flasks reached 90% confluence, the
5 above passaging process was repeated until sufficient cells had been obtained for implantation into mice.

RERF-LC-AI^{IVP} cells were then implanted as above and tumors were permitted to develop *in vivo* until the majority reached an average of 100-200 mm³ in tumor volume, which typically required 2-3 weeks following implantation.
10 Animals with oblong or very small or large tumors were discarded, and only animals carrying tumors that displayed consistent growth rates were selected for studies. Animals were randomized into treatment groups so that the average tumor volumes of each group were similar at the start of dosing.

The HSP90 inhibitor, 17-allylamino-17-demethoxygeldanamycin (17-AAG), was employed as a positive control (Albany Molecular Research, Albany, New York, USA). Stock solutions of test articles were prepared by dissolving the appropriate amounts of each compound in dimethyl sulfoxide (DMSO) by
15 sonication in an ultrasonic water bath. Stock solutions were prepared weekly, stored at -20°C and diluted fresh each day for dosing. A solution of 20% Cremophore RH40 (polyoxyl 40 hydrogenated castor oil; BASF Corp.,
20 Aktiengesellschaft, Ludwigshafen, Germany) in 80% D5W (5% dextrose in water; Abbott Laboratories, North Chicago, Illinois, USA) was also prepared by first heating 100% Cremophore RH40 at 50-60°C until liquefied and clear, diluting 1:5 with 100% D5W, reheating again until clear and then mixing well. This solution
25 was stored at room temperature for up to 3 months prior to use. To prepare formulations for daily dosing, DMSO stock solutions were diluted 1:10 with 20% Cremophore RH40. The final formulation for dosing contained 10% DMSO, 18% Cremophore RH40, 3.6% dextrose, 68.4% water and the appropriate amount of test
30 article. Animals were intraperitoneally (i.p.) injected with this solution at 10 ml per kg body weight on a schedule of 5 days per week (Monday, Tuesday, Wednesday, Thursday and Friday, with no dosing on Saturday and Sunday) for a total of 15 doses.

As shown in Figure 9, treatment with 200 mg/kg body weight of Compound 188 decreased the growth rate of RERF-LC-AI^{IVP} human lung tumor cells in *nude* mice, as did a dose of 75 mg/kg body weight of 17-AAG (an unrelated HSP90 inhibitor). This effect was not associated with overt toxicity, as shown by the
5 minimal effect on body weights depicted in Figure 10.

All publications, patent applications, patents, and other documents cited herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods,
10 and examples are illustrative only and not intended to be limiting.


While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled
15 in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

CLAIMS

What is claimed is:

1. A compound represented by the following structural formula:

5



or a tautomer, pharmaceutically acceptable salt, solvate, clathrate, or a prodrug thereof, wherein:

10 ring A is an aryl or a heteroaryl, wherein the aryl or the heteroaryl are optionally further substituted with one or more substituents in addition to R₃;

R₁ and R₃ are, independently, -OH, -SH, -NR₇H, -OR₂₆, -SR₂₆, -NHR₂₆, -O(CH₂)_mOH, -O(CH₂)_mSH, -O(CH₂)_mNR₇H, -S(CH₂)_mOH, -S(CH₂)_mSH, -S(CH₂)_mNR₇H, -OC(O)NR₁₀R₁₁, -SC(O)NR₁₀R₁₁, -NR₇C(O)NR₁₀R₁₁, -OC(O)R₇, -SC(O)R₇, -NR₇C(O)R₇, -OC(O)OR₇, -SC(O)OR₇, -NR₇C(O)OR₇, -OCH₂C(O)R₇, -SCH₂C(O)R₇, -NR₇CH₂C(O)R₇, -OCH₂C(O)OR₇, -SCH₂C(O)OR₇, -NR₇CH₂C(O)OR₇, -OCH₂C(O)NR₁₀R₁₁, -SCH₂C(O)NR₁₀R₁₁, -NR₇CH₂C(O)NR₁₀R₁₁, -OS(O)_pR₇, -SS(O)_pR₇, -S(O)_pOR₇, -NR₇S(O)_pR₇, -OS(O)_pNR₁₀R₁₁, -SS(O)_pNR₁₀R₁₁, -NR₇S(O)_pNR₁₀R₁₁, -OS(O)_pOR₇, -SS(O)_pOR₇, -NR₇S(O)_pOR₇, -OC(S)R₇, -SC(S)R₇, -NR₇C(S)R₇, -OC(S)OR₇, -SC(S)OR₇, -NR₇C(S)OR₇, -OC(S)NR₁₀R₁₁, -SC(S)NR₁₀R₁₁, -NR₇C(S)NR₁₀R₁₁, -OC(NR₈)R₇, -SC(NR₈)R₇, -NR₇C(NR₈)R₇, -OC(NR₈)OR₇, -SC(NR₈)OR₇, -NR₇C(NR₈)OR₇, -OC(NR₈)NR₁₀R₁₁, -SC(NR₈)NR₁₀R₁₁, -NR₇C(NR₈)NR₁₀R₁₁, -OP(O)(OR₇)₂, or -SP(O)(OR₇)₂;

25

R₅ is an optionally substituted heteroaryl or an optionally substituted 8 to 14 membered aryl;

5 R_7 and R_8 , for each occurrence, are, independently, -H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

10 R_{10} and R_{11} , for each occurrence, are independently -H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R_{10} and R_{11} , taken together with the nitrogen to which they are attached, form an optionally substituted heterocyclyl or an optionally substituted heteroaryl;

15 R_{26} is a C1-C6 alkyl;


p , for each occurrence, is, independently, 0, 1 or 2; and

m , for each occurrence, is independently, 1, 2, 3, or 4;

provided that ring A is not a substituted [1,2,3]triazole; and

20 provided that the compound is not 3-(2,4-dihydroxy-phenyl)-4-(7-naphthalen-1-yl)-5-mercapto-triazole.

2. The compound of Claim 1, wherein R_5 is represented by the following formula:




25 wherein:

R_9 , for each occurrence, is independently a substituent selected from the group consisting of an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally

substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, hydroxyalkyl, alkoxyalkyl, halo, cyano, nitro, guanadino, a haloalkyl, a heteroalkyl, $-NR_{10}R_{11}$, $-OR_7$, $-C(O)R_7$, $-C(O)OR_7$, $-OC(O)R_7$, $-C(O)NR_{10}R_{11}$, $-NR_8C(O)R_7$, $-SR_7$, $-S(O)_pR_7$, $-OS(O)_pR_7$, $-S(O)_pOR_7$, $-NR_8S(O)_pR_7$, or $-S(O)_pNR_{10}R_{11}$, $-S(O)_pOR_7$, $-OP(O)(OR_7)_2$, or $-SP(O)(OR_7)_2$;

or two R_9 groups taken together with the carbon atoms to which they are attached form a fused ring; and
 m is zero or an integer from 1 to 7.


3. The compound of Claim 2, wherein R_5 is represented by one of the following formulas:



wherein:

q is zero or an integer from 1 to 7; and
 u is zero or an integer from 1 to 8.

4. The compound of Claim 1, wherein R_5 is represented by the following structural formula:









wherein:




R₃₃ is a halo, lower alkyl, a lower alkoxy, a lower haloalkyl, a lower haloalkoxy, and lower alkyl sulfanyl;

5 R₃₄ is H, a lower alkyl, or a lower alkylcarbonyl; and


Ring B and Ring C are optionally substituted with one or more substituents.

10 5. The compound of Claim 1, wherein R₅ is selected from the group consisting of:






5



and



10

wherein:

X_6 , for each occurrence, is independently CH, CR₉, N, N(O), N⁺(R₁₇), provided that at least three X_6 groups are independently selected from CH and CR₉;

5 X_7 , for each occurrence, is independently CH, CR₉, N, N(O), N⁺(R₁₇), provided that at least three X_7 groups are independently selected from CH and CR₉;

X_8 , for each occurrence, is independently CH₂, CHR₉, C(R₉)₂, S, S(O)_p, NR₇, or NR₁₇;

X_9 , for each occurrence, is independently N or CH;

10 X_{10} , for each occurrence, is independently CH, CR₉, N, N(O), N⁺(R₁₇), provided that at least one X_{10} is selected from CH and CR₉;

R_9 , for each occurrence, is independently a substituent selected from the group consisting of an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, hydroxyalkyl, alkoxyalkyl, halo, cyano, nitro, guanadino, a haloalkyl, a heteroalkyl, -NR₁₀R₁₁, -OR₇, -C(O)R₇,
15 -C(O)OR₇, -OC(O)R₇, -C(O)NR₁₀R₁₁, -NR₈C(O)R₇, -SR₇, -S(O)_pR₇,
20 -OS(O)_pR₇, -S(O)_pOR₇, -NR₈S(O)_pR₇, or -S(O)_pNR₁₀R₁₁, -S(O)_pOR₇,
-OP(O)(OR₇)₂, or -SP(O)(OR₇)₂, -S(O)_pOR₇, -OP(O)(OR₇)₂, or
-SP(O)(OR₇)₂;


25 or two R_9 groups taken together with the carbon atoms to which they are attached form a fused ring; and

R_{17} , for each occurrence, is independently -H, an alkyl, an aralkyl, -C(O)R₇, -C(O)OR₇, or -C(O)NR₁₀R₁₁.

6. The compound of Claim 5, wherein R_5 is an optionally substituted indolyl,
30 an optionally substituted benzoimidazolyl, an optionally substituted indazolyl, an optionally substituted 3*H*-indazolyl, an optionally substituted indoliziny, an optionally substituted quinoliny, an optionally substituted

isoquinoliny], an optionally substituted benzoxazolyl, an optionally substituted benzo[1,3]dioxolyl, an optionally substituted benzofuryl, an optionally substituted benzothiazolyl, an optionally substituted benzo[d]isoxazolyl, an optionally substituted benzo[d]isothiazolyl, an optionally substituted thiazolo[4,5-c]pyridinyl, an optionally substituted thiazolo[5,4-c]pyridinyl, an optionally substituted thiazolo[4,5-b]pyridinyl, an optionally substituted thiazolo[5,4-b]pyridinyl, an optionally substituted oxazolo[4,5-c]pyridinyl, an optionally substituted oxazolo[5,4-c]pyridinyl, an optionally substituted oxazolo[4,5-b]pyridinyl, an optionally substituted oxazolo[5,4-b]pyridinyl, an optionally substituted imidazopyridinyl, an optionally substituted benzothiadiazolyl, benzoxadiazolyl, an optionally substituted benzotriazolyl, an optionally substituted tetrahydroindolyl, an optionally substituted azaindolyl, an optionally substituted quinazolinyl, an optionally substituted purinyl, an optionally substituted imidazo[4,5-a]pyridinyl, an optionally substituted imidazo[1,2-a]pyridinyl, an optionally substituted 3*H*-imidazo[4,5-b]pyridinyl, an optionally substituted 1*H*-imidazo[4,5-b]pyridinyl, an optionally substituted 1*H*-imidazo[4,5-c]pyridinyl, an optionally substituted 3*H*-imidazo[4,5-c]pyridinyl, an optionally substituted pyridopyrdazinyl, and optionally substituted pyridopyrimidinyl, an optionally substituted pyrrolo[2,3]pyrimidyl, an optionally substituted pyrazolo[3,4]pyrimidyl an optionally substituted cyclopentaimidazolyl, an optionally substituted cyclopentatriazolyl, an optionally substituted pyrrolopyrazolyl, an optionally substituted pyrroloimidazolyl, an optionally substituted pyrrolotriazolyl, or an optionally substituted benzo(b)thienyl.

7. The compound of Claim 1, wherein R₅ is selected from the group consisting of:



wherein:

X_{11} , for each occurrence, is independently CH, CR₉, N, N(O), or N⁺(R₁₇), provided that at least one X_{11} is N, N(O), or N⁺(R₁₇) and at least two X_{11} groups are independently selected from CH and CR₉;

X_{12} , for each occurrence, is independently CH, CR₉, N, N(O), N⁺(R₁₇), provided that at least one X_{12} group is independently selected from CH and CR₉;


X_{13} , for each occurrence, is independently O, S, S(O)_p, NR₇, or NR₁₇;

R₉, for each occurrence, is independently a substituent selected from the group consisting of an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, halo, cyano, nitro, guanadino, a hydroxyalkyl, alkoxyalkyl, haloalkyl, a heteroalkyl, -NR₁₀R₁₁, -OR₇, -C(O)R₇, -C(O)OR₇, -OC(O)R₇, -C(O)NR₁₀R₁₁, -NR₈C(O)R₇, -SR₇, -S(O)_pR₇, -OS(O)_pR₇, -S(O)_pOR₇, -NR₈S(O)_pR₇, or -S(O)_pNR₁₀R₁₁, -S(O)_pOR₇, -OP(O)(OR₇)₂, or -SP(O)(OR₇)₂, -S(O)_pOR₇, -OP(O)(OR₇)₂, or -SP(O)(OR₇)₂;

or two R₉ groups taken together with the carbon atoms to which they are attached form a fused ring; and

R₁₇, for each occurrence, is independently an alkyl or an aralkyl.

8. The compound of Claim 1, wherein the compound is represented by the following structural formula:



wherein:

5 R_6 , for each occurrence, is independently an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteroaralkyl, halo, cyano, nitro, guanadino, a
 10 haloalkyl, a heteroalkyl, alkoxy, haloalkoxy, $-NR_{10}R_{11}$, $-OR_7$, $-C(O)R_7$, $-C(O)OR_7$, $-C(S)R_7$, $-C(O)SR_7$, $-C(S)SR_7$, $-C(S)OR_7$, $-C(S)NR_{10}R_{11}$, $-C(NR_8)OR_7$, $-C(NR_8)R_7$, $-C(NR_8)NR_{10}R_{11}$, $-C(NR_8)SR_7$, $-OC(O)R_7$, $-OC(O)OR_7$, $-OC(S)OR_7$, $-OC(NR_8)OR_7$, $-SC(O)R_7$, $-SC(O)OR_7$, $-SC(NR_8)OR_7$, $-OC(S)R_7$, $-SC(S)R_7$, $-SC(S)OR_7$, $-OC(O)NR_{10}R_{11}$,
 15 $-OC(S)NR_{10}R_{11}$, $-OC(NR_8)NR_{10}R_{11}$, $-SC(O)NR_{10}R_{11}$, $-SC(NR_8)NR_{10}R_{11}$, $-SC(S)NR_{10}R_{11}$, $-OC(NR_8)R_7$, $-SC(NR_8)R_7$, $-C(O)NR_{10}R_{11}$, $-NR_8C(O)R_7$, $-NR_7C(S)R_7$, $-NR_7C(S)OR_7$, $-NR_7C(NR_8)R_7$, $-NR_7C(O)OR_7$, $-NR_7C(NR_8)OR_7$, $-NR_7C(O)NR_{10}R_{11}$, $-NR_7C(S)NR_{10}R_{11}$, $-NR_7C(NR_8)NR_{10}R_{11}$, $-SR_7$, $-S(O)_pR_7$, $-OS(O)_pR_7$, $-OS(O)_pOR_7$,
 20 $-OS(O)_pNR_{10}R_{11}$, $-S(O)_pOR_7$, $-NR_8S(O)_pR_7$, $-NR_7S(O)_pNR_{10}R_{11}$, $-NR_7S(O)_pOR_7$, $-S(O)_pNR_{10}R_{11}$, $-SS(O)_pR_7$, $-SS(O)_pOR_7$, $-SS(O)_pNR_{10}R_{11}$, $-OP(O)(OR_7)_2$, or $-SP(O)(OR_7)_2$; and

n is zero or an integer from 1 to 4.

- 25 9. The compound of Claim 8, wherein R_5 is moiety R_2 , and wherein R_2 is a substituted phenyl, wherein the phenyl group is substituted with:
 i) one substituent selected from nitro, cyano, a haloalkoxy, an optionally substituted alkenyl, an optionally substituted alkynyl, an

- optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, hydroxylalkyl, alkoxyalkyl, guanadino, $-NR_{10}R_{11}$, $-O-R_{20}$, $-C(O)R_7$, $-C(O)OR_{20}$, $-OC(O)R_7$, $-C(O)NR_{10}R_{11}$, $-NR_8C(O)R_7$, $-SR_7$, $-S(O)_pR_7$, $-OS(O)_pR_7$, $-S(O)_pOR_7$, $-NR_8S(O)_pR_7$, $-S(O)_pNR_{10}R_{11}$, $-S(O)_pOR_7$, $-OP(O)(OR_7)_2$, or $-SP(O)(OR_7)_2$; or
- ii) two to five substituents selected from the group consisting of an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, hydroxylalkyl, alkoxyalkyl, $-F$, $-Br$, $-I$, cyano, nitro, guanadino, a haloalkyl, a heteroalkyl, $-NR_{10}R_{11}$, $-OR_7$, $-C(O)R_7$, $-C(O)OR_7$, $-OC(O)R_7$, $-C(O)NR_{10}R_{11}$, $-NR_8C(O)R_7$, $-SR_7$, $-S(O)_pR_7$, $-OS(O)_pR_7$, $-S(O)_pOR_7$, $-NR_8S(O)_pR_7$, $-S(O)_pNR_{10}R_{11}$, $-OP(O)(OR_7)_2$, or $-SP(O)(OR_7)_2$, $-S(O)_pOR_7$.
10. A compound of Claim 1, wherein R_5 is a moiety R_{18} , and further wherein R_{18} is an optionally substituted cycloalkyl, and optionally substituted cycloalkenyl, or a substituted alkyl, wherein the alkyl group is substituted with one or more substituents independently selected from the group consisting of an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, halo, cyano, nitro, guanadino, a haloalkyl, $-NR_{10}R_{11}$, $-OR_7$, $-C(O)R_7$, $-C(O)OR_7$, $-OC(O)R_7$, $-C(O)NR_{10}R_{11}$, $-NR_8C(O)R_7$, $-SR_7$, $-S(O)_pR_7$, $-OS(O)_pR_7$, $-S(O)_pOR_7$, $-NR_8S(O)_pR_7$, or $-S(O)_pNR_{10}R_{11}$, $-S(O)_pOR_7$, $-OP(O)(OR_7)_2$, or $-SP(O)(OR_7)_2$; and
- m , for each occurrence, is independently, 1, 2, 3, or 4;


provided that R_{18} is not cyclohexyl.

11. The compound of Claim 10, wherein R_{18} is an optionally substituted cycloalkyl or an optionally substituted cycloalkenyl.

5

12. The compound of Claim 10, wherein R_{18} is a substituted alkyl.

13. The compound of Claim 8, wherein the compound is represented by the following structural formula:



10

wherein:

- R_{25} is an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteroaralkyl, halo, cyano, nitro, guanadino, a haloalkyl, a heteroalkyl, alkoxy, haloalkoxy, $-NR_{10}R_{11}$, $-OR_7$, $-C(O)R_7$, $-C(O)OR_7$, $-C(S)R_7$, $-C(O)SR_7$, $-C(S)SR_7$, $-C(S)OR_7$, $-C(S)NR_{10}R_{11}$, $-C(NR_8)OR_7$, $-C(NR_8)R_7$, $-C(NR_8)NR_{10}R_{11}$, $-C(NR_8)SR_7$, $-OC(O)R_7$, $-OC(O)OR_7$, $-OC(S)OR_7$, $-OC(NR_8)OR_7$, $-SC(O)R_7$, $-SC(O)OR_7$, $-SC(NR_8)OR_7$, $-OC(S)R_7$, $-SC(S)R_7$, $-SC(S)OR_7$, $-OC(O)NR_{10}R_{11}$, $-OC(S)NR_{10}R_{11}$, $-OC(NR_8)NR_{10}R_{11}$, $-SC(O)NR_{10}R_{11}$, $-SC(NR_8)NR_{10}R_{11}$, $-SC(S)NR_{10}R_{11}$, $-OC(NR_8)R_7$, $-SC(NR_8)R_7$, $-C(O)NR_{10}R_{11}$, $-NR_8C(O)R_7$, $-NR_7C(S)R_7$, $-NR_7C(S)OR_7$, $-NR_7C(NR_8)R_7$, $-NR_7C(O)OR_7$, $-NR_7C(NR_8)OR_7$, $-NR_7C(O)NR_{10}R_{11}$, $-NR_7C(S)NR_{10}R_{11}$, $-NR_7C(NR_8)NR_{10}R_{11}$, $-SR_7$, $-S(O)_pR_7$, $-OS(O)_pR_7$,
- 15
- 20
- 25

-OS(O)_pOR₇, -OS(O)_pNR₁₀R₁₁, -S(O)_pOR₇, -NR₈S(O)_pR₇,
 -NR₇S(O)_pNR₁₀R₁₁, -NR₇S(O)_pOR₇, -S(O)_pNR₁₀R₁₁, -SS(O)_pR₇,
 -SS(O)_pOR₇, -SS(O)_pNR₁₀R₁₁, -OP(O)(OR₇)₂, or -SP(O)(OR₇)₂;


k is 1, 2, 3, or 4; and

5 r is zero or an integer from 1 to 3.

14. The compound of Claim 13 wherein R₁, R₃ and R₂₅ are each independently
 -OH, -SH, -NHR₇, -OC(O)NR₁₀R₁₁, -SC(O)NR₁₀R₁₁, -OC(O)R₇, -SC(O)R₇,
 -OC(O)OR₇, -SC(O)OR₇, -OS(O)_pR₇, -S(O)_pOR₇, -SS(O)_pR₇, -OS(O)_pOR₇,
 10 -SS(O)_pOR₇, -OC(S)R₇, -SC(S)R₇, -OC(S)OR₇, -SC(S)OR₇, -OC(S)NR₁₀R₁₁,
 -SC(S)NR₁₀R₁₁, -OC(NR₈)R₇, -SC(NR₈)R₇, -OC(NR₈)OR₇, -SC(NR₈)OR₇,
 -OP(O)(OR₇)₂ or -SP(O)(OR₇)₂.

15. The compound of Claim 3, wherein R₁ and R₃ are each, independently, -OH,
 15 -SH, or -NHR₇.

16. The compound of Claim 15, wherein the compound is represented by the
 following structural formula:



20 wherein R₆ is an optionally substituted alkyl, an optionally
 substituted alkenyl, an optionally substituted alkynyl, cyano, halo, nitro, an
 optionally substituted cycloalkyl, haloalkyl, an optionally substituted
 heterocyclyl, an optionally substituted aryl, an optionally substituted
 heteroaryl, an optionally substituted aralkyl, an optionally substituted
 25 heteroaralkyl, -OR₇, -SR₇, -NR₁₀R₁₁, -OC(O)NR₁₀R₁₁, -SC(O)NR₁₀R₁₁,
 -NR₇C(O)NR₁₀R₁₁, -OC(O)R₇, -SC(O)R₇, -NR₇C(O)R₇, -OC(O)OR₇,
 -SC(O)OR₇, -NR₇C(O)OR₇, -OCH₂C(O)R₇, -SCH₂C(O)R₇,

- 5
10
- NR₇CH₂C(O)R₇, -OCH₂C(O)OR₇, -SCH₂C(O)OR₇, -NR₇CH₂C(O)OR₇,
 -OCH₂C(O)NR₁₀R₁₁, -SCH₂C(O)NR₁₀R₁₁, -NR₇CH₂C(O)NR₁₀R₁₁,
 -OS(O)_pR₇, -SS(O)_pR₇, -NR₇S(O)_pR₇, -OS(O)_pNR₁₀R₁₁, -SS(O)_pNR₁₀R₁₁,
 -NR₇S(O)_pNR₁₀R₁₁, -OS(O)_pOR₇, -SS(O)_pOR₇, -NR₇S(O)_pOR₇, -OC(S)R₇,
 -SC(S)R₇, -NR₇C(S)R₇, -OC(S)OR₇, -SC(S)OR₇, -NR₇C(S)OR₇,
 -OC(S)NR₁₀R₁₁, -SC(S)NR₁₀R₁₁, -NR₇C(S)NR₁₀R₁₁, -OC(NR₈)R₇,
 -SC(NR₈)R₇, -NR₇C(NR₈)R₇, -OC(NR₈)OR₇, -SC(NR₈)OR₇,
 -NR₇C(NR₈)OR₇, -OC(NR₈)NR₁₀R₁₁, -SC(NR₈)NR₁₀R₁₁,
 -NR₇C(NR₈)NR₁₀R₁₁, -C(O)R₇, -C(O)OR₇, -C(O)NR₁₀R₁₁, -C(O)SR₇,
 -C(S)R₇, -C(S)OR₇, -C(S)NR₁₀R₁₁, -C(S)SR₇, -C(NR₈)OR₇, -C(NR₈)R₇,
 -C(NR₈)NR₁₀R₁₁, -C(NR₈)SR₇, -S(O)_pOR₇, -S(O)_pNR₁₀R₁₁, or -S(O)_pR₇.

17. The compound of Claim 16, wherein:


R₁ is -SH or -OH;

R₃ and R₂₅ are -OH;



R₆ is a lower alkyl, C3-C6 cycloalkyl, lower alkoxy, a lower alkyl sulfanyl, or -NR₁₀R₁₁; and

R₉, for each occurrence, is independently selected from the group consisting of -OH, -SH, halo, a lower haloalkyl, cyano, a lower alkyl, a lower alkoxy, and a lower alkyl sulfanyl.

18. The compound of Claim 14, wherein the compound is represented by the following structural formula:



19. The compound of Claim 18, wherein R_6 is a C1-C6 alkyl, a C1-C6 haloalkyl, a C1-C6 alkoxy, a C1-C6 haloalkoxy, a C1-C6 alkyl sulfanyl or a C3-C6 cycloalkyl.
- 5 20. The compound of Claim 19, wherein R_1 and R_3 are each, independently, -OH, -SH, or -NHR₇.
21. The compound of Claim 20, wherein:
- R_1 is -SH or -OH;
- 10 R_3 and R_{25} are -OH;
- R_6 is a lower alkyl, C3-C6 cycloalkyl, lower alkoxy, a lower alkyl sulfanyl, or -NR₁₀R₁₁.
22. The compound of Claim 1, wherein the compound is represented by one of the following structural formulas:
- 15



wherein:

- X_3 and X_4 are each, independently, N, N(O), N⁺(R₁₇), CH or CR₆;
- X_5 is O, S, NR₁₇, CH=CH, CH=CR₆, CR₆=CH, CR₆=CR₆, CH=N, CR₆=N, CH=N(O), CR₆=N(O), N=CH, N=CR₆, N(O)=CH, N(O)=CR₆, N⁺(R₁₇)=CH, N⁺(R₁₇)=CR₆, CH=N⁺(R₁₇), CR₆=N⁺(R₁₇), or N=N;
- 20 R_6 , for each occurrence, is independently a substituent selected from the group consisting of an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, halo, cyano, nitro, guanadino, a haloalkyl, a heteroalkyl, -NR₁₀R₁₁, -OR₇, -C(O)R₇, -C(O)OR₇, -OC(O)R₇,
- 25



-C(O)NR₁₀R₁₁, -NR₈C(O)R₇, -SR₇, -S(O)_pR₇, -OS(O)_pR₇, -S(O)_pOR₇,
 -NR₈S(O)_pR₇, or -S(O)_pNR₁₀R₁₁, -S(O)_pOR₇, -OP(O)(OR₇)₂, or
 -SP(O)(OR₇)₂;

R₁₇, for each occurrence, is independently an alkyl or an aralkyl; and







5

n is zero or an integer from 1 to 4.


23. The compound of Claim 22, wherein the compound is selected from the group consisting of:



10



15



wherein:

- R_{25} is a halo, a haloalkyl, a haloalkoxy, a heteroalkyl, -OH, -SH,
 -NHR₇, -(CH₂)_kOH, -(CH₂)_kSH, -(CH₂)_kNR₇H, -OCH₃, -SCH₃, -NHCH₃,
 5 -OCH₂CH₂OH, -OCH₂CH₂SH, -OCH₂CH₂NR₇H, -SCH₂CH₂OH,
 -SCH₂CH₂SH, -SCH₂CH₂NR₇H, -OC(O)NR₁₀R₁₁, -SC(O)NR₁₀R₁₁,
 -NR₇C(O)NR₁₀R₁₁, -OC(O)R₇, -SC(O)R₇, -NR₇C(O)R₇, -OC(O)OR₇,
 -SC(O)OR₇, -NR₇C(O)OR₇, -OCH₂C(O)R₇, -SCH₂C(O)R₇,
 -NR₇CH₂C(O)R₇, -OCH₂C(O)OR₇, -SCH₂C(O)OR₇, -NR₇CH₂C(O)OR₇,
 10 -OCH₂C(O)NR₁₀R₁₁, -SCH₂C(O)NR₁₀R₁₁, -NR₇CH₂C(O)NR₁₀R₁₁,
 -OS(O)_pR₇, -SS(O)_pR₇, -NR₇S(O)_pR₇, -OS(O)_pNR₁₀R₁₁, -SS(O)_pNR₁₀R₁₁,
 -NR₇S(O)_pNR₁₀R₁₁, -OS(O)_pOR₇, -SS(O)_pOR₇, -NR₇S(O)_pOR₇, -OC(S)R₇,
 -SC(S)R₇, -NR₇C(S)R₇, -OC(S)OR₇, -SC(S)OR₇, -NR₇C(S)OR₇,
 -OC(S)NR₁₀R₁₁, -SC(S)NR₁₀R₁₁, -NR₇C(S)NR₁₀R₁₁, -OC(NR₈)R₇,
 15 -SC(NR₈)R₇, -NR₇C(NR₈)R₇, -OC(NR₈)OR₇, -SC(NR₈)OR₇,
 -NR₇C(NR₈)OR₇, -OC(NR₈)NR₁₀R₁₁, -SC(NR₈)NR₁₀R₁₁,
 -NR₇C(NR₈)NR₁₀R₁₁, -C(O)R₇, -C(O)OR₇, -C(O)NR₁₀R₁₁, -C(O)SR₇,
 -C(S)R₇, -C(S)OR₇, -C(S)NR₁₀R₁₁, -C(S)SR₇, -C(NR₈)OR₇, -C(NR₈)R₇,
 -C(NR₈)NR₁₀R₁₁, -C(NR₈)SR₇, -S(O)_pOR₇, -S(O)_pNR₁₀R₁₁, or -S(O)_pR₇,
 20 -S(O)_pOR₇, -OP(O)(OR₇)₂, or -SP(O)(OR₇)₂; and

k is 1, 2, 3, or 4.

24. The compound of Claim 1, selected from the group consisting of
 3-(2-Hydroxyphenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
 25 3-(2,4-Dihydroxyphenyl)-4-[4-(2-methoxyethoxy)-naphthalen-1-yl]-5-mercapto-
 triazole;
 3-(2,4-Dihydroxyphenyl)-4-(2-methyl-4-bromophenyl)-5-mercapto-triazole;
 3-(3,4-Dihydroxyphenyl)-4-(6-methoxy-naphthalen-1-yl)-5-mercapto-triazole;
 3-(3,4-Dihydroxyphenyl)-4-(6-ethoxy-naphthalen-1-yl)-5-mercapto-triazole;

- 3-(3,4-Dihydroxyphenyl)-4-(6-propoxy-naphthalen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(5-methoxy-naphthalen-1-yl)-5-mercapto-triazole;
3-(3,4-Dihydroxyphenyl)-4-(6-isopropoxy-naphthalen-1-yl)-5-mercapto-triazole;
5 3-(2,4-Dihydroxyphenyl)-4-(2,6-diethylphenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(2-methy-6-ethylphenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(2,6-diisopropylphenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(1-ethyl-indol-4-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(2,3-dihydro-benzo[1,4]dioxin-5-yl)-5-mercapto-triazole;
10 triazole;
3-(2,4-Dihydroxyphenyl)-4-(3-methylphenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(4-methylphenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(2-chlorophenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(3-chlorophenyl)-5-mercapto-triazole;
15 3-(2,4-Dihydroxyphenyl)-4-(4-chlorophenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(2-methoxyphenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(3-methoxyphenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(3-fluorophenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(2-ethylphenyl)-5-mercapto-triazole;
20 3-(2-Hydroxy-4-fluorophenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
3-(2-Hydroxy-4-aminophenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(2-methyl-4-butyl-phenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(2,4-dimethyl-phenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(2,6-dimethyl-phenyl)-5-mercapto-triazole;
25 3-(2,4-Dihydroxyphenyl)-4-(2,6-dimethyl-phenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(4-fluorophenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(2-methylsulfanylphenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(naphthalene-2-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(2,3-dimethylphenyl)-5-mercapto-triazole;
30 3-(2,4-Dihydroxyphenyl)-4-(2-methyl-4-fluorophenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(acenaphthalen-5-yl)-5-mercapto-triazole;
3-(2-Hydroxy-4-methoxy-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;

- 3-(2,4-Dihydroxyphenyl)-4-(2,3-dichlorophenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(5-methoxynaphthalen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(pyren-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(quinolin-5-yl)-5-mercapto-triazole;
5 3-(2,4-Dihydroxyphenyl)-4-(1,2,3,4-tetrahydronaphthalen-5-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(anthracen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(biphenyl-2-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-6-methyl-phenyl)-4-(naphthalene-1-yl)-5-mercapto-triazole;
10 3-(2,4-Dihydroxyphenyl)-4-(4-pentyloxyphenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(4-octyloxyphenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(4-chloronaphthalen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(7-carboxymethoxy-naphthalen-1-yl)-5-
15 mercapto-triazole;
3-(2,4-Dihydroxyphenyl)-4-(2-methyl-quinolin-4-yl)-5-mercapto-triazole;
3-(3-Hydroxypyridin-4-yl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
3-(2-Hydroxy-4-acetylamino-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-phenyl)-4-(1,2,3,4-tetrahydronaphthalen-1-yl)-5-mercapto-
20 triazole;
3-(2,4-Dihydroxy-phenyl)-4-(2,3-dihydro-benzo[1,4]dioxin-5-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-phenyl)-4-(3,5-dimethoxyphenyl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-phenyl)-4-(2,3-dimethyl-1H-indol-4-yl)-5-mercapto-triazole;
25 3-(2,4-Dihydroxy-3-propyl-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
3-(4,6-Dihydroxy-1-ethyl-pyridin-3-yl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
3-(4,6-Dihydroxy-1-methyl-pyridin-3-yl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-phenyl)-4-(3,5-di-tert-butylphenyl)-5-mercapto-triazole;
30 3-(2,6-Dihydroxy-5-fluoro-pyridin-3-yl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-5-methyl-phenyl)-4-(naphthalene-1-yl)-5-mercapto-triazole;
3-[2,4-Dihydroxy-phenyl]-4-(3-benzoylphenyl)-5-mercapto-triazole;

- 3-(2,4-Dihydroxy-phenyl)-4-(4-carboxy-naphthalen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-phenyl)-4-[4-(N,N-dimethylcarbamoyl)-naphthalen-1-yl]-5-mercapto-triazole;
3-(2,4-Dihydroxy-phenyl)-4-(4-propoxy-naphthalen-1-yl)-5-mercapto-triazole;
5 3-(2,4-Dihydroxy-phenyl)-4-(4-isopropoxy-naphthalen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-phenyl)-4-(5-isopropoxy-naphthalen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-phenyl)-4-(isoquinolin-5-yl)-5-mercapto-triazole;
10 3-(2,4-Dihydroxy-phenyl)-4-(5-propoxy-naphthalen-1-yl)-5-mercapto-triazole;
3-(2-Hydroxy-4-methanesulfonamino-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-3,6-dimethyl-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
15 3-(2,4-Dihydroxy-phenyl)-4-[7-(2-methoxyethoxy)-naphthalen-1-yl]-5-mercapto-triazole;
3-(2,4-Dihydroxy-5-hexyl-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(4-methoxy-naphthalen-1-yl)-5-mercapto-triazole;
20 3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(6-methoxy-naphthalin-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-3-chloro-5-ethyl-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(2,3-dimethoxy-4-methoxy-phenyl)-5-mercapto-triazole;
25 3-(2,4-Dihydroxy-phenyl)-4-(7-isopropoxy-naphthalen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-phenyl)-4-(7-ethoxy-naphthalen-1-yl)-5-mercapto-triazole;
3-(2,4-Dihydroxy-phenyl)-4-(7-propoxy-naphthalen-1-yl)-5-mercapto-triazole;
30 3-(2-Hydroxy-4-methoxymethoxy-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;

- 3-[2-Hydroxy-4-(2-hydroxy-ethoxy)-phenyl]-4-(naphthalen-1-yl)-5-mercapto-triazole;
- 3-(2,4-Dihydroxyphenyl)-4-(7-methoxy-naphthalen-1-yl)-5-mercapto-triazole;
- 3-(2,4-Dihydroxyphenyl)-4-(5-methoxy-naphthalen-1-yl)-5-mercapto-triazole;
- 5 3-(2,4-Dihydroxyphenyl)-4-(4-hydroxy-naphthalen-1-yl)-5-mercapto-triazole;
- 3-(2,4-Dihydroxyphenyl)-4-(1-isopropyl-indol-4-yl)-5-mercapto-triazole;
- 3-(2,4-Dihydroxy-5-tert-butyl-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
- 3-(2,4-Dihydroxy-5-propyl-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
- 3-(2,4-Dihydroxy-3-methyl-5-ethyl-phenyl)-4-(naphthalen-1-yl)-5-mercapto-
- 10 triazole;
- 3-(2,4-Dihydroxy-5-isobutyl-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
- 3-(2,4-Dihydroxy-phenyl)-4-(2,3-dimethoxy-phenyl)-5-mercapto-triazole;
- 3-(2,4-Dihydroxy-phenyl)-4-(2-methoxy-3-chloro-phenyl)-5-mercapto-triazole;
- 3-(2,4-Dihydroxy-phenyl)-4-(indol-4-yl)-5-mercapto-triazole;
- 15 3-(2,4-Dihydroxy-phenyl)-4-[1-(2-methoxyethoxy)-indol-4-yl]-5-mercapto-triazole;
- 3-(2,4-Dihydroxy-phenyl)-4-(naphthalen-1-yl)-5-hydroxy-triazole;
- 3-(1-Oxo-3-hydroxy-pyridin-4-yl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
- 3-(2,5-Dihydroxy-4-carboxy)-4-(naphthalen-1-yl)-5-mercapto-triazole;
- 20 3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-indol-4-yl)-5-mercapto-triazole;
- 3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-[1-(dimethyl-carbamoyl)-indol-4-yl]-5-mercapto-triazole;
- 3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(1-ethyl-benzoimidazol-4-yl)-5-mercapto-
- 25 triazole;
- 3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(1,2,3-trimethyl-indol-5-yl)-5-mercapto-triazole;
- 3-(2,5-Dihydroxy-4-hydroxymethyl-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
- 30 3-(2-Hydroxy-4-amino-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
- 3-(2-Hydroxy-4-acetyl-amino-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
- 3-(2,4-Dihydroxy-3-chloro-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;

- 3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(3-methoxy-phenyl)-5-hydroxy-triazole;
3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(naphthalen-1-yl)-5-hydroxy-triazole;
3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-indol-3-yl)-5-hydroxy-triazole;
3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-indol-4-yl)-5-amino-triazole;
5 3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(3-methoxy-phenyl)-5-amino-triazole;
3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(naphthalen-1-yl)-5-amino-triazole;
3-(2-Hydroxy-5-ethyloxy-phenyl)-4-(naphthalen-1-yl)-5-hydroxy-triazole;
3-(2-Hydroxy-5-isopropyl-phenyl)-4-(naphthalen-1-yl)-5-hydroxy-triazole;
3-(2-Dihydroxy-phenyl)-4-(7-fluoro-naphthalen-1-yl)-5-hydroxy-triazole;
10 3-(2,4-Dihydroxy-phenyl)-4-(2,3-difluorophenyl)-5-hydroxy-triazole;
3-(2,4-Dihydroxy-phenyl)-4-[2-(1H-tetrazol-5-yl)-phenyl]-5-hydroxy-triazole;
3-(2,4-Dihydroxy-phenyl)-4-(benzothiazol-4-yl)-5-hydroxy-triazole;
3-(2,4-Dihydroxy-phenyl)-4-(9H-purin-6-yl)-5-hydroxy-triazole;
3-(2,4-Dihydroxy-phenyl)-4-{4-[2-(morpholin-1-yl)-ethoxy]-phenyl}-5-
15 hydroxy-triazole;
3-(2,4-Dihydroxy-phenyl)-4-cyclopentyl-5-hydroxy-triazole;
3-(2,4-Dihydroxy-phenyl)-4-phenyl-5-(sulfamoylamino)-triazole;
3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(naphthalene-1-yl)-5-ureido-triazole;
3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(2,3-difluorophenyl)-5-ureido-triazole;
20 3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-indol-4-yl)-5-ureido-triazole;
3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(quinolin-5-yl)-5-ureido-triazole;
3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(naphthalene-1-yl)-5-carbamoyloxy-
triazole;
3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(3-trifluoromethyl-phenyl)-5-
25 carbamoyloxy-triazole;
3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(1-methyl-indol-4-yl)-5-carbamoyloxy-
triazole;
3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(8-methoxy-quinolin-5-yl)-5-
carbamoyloxy-triazole;
30 3-(2,4-Dihydroxy-5-isopropyl-phenyl)-4-(3-methyl-quinolin-5-yl)-5-
carboxy-amino-triazole;


- 3-(2,4-Dihydroxy-phenyl)-4-(1-methyl-2-chloro-indol-4-yl)-5-carbamoyloxy-triazole;
- 3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-[3,5-di-(trifluoromethyl)-phenyl]-5-carbamoyloxy-triazole;
- 5 3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(3-trifluoromethyl-phenyl)-5-(sulfamoylamino)-triazole;
- 3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(naphthalene-1-yl)-5-(sulfamoylamino)-triazole;
- 3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(1-isopropyl-benzoimidazol-4-yl)-5-(sulfamoylamino)-triazole;
- 10 3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(3-isopropylphenyl)-5-(thiocarboxy-amino)-triazole;
- 3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(3-isopropoxy-phenyl)-5-(sulfamoyloxy)-triazole;
- 15 3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(naphthalene-1-yl)-5-(sulfamoyloxy)-triazole;
- 3-(2,4-Dihydroxy-5-methoxy-phenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
- 3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(5-hydroxy-naphthalen-1-yl)-5-mercapto-triazole;
- 20 3-(2,4-Dihydroxy-phenyl)-4-(naphthalen-1-ylmethyl)-5-mercapto-triazole;
- 3-(2-Hydroxy-4-methoxyphenyl)-4-(naphthalen-1-yl)-5-mercapto-triazole;
- 3-(2,4-Dihydroxy-phenyl)-4-(biphenyl-3-yl)-5-mercapto-triazole;
- 3-(2,4-Dihydroxy-phenyl)-4-(2-methyl-5-hydroxymethyl-phenyl)-5-mercapto-triazole;
- 25 3-(2,4-Dihydroxy-phenyl)-4-(1-dimethylcarbamoyl-indol-4-yl)-5-mercapto-triazole;
- 3-(2,4,5-Trihydroxy-phenyl)-4-(naphthalene-1-yl)-5-mercapto-triazole;
- 3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(2,3-dimethyl-indol-5-yl)-5-mercapto-triazole;
- 30 3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(3-t-butyl-4-methoxy-phenyl)-5-mercapto-triazole;

3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(1-ethyl-1H-benzimidazol-4-yl)-5-mercapto-triazole, HCl salt;

3-(2,4-Dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-5-mercapto-triazole; and

- 5 3-(2,4-Dihydroxy-5-cyclopropyl-phenyl)-4-(naphthalene-1-yl)-5-mercapto-triazole
or a tautomer, pharmaceutically acceptable salt, solvate, clathrate, or a prodrug thereof.

- 10 25. A compound represented by the following structural formula:



or a tautomer, pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein:

X_{41} is O, S, or NR_{42} ;

- 15 X_{42} is CR_{44} or N;

Y_{40} is N or CR_{43} ;

Y_{41} is N or CR_{45} ;

Y_{42} , for each occurrence, is independently N, C or CR_{46} ;

Z is OH, SH, or NHR_7 ;


- 20 R_{41} is -H, -OH, -SH, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, halo, cyano, nitro, guanadino, a haloalkyl, a heteroalkyl, an
25 alkoxy or cycloalkoxy, a haloalkoxy, $-NR_{10}R_{11}$, $-OR_7$, $-C(O)R_7$, $-C(O)OR_7$,

- C(S)R₇, -C(O)SR₇, -C(S)SR₇, -C(S)OR₇, -C(S)NR₁₀R₁₁, -C(NR₈)OR₇,
 -C(NR₈)R₇, -C(NR₈)NR₁₀R₁₁, -C(NR₈)SR₇, -OC(O)R₇, -OC(O)OR₇,
 -OC(S)OR₇, -OC(NR₈)OR₇, -SC(O)R₇, -SC(O)OR₇, -SC(NR₈)OR₇,
 -OC(S)R₇, -SC(S)R₇, -SC(S)OR₇, -OC(O)NR₁₀R₁₁, -OC(S)NR₁₀R₁₁,
 5 -OC(NR₈)NR₁₀R₁₁, -SC(O)NR₁₀R₁₁, -SC(NR₈)NR₁₀R₁₁, -SC(S)NR₁₀R₁₁,
 -OC(NR₈)R₇, -SC(NR₈)R₇, -C(O)NR₁₀R₁₁, -NR₈C(O)R₇, -NR₇C(S)R₇,
 -NR₇C(S)OR₇, -NR₇C(NR₈)R₇, -NR₇C(O)OR₇, -NR₇C(NR₈)OR₇,
 -NR₇C(O)NR₁₀R₁₁, -NR₇C(S)NR₁₀R₁₁, -NR₇C(NR₈)NR₁₀R₁₁, -SR₇,
 -S(O)_pR₇, -OS(O)_pR₇, -OS(O)_pOR₇, -OS(O)_pNR₁₀R₁₁, -S(O)_pOR₇,
 10 -NR₈S(O)_pR₇, -NR₇S(O)_pNR₁₀R₁₁, -NR₇S(O)_pOR₇, -S(O)_pNR₁₀R₁₁,
 -SS(O)_pR₇, -SS(O)_pOR₇, -SS(O)_pNR₁₀R₁₁, -OP(O)(OR₇)₂, or -SP(O)(OR₇)₂;
- R₄₂ is -H, an optionally substituted alkyl, an optionally substituted alkenyl, an
 optionally substituted alkynyl, an optionally substituted cycloalkyl, an
 optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an
 15 optionally substituted aryl, an optionally substituted heteroaryl, an optionally
 substituted aralkyl, an optionally substituted heteraralkyl, hydroxyalkyl,
 alkoxyalkyl, a haloalkyl, a heteroalkyl, -C(O)R₇, -(CH₂)_mC(O)OR₇,
 -C(O)OR₇, -OC(O)R₇, -C(O)NR₁₀R₁₁, -S(O)_pR₇, -S(O)_pOR₇, or
 -S(O)_pNR₁₀R₁₁;
- 20 R₄₃ and R₄₄ are, independently, -H, -OH, an optionally substituted alkyl, an
 optionally substituted alkenyl, an optionally substituted alkynyl, an
 optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an
 optionally substituted heterocyclyl, an optionally substituted aryl, an
 optionally substituted heteroaryl, an optionally substituted aralkyl, an
 25 optionally substituted heteraralkyl, hydroxyalkyl, alkoxyalkyl, halo, cyano,
 nitro, guanidino, a haloalkyl, a heteroalkyl, -C(O)R₇, -C(O)OR₇,
 -OC(O)R₇, -C(O)NR₁₀R₁₁, -NR₈C(O)R₇, -SR₇, -S(O)_pR₇, -OS(O)_pR₇,
 -S(O)_pOR₇, -NR₈S(O)_pR₇, -S(O)_pNR₁₀R₁₁, or R₄₃ and R₄₄ taken together
 with the carbon atoms to which they are attached form an optionally
 30 substituted cycloalkenyl, an optionally substituted aryl, an optionally
 substituted heterocyclyl, or an optionally substituted heteroaryl;

- R_{45} is -H, -OH, -SH, -NR₇H, -OR₂₆, -SR₂₆, -NHR₂₆, -O(CH₂)_mOH,
 -O(CH₂)_mSH, -O(CH₂)_mNR₇H, -S(CH₂)_mOH, -S(CH₂)_mSH,
 -S(CH₂)_mNR₇H, -OC(O)NR₁₀R₁₁, -SC(O)NR₁₀R₁₁, -NR₇C(O)NR₁₀R₁₁,
 -OC(O)R₇, -SC(O)R₇, -NR₇C(O)R₇, -OC(O)OR₇, -SC(O)OR₇,
 5 -NR₇C(O)OR₇, -OCH₂C(O)R₇, -SCH₂C(O)R₇, -NR₇CH₂C(O)R₇,
 -OCH₂C(O)OR₇, -SCH₂C(O)OR₇, -NR₇CH₂C(O)OR₇,
 -OCH₂C(O)NR₁₀R₁₁, -SCH₂C(O)NR₁₀R₁₁, -NR₇CH₂C(O)NR₁₀R₁₁,
 -OS(O)_pR₇, -SS(O)_pR₇, -NR₇S(O)_pR₇, -OS(O)_pNR₁₀R₁₁, -SS(O)_pNR₁₀R₁₁,
 -NR₇S(O)_pNR₁₀R₁₁, -OS(O)_pOR₇, -SS(O)_pOR₇, -NR₇S(O)_pOR₇, -OC(S)R₇,
 10 -SC(S)R₇, -NR₇C(S)R₇, -OC(S)OR₇, -SC(S)OR₇, -NR₇C(S)OR₇,
 -OC(S)NR₁₀R₁₁, -SC(S)NR₁₀R₁₁, -NR₇C(S)NR₁₀R₁₁, -OC(NR₈)R₇,
 -SC(NR₈)R₇, -NR₇C(NR₈)R₇, -OC(NR₈)OR₇, -SC(NR₈)OR₇,
 -NR₇C(NR₈)OR₇, -OC(NR₈)NR₁₀R₁₁, -SC(NR₈)NR₁₀R₁₁, or
 -NR₇C(NR₈)NR₁₀R₁₁;
- 15 R_{46} , for each occurrence, is independently, selected from the group consisting of H,
 an optionally substituted alkyl, an optionally substituted alkenyl, an
 optionally substituted alkynyl, an optionally substituted cycloalkyl, an
 optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an
 optionally substituted aryl, an optionally substituted heteroaryl, an optionally
 20 substituted aralkyl, an optionally substituted heteraralkyl, halo, cyano, nitro,
 guanadino, a haloalkyl, a heteroalkyl, -NR₁₀R₁₁, -OR₇, -C(O)R₇,
 -C(O)OR₇, -OC(O)R₇, -C(O)NR₁₀R₁₁, -NR₈C(O)R₇, -SR₇, -S(O)_pR₇,
 -OS(O)_pR₇, -S(O)_pOR₇, -NR₈S(O)_pR₇, or -S(O)_pNR₁₀R₁₁;
- R_7 and R_8 , for each occurrence, are, independently, -H, an optionally substituted
 25 alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an
 optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an
 optionally substituted heterocyclyl, an optionally substituted aryl, an
 optionally substituted heteroaryl, an optionally substituted aralkyl, or an
 optionally substituted heteraralkyl;
- 30 R_{10} and R_{11} , for each occurrence, are independently -H, an optionally substituted
 alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an
 optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an


- optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R₁₀ and R₁₁, taken together with the nitrogen to which they are attached, form an optionally substituted heterocyclyl or an optionally substituted heteroaryl;
- 5 R₂₆, for each occurrence is, is independently, a lower alkyl;
- p, for each occurrence, is, independently, 1 or 2; and
- m, for each occurrence, is independently, 1, 2, 3, or 4.
- 10 26. The compound of claim 25, wherein X₄₁ is NR₄₂ and X₄₂ is CR₄₄.
27. The compound of claim 25, wherein X₄₁ is NR₄₂ and X₄₂ is N.
- 15 28. The compound of claim 25, wherein R₄₁ is selected from the group consisting of -H, lower alkyl, lower alkoxy, lower cycloalkyl, and lower cycloalkoxy.
29. The compound of claim 25, wherein R₄₁ is selected from the group consisting of -H, methyl, ethyl, propyl, isopropyl, cyclopropyl, methoxy, ethoxy, propoxy, and cyclopropoxy.
- 20 30. The compound of claim 25, wherein X₄₁ is NR₄₂, and R₄₂ is selected from the group consisting of -H, a lower alkyl, a lower cycloalkyl, -C(O)N(R₂₇)₂, and -C(O)OH, wherein each R₂₇ is independently -H or a lower alkyl.
- 25 31. The compound of claim 25, wherein X₄₁ is NR₄₂, and R₄₂ is selected from the group consisting of -H, methyl, ethyl, n-propyl, isopropyl, cyclopropyl, n-butyl, *sec*-butyl, *tert*-butyl, n-pentyl, n-hexyl, -C(O)OH, -(CH₂)_mC(O)OH, -CH₂OCH₃, -CH₂CH₂OCH₃, and -C(O)N(CH₃)₂.
- 30

32. The compound of claim 25, wherein R_{43} and R_{44} are, independently, selected from the group consisting of -H, methyl, ethyl, propyl, isopropyl, cyclopropyl, methoxy, ethoxy, propoxy, and cyclopropoxy.
- 5 33. The compound of claim 25, wherein X_{42} is CR_{44} ; Y_{40} is CR_{43} ; and R_{43} and R_{44} together with the carbon atoms to which they are attached form a cycloalkenyl, an aryl, heterocyclyl, or heteroaryl ring.
- 10 34. The compound of claim 33, wherein R_{43} and R_{44} together with the carbon atoms to which they are attached form a C_5 - C_8 cycloalkenyl or a C_5 - C_8 aryl.
35. The compound of claim 33, wherein R_{45} or CR_{45} is selected from the group consisting of -H, -OH, -SH, - NH_2 , a lower alkoxy, a lower alkyl amino, and a lower dialkyl amino.
- 15 36. The compound of Claim 35, wherein R_{45} is selected from the group consisting of -H, -OH, methoxy and ethoxy.
37. The compound of claim 33, wherein X_{41} is O.
- 20 38. The compound of claim 37, wherein said compound is selected from the group consisting of:
3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(2-methyl-7-methoxy-benzofuran-4-yl)-5-mercapto-[1,2,4]triazole,
25 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(benzofuran-5-yl)-5-mercapto-[1,2,4]triazole, and
3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(2-methyl-1,3-benzoxaz-5-yl)-5-mercapto-[1,2,4]triazole.
- 30 39. The compound of claim 25, wherein Z is -OH or -SH.
40. A compound of Claim 25 represented by the following structural formula:



or a tautomer, pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein Z_1 is $-OH$ or $-SH$.

- 5 41. The compound of claim 40, wherein X_{42} is CR_{44} , and R_{43} and R_{44} are, independently, selected from the group consisting of $-H$, methyl, ethyl, propyl, isopropyl, cyclopropyl, methoxy, ethoxy, propoxy, and cyclopropoxy.
- 10 42. The compound of claim 40, wherein X_{42} is CR_{44} , and R_{43} and R_{44} , taken together with the carbon atoms to which they are attached, form a cycloalkenyl, aryl, heterocyclyl, or heteroaryl ring.
- 15 43. The compound of claim 42, wherein R_{43} and R_{44} , taken together with the carbon atoms to which they are attached, form a C_5 - C_8 cycloalkenyl or a C_5 - C_8 aryl.
44. The compound of claim 40, wherein X_{42} is CR_{44} .
- 20 45. The compound of claim 40, wherein X_{42} is N .
46. A compound of Claims 40 represented by the following structural formula:



or a tautomer, pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein:

X₄₅ is CR₅₄ or N;

5 Z₁ is -OH or -SH;

R₅₆ is selected from the group consisting of -H, methyl, ethyl, isopropyl, and cyclopropyl;

R₅₂ is selected from the group consisting of -H, methyl, ethyl, n-propyl, isopropyl, n-butyl, n-pentyl, n-hexyl, -(CH₂)₂OCH₃, -CH₂C(O)OH, and -C(O)N(CH₃)₂;

10 R₅₃ and R₅₄ are each, independently, -H, methyl, ethyl, or isopropyl; or R₅₃ and R₅₄ taken together with the carbon atoms to which they are attached form a phenyl, cyclohexenyl, or cyclooctenyl ring; and

R₅₅ is selected from the group consisting of -H, -OH, -OCH₃, and -OCH₂CH₃.

15 47. The compound of claim 25, wherein said compound is selected from the group consisting of:

3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole,

20 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole,

3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-hydroxy-[1,2,4]triazole, and

3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-isopropyl-indol-4-yl)-5-hydroxy-[1,2,4]triazole.

25

48. The compound of claim 25, wherein said compound is selected from the group consisting of:
3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indazol-5-yl)-5-mercapto-[1,2,4]triazole and
5 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-indazol-6-yl)-5-mercapto-[1,2,4]triazole.
49. The compound of claim 25, wherein said compound is selected from the group consisting of:
10 3-(2,4-dihydroxyphenyl)-4-(1-ethyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,
3-(2,4-dihydroxyphenyl)-4-(1-isopropyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,
3-(2,4-dihydroxyphenyl)-4-(indol-4-yl)-5-mercapto-[1,2,4]triazole,
3-(2,4-dihydroxyphenyl)-4-(1-methoxyethyl-indol-4-yl)-5-mercapto-
15 [1,2,4]triazole,
3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,
3-(2,4-dihydroxyphenyl)-4-(1-dimethylcarbamoyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,
20 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-propyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,
3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,2,3-trimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(2,3-dimethyl-indol-5-yl)-5-mercapto-
25 [1,2,4]triazole,
3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-acetyl-2,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-5-mercapto-[1,2,4]triazole,
30 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-propyl-2,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,

- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(N-methyl-tetrahydrocarbozol-7-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(N-methyl-cyclononan[a]indol-5-yl)-5-mercapto-[1,2,4]triazole,
- 5 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-n-butyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-n-pentyl-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-n-hexyl-indol-4-yl)-5-mercapto-
- 10 [1,2,4]triazole,
- 3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1-(1-methylcyclopropyl)-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- 15 3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1,2,3-trimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-5-mercapto-[1,2,4]triazole disodium salt,
- 3-(2,4-dihydroxy-5-*tert*-butyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-
- 20 5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1-propyl-7-methoxy-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-methyl-3-ethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- 25 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-isopropyl-7-methoxy-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-methyl-3-isopropyl-indol-5-yl)-5-
- 30 mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(N-ethyl-carbozol-7-yl)-5-mercapto-[1,2,4]triazole,

- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-7-hydroxy-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-isopropyl-7-ethoxy-indol-4-yl)-5-mercapto-[1,2,4]triazole,
- 5 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,2-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(N-methyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- 10 3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1,3-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-cyclopropyl-phenyl)-4-(1-methyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- 15 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1H-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1,2-dimethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-ethyl-indol-5-yl)-5-mercapto-[1,2,4]triazole, and
- 20 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-propyl-indol-5-yl)-5-mercapto-[1,2,4]triazole.
50. The compound of claim 25, wherein said compound is selected from the group consisting of
- 25 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-ethyl-benzimidazol-4-yl)-5-mercapto-[1,2,4]triazole,
- 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-ethyl-benzimidazol-4-yl)-5-mercapto-[1,2,4]triazole HCL salt,
- 30 3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(2-methyl-3-ethyl-benzimidazol-5-yl)-5-mercapto-[1,2,4]triazole,

3-(2,4-dihydroxy-5-ethyl-phenyl)-4-(1-ethyl-2-methyl-benzimidazol-5-yl)-5-mercapto-[1,2,4]triazole, and

3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(1-methyl-2-trifluoromethyl-benzimidazol-5-yl)-5-mercapto-[1,2,4]triazole.

5

51. The compound of claim 25, wherein said compound is 3-(2,4-dihydroxy-5-isopropyl-phenyl)-4-(N-methyl-indol-5-yl)-5-mercapto-[1,2,4]triazole.
52. A method for inhibiting Hsp90 in a cell, comprising administering to the cell an effective amount of any of the compounds of Claims 1-51 or a tautomer, pharmaceutically acceptable salt, solvate, clathrate or a prodrug thereof.
53. A method for treating or preventing a proliferation disorder in a mammal, comprising administering to the cell an effective amount of any of the compounds of Claims 1-51 or a tautomer, pharmaceutically acceptable salt, solvate, clathrate or a prodrug thereof.
54. A method for treating cancer in a mammal, comprising administering to the mammal an effective amount of any of the compounds of Claims 1-51 or a tautomer, pharmaceutically acceptable salt, solvate, clathrate or a prodrug thereof.
55. A method of inducing degradation of a c-kit protein, comprising administering to the mammal an effective amount of any of the compounds of Claims 1-51 or a tautomer, pharmaceutically acceptable salt, solvate, clathrate or a prodrug thereof.
56. A method of treating a c-kit associated cancer in a mammal, comprising administering to the mammal an effective amount of any of the compounds of Claims 1-51 or a tautomer, pharmaceutically acceptable salt, solvate, clathrate or a prodrug thereof.

57. A method of inducing degradation of a c-met protein, comprising administering to the mammal an effective amount of any of the compounds of Claims 1-51 or a tautomer, pharmaceutically acceptable salt, solvate, clathrate or a prodrug thereof.
- 5
58. A method of treating a c-met associated cancer in a mammal, comprising administering to the mammal an effective amount of any of the compounds of Claims 1-51 or a tautomer, pharmaceutically acceptable salt, solvate, clathrate or a prodrug thereof.
- 10
59. A pharmaceutical composition, comprising a pharmaceutically acceptable carrier and one or more compounds of Claims 1-51.
- 15

1/10




Fig. 1

2/10




Fig. 2




Fig. 3

Hsp90 inhibitors of the invention induce degradation
of c-Kit in Kasumi-1 cells




Fig. 4

5/10

Hsp90 inhibitors induce degradation of c-Kit in Kasumi-1 cells




Fig. 5

6/10

C-Met Degradation induced by Hsp90 inhibitors in NCI-H1193 cells




Fig. 6

7/10




Fig. 7

8/10




Fig. 8

9/10




Fig. 9

10/10




Fig. 10

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2005/041779

A. CLASSIFICATION OF SUBJECT MATTER
 INV. C07D249/12 C07D401/04 C07D405/04 C07D403/04 A61K31/4196
 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 2005/000300 A (VERNALIS LIMITED; CANCER RESEARCH TECHNOLOGY LTD; THE INSTITUTE OF CA) 6 January 2005 (2005-01-06) the whole document	1-59
Y	WO 03/055860 A (RIBOTARGETS LIMITED; CANCER RESEARCH TECHNOLOGY LTD; THE INSTITUTE OF) 10 July 2003 (2003-07-10) the whole document ----- -/--	1-59

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

17 March 2006

Date of mailing of the international search report

20/04/2006

Name and mailing address of the ISA/
 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Lauro, P

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2005/041779

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>ROSTOM S A F ET AL: "Polysubstituted pyrazoles, part 5. Synthesis of new 1-(4-chlorophenyl)-4-hydroxy-1H-pyrazole-3-carboxylic acid hydrazide analogs and some derived ring systems.. A novel class of potential antitumor and anti-HCV agents"</p> <p>EUROPEAN JOURNAL OF MEDICINAL CHEMISTRY, EDITIONS SCIENTIFIQUE ELSEVIER, PARIS, FR, vol. 38, no. 11-12, November 2003 (2003-11), pages 959-974, XP004475841 ISSN: 0223-5234 examples 7a,7b</p>	1-59
Y	<p>B. W. DYMOCK ET AL.: "Inhibitors of HSP90 and other chaperones for the treatment of cancer"</p> <p>EXPERT OPIN. THER. PATENTS, vol. 14, no. 6, June 2004 (2004-06), pages 837-847, XP002372377 the whole document</p>	1-59
X	<p>L. LABANAUSKAS ET AL.: "Synthesis of 5-(2-,3- and 4-methoxyphenyl)-4H-1,2,4-triazole-3-thiol derivatives exhibiting anti-inflammatory activity"</p> <p>IL FARMACO, vol. 59, April 2004 (2004-04), pages 255-259, XP002372378 examples 4a,7a,8a,9a</p>	8,9,24
X	<p>US 5 489 598 A (CONNOR ET AL) 6 February 1996 (1996-02-06) examples 24,35,37</p>	8,9,24
X	<p>M. TANDON ET AL.: INDIAN JOURNAL OF CHEMISTRY, SECTION B: ORGANIC CHEMISTRY INCLUDING MEDICINAL CHEMISTRY, vol. 20b, no. 11, 1981, pages 1017-1018, XP009063370 table 1</p>	8,9,24
X	<p>DATABASE CAPLUS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; XP002372384 retrieved from STN Database accession no. 1989:8130 abstract</p> <p style="text-align: center;">-/--</p>	8,9,24

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2005/041779

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	<p>& GAWANDE ET AL.: "Synthesis of some thiosemicarbazides and related compounds" ACTA CIENCIA INDICA, CHEMISTRY, vol. 13, no. 2, 1987, pages 109-111, * see following compounds: RN=117839-88-0, 117839-89-1, 117839-90-4, 117839-91-5, 117839-92-6 *</p> <p>-----</p>	
X	<p>DATABASE BEILSTEIN [Online] Beilstein Institut zur Förderung der Chemischen Wissenschaften, Frankfurt am Main, DE; XP002372385 Database accession no. 625992 (BRN) abstract & SHAH ET AL.: INDIAN J. CHEM., vol. 5, 1967, pages 391-392,</p> <p>-----</p>	8,9,24
X	<p>DATABASE BEILSTEIN [Online] Beilstein Institut zur Förderung der Chemischen Wissenschaften, Frankfurt am Main, DE; XP002372386 Database accession no. 574001 (BRN) abstract & DOLESCHALL: ACTA CHIM. ACAD. SCI. HUNG., vol. 90, 1976, pages 419-422,</p> <p>-----</p>	8,9,24
P,X	<p>WO 2005/097758 A (AMPHORA DISCOVERY CORPORATION; HODGE, CARL, NICHOLAS; DICKSON, JOHN, K) 20 October 2005 (2005-10-20) the whole document</p> <p>-----</p>	1-59

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2005/041779

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 2005000300	A	06-01-2005	EP 1638555 A1	29-03-2006
WO 03055860	A	10-07-2003	AU 2002356301 A1	15-07-2003
			EP 1456180 A1	15-09-2004
			JP 2005517675 T	16-06-2005
			US 2005222230 A1	06-10-2005
US 5489598	A	06-02-1996	NONE	
WO 2005097758	A	20-10-2005	NONE	